

b
q=0.0028
q=0.0028
q=0.0028
q=0.038
ref.
Share of LFQ intensity (%)
100
75
50
25
0
Region 1
Region 2
Region 3
Region 4
Stool
Human-assigned
Microbial
Unresolved
c
Capsule–stool functional profile distance
0.8
0.7
0.6
0.5
0.4
0.3
Own stool
Other stools
d
Share of profile variation (%)
100
80
60
40
20
0
4.2%
10.7%
45.0%
5.3%
56.9%
Capsules + stool
Capsules only
Participant
Residual
Capsules vs stool
Capsule regions
e
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
Glyceraldehyde-3-phosphate dehydrogenase
Elongation factor Tu
Chaperonin GroEL
Starch-binding protein SusD
Pyruvate–ferredoxin oxidoreductase
Elongation factor G
Outer-membrane porin OmpA/OmpF
Pyruvate formate-lyase
Triosephosphate isomerase
Molecular chaperone DnaK
Glutamate dehydrogenase (NADP)
Phosphoglycerate kinase
Enolase
Glutamate dehydrogenase (NAD or NADP)
Formate–tetrahydrofolate ligase
Fucose/arabinose isomerase
Oligosaccharide-binding protein MsmE
Ribosomal protein S1
Ribosomal protein S3
Acetyl-CoA acetyltransferase
Oligosaccharide transporter permease
RNA polymerase subunit alpha
Cell division protease FtsH
Dihydroxy-acid dehydratase
1234S
0
1
2
Mean share of total LFQ (%)
B. wexlerae
F. saccharivorans
P. dorei
S. thermophilus
E. coli
Prevotella sp900557255
A. rectalis
Ruminococcus sp000437175
L. eligens
A. soehngenii
B. adolescentis
A. hallii
Other species
Not resolved or not sampled
f
Same leading species, capsule and own stool (%)
100
75
50
25
0
46.4
57.6
All eligible
Strict subset
g
−1
0
1
Native
eggNOG OG
KEGG KO
1 DNA
2 Protein
3 Host protein
4 Metabolites
1.00
0.51*
0.35*
0.55*
0.27*
0.38*
0.52*
0.67*
0.36*
0.43*
0.28*
0.70*
0.37*
0.49*
0.34*
—
1
2
3
4