
a
Peptides with a non-K/R C-terminus (%)
12
8
4
0
tryptic search
Duodenum
Jejunum
Ileum
Colon
Stool
De novo (top 30%): capsule > stool in 15/15, q < 0.001
Sage semi-tryptic: capsule > stool in 15/15, q < 0.001
Sage non-specific: capsule > stool in 15/15, q < 0.001
b
Histidine among non-K/R C-termini (%)
50
40
30
20
10
0
Duodenum
Jejunum
Ileum
Colon
Stool
Host: capsule > stool in 15/15, q < 0.001
Microbial: capsule > stool in 15/15, q < 0.001
c
Share of non-K/R C-termini (%)
Histidine
Proximal fluid (duodenum + jejunum)
Distal fluid (ileum + colon)
Stool
40
30
20
10
0
H
N
F
Y
M
A
G
Q
S
L
D
E
C
V
P
T
I
W
d
Duodenum
Jejunum
Ileum
Colon
Stool
PREDICT stool
0
10
20
30
40
Histidine at C-boundaries, observed – expected (points)
e
0
2
4
Histidine at N-boundaries, observed – expected (points)
f
0
2
4
Complementary products, histidine – control (points)
De novo (top 30%)
Sage semi-tryptic
Sage non-specific