

5–50 (published)
7–50, filtered
7–50, new search
9–50, filtered
9–50 subset of new search
a
Peptide gain over MG (%)
8 >250
7 >250
4 >250
12 >250
11 >250
10 >250
18.8
14.5
9.41
51.7
45.0
42.4
PREDICT (n = 296)
CapScan (n = 282)
b
MG identities recovered (%)
85.6
85.4
87.7
84.3
84.0
86.4
PREDICT (n = 296)
CapScan (n = 282)
c
PREDICT FL
PREDICT MG
CapScan FL
CapScan MG
Healthy FL
CAMPI FL
Median change in peptides vs 5–50 (%)
d
Member-specific peptides (pooled, +0.5)
FL
MG
Reference
L. plantarum
C. butyricum
B. longum
e
Pooled fusion diagnostic (%)
2.79
2.43
2.60
1.55
1.66
2.59
2.26
2.36
1.56
1.63
seed 20260907
seed 20260908
f
Median peptide change vs own FASTA (%)
7.1%
88.8% kept
9.3%
90.0% kept
32.9%
92.5% kept
34.6%
94.4% kept
Common FASTA
Joint search
g
Study groups, 7–50 vs 5–50 (%)
-0.1
+0.4
+0.4
-1.2
+0.0
+0.1
+0.6
-0.3
Healthy (58)
PREDICT (306)
CapScan (347)
CAMPI FL
CAMPI FL + Reference
CAMPI FL + Reference + MG
CAMPI Reference
CAMPI MG