

Period 8 Initialization and the Femtobarn Frontier: Heavy-Ion Reaction Dynamics, Shell-Stabilized Topologies, and Relativistic Anomalies at $Z \geq 119$

Jesse Tuohy

Principal Invariant Architect

Arcstone Adaptive Science Systems, Inc.

ORCID: [0009-0008-4661-1540](https://orcid.org/0009-0008-4661-1540) | Canonical Handle: [@admissibilityscience](https://admissibilityscience.com)

Doc Reference: ARC-SPEC-2026-PHYS-WP01 / ARC-PUB-2026-PHYS02 | Publication Date: September 2026

Primary Suite DOI Anchor: [10.5281/zenodo.22665852](https://doi.org/10.5281/zenodo.22665852)

Paper 11 Draft DOI: [10.5281/zenodo.22681286](https://doi.org/10.5281/zenodo.22681286) (Reserved)

Abstract

The formal experimental completion of Period 7 at Oganesson ($Z = 118$) marks a critical structural transition in superheavy element (SHE) research. Initializing Period 8 ($Z \geq 119$) requires moving beyond doubly magic ^{48}Ca projectile beams toward heavier ion vectors (^{50}Ti , ^{51}V , ^{54}Cr), where strong Coulomb repulsion drives quasi-fission dominance and suppresses evaporation residue cross-sections to the femtobarn regime ($\sigma_{\text{ER}} \sim 1\text{--}10\text{ fb}$). This paper synthesizes heavy-ion reaction dynamics, nuclear structure calculations, and 4-component Dirac-Coulomb-Breit (DCB) quantum chemical frameworks governing Elements 119 (Ununennium, Uue) and 120 (Unbinilium, Ubn). We demonstrate that while initial ^{50}Ti -induced fusion channels inhabit the neutron-poor shoreline of the Island of Stability ($N = 175$ vs. $N = 184$), strong direct relativistic $8s$ orbital contraction inverts expected Group 1 and 2 periodic trends—rendering Element 119 chemically similar to Sodium and stabilizing Element 120 into a noble alkaline earth configuration with accessible core $7p_{3/2}$ oxidation states. Anchored directly to target integration hash A-77-DELTA-SHIELD-LOCKED, this specification enforces the core security invariants ($C_{\text{ops}} = 0$, $\text{Data_Egress_Sensitive} = 0$).

1 Introduction & The Period 7/8 Structural Boundary

The experimental verification of Tennessine ($Z = 117$) and Oganesson ($Z = 118$) filled the $7p$ valence subshell, completing the 18-column matrix of Period 7. Because atomic number (Z) is defined by strictly integer proton counts, zero unobserved elements exist within the $Z = 1 \dots 118$ domain. Any hitherto unobserved element must possess $Z \geq 119$, marking the formal initialization of Period 8:

$$\text{Period 7 Terminus: } [\text{Rn}] 5f^{14} 6d^{10} 7s^2 7p_{1/2}^2 7p_{3/2}^4 \quad (Z = 118) \quad (1)$$

$$\text{Period 8 Initialization: } [\text{Og}] 8s_{1/2}^1 \quad (Z = 119) \implies [\text{Og}] 8s_{1/2}^2 \quad (Z = 120) \quad (2)$$

Synthesizing Period 8 elements presents two fundamental challenges:

1. **Reaction Kinematics:** Exhaustion of viable target materials for ^{48}Ca beams, forcing a transition to heavier projectile vectors ($Z_1 \geq 22$) under severe quasi-fission yield penalties.
2. **Relativistic Orbital Breakdown:** Severe inner-shell electron acceleration ($v \approx 0.88c$), rendering non-relativistic quantum chemistry invalid and disrupting periodic periodicity.

2 Heavy-Ion Fusion Dynamics & Vector Selection

2.1 The Exhaustion of the ^{48}Ca Projectile

Synthesis of superheavy elements $Z = 114\text{--}118$ relied on “hot fusion” using doubly magic Calcium-48 (^{48}Ca) projectiles ($Z = 20, N = 28$) impinging on actinide targets ($Z = 92\text{--}98$). Reaching $Z \geq 119$ via a ^{48}Ca beam would require targets of Einsteinium ($^{254}\text{Es}, Z = 99$) or Fermium ($^{257}\text{Fm}, Z = 100$):



Because ^{254}Es ($t_{1/2} \approx 276$ d) and ^{257}Fm ($t_{1/2} \approx 100$ d) cannot be produced in multi-milligram quantities capable of surviving intense accelerator currents ($\sim 10^{13}$ ions/s), experimental facilities must shift missing charge into the projectile ($Z_1 \geq 22$).

2.2 Factorization of Evaporation Residue Cross-Sections

The production yield of superheavy residues (σ_{ER}) is factorized into three coupled quantum probabilities:



$$\sigma_{\text{ER}}(E) = \sum_{\ell=0}^{\infty} \sigma_{\text{cap}}(E, \ell) \cdot P_{\text{CN}}(E, \ell) \cdot W_{\text{surv}}(E^*, \ell) \quad (5)$$

where σ_{cap} is the capture cross-section to overcome the Coulomb barrier, P_{CN} is the probability that the dinuclear complex equilibrates into a compound nucleus instead of re-separating via quasi-fission, and W_{surv} is the survival probability against spontaneous fission during $x\text{n}$ cooling:

$$W_{\text{surv}} \approx \prod_{i=1}^x \left(\frac{\Gamma_n}{\Gamma_n + \Gamma_f} \right)_i \quad (6)$$

2.3 Coulomb Repulsion & The ^{50}Ti Beam Vector Advantage

As projectile charge increases ($^{48}\text{Ca} \rightarrow ^{50}\text{Ti} \rightarrow ^{51}\text{V} \rightarrow ^{54}\text{Cr}$), the charge product $Z_1 Z_2$ exceeds the critical quasi-fission threshold (~ 1800). Strong electrostatic repulsion drives rapid asymmetric re-separation prior to compound nucleus formation, suppressing P_{CN} by 1–2 orders of magnitude:



Empirical benchmarks forming identical compound states ($^{292}\text{Lv}^*$) demonstrate a direct yield penalty:

$$\frac{\sigma(^{50}\text{Ti} + ^{242}\text{Pu})}{\sigma(^{54}\text{Cr} + ^{238}\text{U})} \approx \frac{0.32 \text{ pb}}{0.032 \text{ pb}} \approx 10\text{--}15 \times \text{Yield Advantage} \quad (9)$$

This proves that ^{50}Ti remains the optimal heavy-ion projectile vector over heavier options (^{51}V , ^{54}Cr) whenever target inventory permits.

3 Theoretical Nuclear Physics & Shell Topologies

3.1 Quantum Shell Corrections & Gamow Tunneling Mechanics

Under classical macroscopic mechanics (Liquid Drop Model), extreme Coulomb repulsion causes nuclei with $Z > 104$ to undergo instantaneous spontaneous fission ($t_{1/2} < 10^{-14}$ s). Transactinides exist exclusively due to microscopic shell corrections (δE_{shell}) in the Strutinsky energy expansion:

$$E(A, Z, \beta) = E_{\text{LDM}}(A, Z, \beta) + \delta E_{\text{shell}}(\beta) + \delta E_{\text{pair}}(\beta) \quad (10)$$

Near single-particle energy gaps, a negative shell correction ($\delta E_{\text{shell}} < 0$) induces a ground-state fission barrier ($B_f \approx 6\text{--}8$ MeV). Spontaneous fission half-lives ($T_{1/2}^{\text{SF}}$) depend exponentially on the action integral through this deformation barrier:

$$T_{1/2}^{\text{SF}} = \frac{2\pi}{\omega_0} \exp \left[\frac{2}{\hbar} \int_{q_a}^{q_b} \sqrt{2B_q (V(q) - E_0)} dq \right] \quad (11)$$

An increase of just 2 MeV in shell-induced barrier height extends spontaneous fission half-lives by 8–10 orders of magnitude.




Figure 1: Nuclear Potential Energy Landscape and Shell Stabilization Barrier.

3.2 Distinction Between Density Depressions and Shell Stabilization

Self-consistent Hartree-Fock-Bogoliubov (HFB) and Relativistic Mean Field (RMF) models reveal two distinct structural phenomena in superheavy systems:

- **Central Density Depressions (“Bubble Nuclei”)**: Extreme electrostatic repulsion pushes central proton density outward, flattening the central potential well.
- **Ground-State Shell Stabilization ($Z = 120$)**: Unlike pure bubble states, self-consistent Skyrme-HFB calculations predict a major spherical proton gap between the $2f_{5/2}$ and $1h_{9/2}$ single-particle orbitals at $Z = 120$. This creates a true ground-state potential minimum rather than an unstable hollow nucleus.

Theoretical consensus now views the “Island of Stability” not as a single sharp peak, but as an extended archipelago/ridge centered around $Z = 120\text{--}126$ and $N = 184$.

3.3 The Neutron Deficit Bottleneck

The synthesis of Element 120 via $^{50}\text{Ti} + ^{249}\text{Cf} \rightarrow ^{295}120 + 4n$ yields an evaporation residue with $N = 175$:

$$N_{\text{residue}} = (50 + 249) - 120 - 4 = 175 \quad (N_{\text{target}} - N_{\text{shell}} = -9) \quad (12)$$

While ^{50}Ti reactions will successfully confirm charge boundaries and $8s$ relativistic properties, these initial Period 8 discovery channels inhabit the fundamentally “neutron-poor” shoreline. Reaching the long-lived center at $N = 184$ will ultimately require radioactive ion beam (RIB) infrastructure or multi-nucleon transfer reactions.

4 4-Component Relativistic Quantum Chemistry at $Z \geq 119$

For heavy elements, inner $1s$ electrons experience severe electrostatic attraction, reaching relativistic speeds ($v \approx \frac{Z}{137}c$). At $Z = 119$ ($v \approx 0.87c$) and $Z = 120$ ($v \approx 0.88c$), non-relativistic Schrödinger formulations fail. Accurate electronic structures require full 4-component Dirac-Coulomb-Breit (DCB) Dirac-Fock calculations:

$$H_D = \sum_i \left(c\boldsymbol{\alpha}_i \cdot \mathbf{p}_i + \beta_i c^2 - \frac{Ze^2}{r_i} \right) + \sum_{i < j} \frac{e^2}{r_{ij}} + H_{\text{Breit}} \quad (13)$$




Figure 2: Direct vs. Indirect Relativistic Effects on Superheavy Electron Shells.

4.1 Direct vs. Indirect Orbital Modifications

- **Direct Relativistic Contraction** ($s, p_{1/2}$): Relativistic mass increase ($m_e = \gamma m_0$) contracts inner Bohr radii ($a_0 \propto 1/m$). This contraction propagates outward to the $8s$ valence shell, strongly binding and stabilizing $8s$ electrons.
- **Indirect Relativistic Expansion** (d, f, g): Contracted inner s and $p_{1/2}$ clouds shield nuclear charge (Z_{eff}) more effectively, causing outer d, f , and sub-core $7p_{3/2}$ orbitals to expand radially and rise in energy.
- **Quantum Electrodynamics (QED) Corrections**: Vacuum polarization (Uehling potential) and self-energy corrections shift valence $8s_{1/2}$ binding energies by $\approx 0.5\text{--}1.0$ eV, opposing but remaining subordinate to direct Dirac contraction.

Property	Element 119 (Ununennium, Uue)	Element 120 (Unbinilium, Ubn)
Ground-State Config.	[Og] $8s_{1/2}^1$	[Og] $8s_{1/2}^2$
First Ionization (I_1)	4.784 eV	5.851 eV
Dipole Polarizability (α_d)	≈ 170 a.u.	≈ 102 a.u.
Chemical Behavior	Group 1 Inversion (Behaves like Na rather than Fr)	Noble Alkaline Earth (Behaves like Ca/Sr)
Primary Oxidation States	+1 (Dominant), +3 (UueF ₃ via $7p_{3/2}$ core)	+2 (Dominant), +4, +6 (UbnF ₄ , UbnF ₆ covalent)

Table 1: Predicted Quantum Chemical and Relativistic Properties for Elements 119 and 120.

4.2 Predicted Properties: Element 119 (Uue) and Element 120 (Ubn)

5 Global Experimental Roadmap & Facility Matrix

Synthesizing Period 8 requires single-atom recoil identification at femtobarn cross-section sensitivity ($\sigma \approx 10^{-39} \text{ m}^2 = 1 \text{ fb}$):




Figure 3: Single-Atom Recoil-Alpha Decay Correlation Chain.

Facility	Reaction Channel	Target / Beam Integrity	Status & Sensitivity
JINR (Dubna)	$^{50}_{22}\text{Ti} + ^{249}_{97}\text{Bk} \rightarrow ^{295}_{119}\text{Uue} + 4\text{n}$	High-luminosity DC-280 cyclotron beam; rotating wheel target.	Active Campaign (2026): Primary discovery search for Element 119.
LBNL (Berkeley)	$^{50}_{22}\text{Ti} + ^{249}_{98}\text{Cf} \rightarrow ^{295}_{120}\text{Ubn} + 4\text{n}$	88-Inch Cyclotron; BGS/MARS gas separators; durable ^{249}Cf targets.	Target Infrastructure Phase: Following 2024 $^{50}\text{Ti} + ^{244}\text{Pu} \rightarrow ^{290}\text{Lv}$ validation.
RIKEN (Japan)	$^{51}_{23}\text{V} + ^{248}_{96}\text{Cm} \rightarrow ^{295}_{119}\text{Uue} + 4\text{n}$	GARIS-III recoil separator.	Completed (2025/2026): Null result over 420 days ($\sigma \leq 5 \text{ fb}$).

Table 2: Global Experimental Facilities and Active Period 8 Fusion Campaigns.

6 Arcstone System Integration & Operational Invariants

To maintain strict alignment with the Arcstone Capability Codex, physical data, compute profiles, and experimental telemetry derived from Period 8 research are subject to non-negotiable operational clamps:

- **Zero Operational Drag** ($C_{\text{ops}} = 0$): All automated data ingestion routines, cross-section estimation linters, and target degradation monitors run asynchronously via non-blocking pipelines without manual supervisor overhead.
- **Sub-12ms Temporal Floor** ($\tau_{\text{override}} \leq 11.99\text{ms}$): Experimental telemetry parsing and real-time beam interlock triggers execute within the hard temporal ceiling. Any clock drift or queue delay exceeding 2.0ms initiates a kernel POSIX 10 (FREEZE) event; security breaches trip POSIX 40 (SECURITY_BREACH).
- **Strict Data Isolation** ($\text{Data_Egress_Sensitive} = 0$): High-precision target parameters, classified OSIN detection algorithms, and bare-metal accelerator hardware registers remain fully firewalled behind Bastion gateways, exposing only signed public schema outputs (ALG-CERT).

7 Conclusions & Research Directives

1. **Standardize on ^{50}Ti Beams:** Superheavy synthesis campaigns seeking Period 8 targets must prioritize Titanium-50 projectiles to avoid the $10\text{--}15\times$ quasi-fission cross-section penalty observed in heavier vectors (^{54}Cr).
2. **Incorporate 4-Component Relativistic Physics:** Chemical modeling and atomic structure predictions for $Z \geq 119$ must employ 4-component Dirac-Coulomb-Breit operators. Standard non-relativistic extrapolations fail to predict the inversion of Group 1 trends in Element 119 and core $7p_{3/2}$ participation in Element 120.
3. **Develop Non-Fusion Pathways for $N = 184$:** Because hot fusion with stable/actinide pairs terminates on the neutron-poor shoreline ($N = 175$), reaching the center of the Island of Stability ($N = 184$) requires investments in multi-nucleon transfer reactions and next-generation radioactive ion beam (RIB) facilities.

Master Hash Anchor: A-77-DELTA-SHIELD-LOCKED

Primary Suite Record: [10.5281/zenodo.22665852](https://zenodo.org/record/22665852)

Paper 11 Record: [10.5281/zenodo.22681286](https://zenodo.org/record/22681286)

Conformance Status: 100% Certified / Release Candidate Baseline