

# CHAPTER 6: CONTINUOUS PROBABILITY DISTRIBUTIONS

Uniform, Normal, and Exponential Models

## I. Continuous Probability Basics

A continuous random variable can take on any value in an interval. Because there are infinite possible values, we use a **Probability Density Function (PDF)**, denoted as  $f(x)$ .

- The total area under the PDF curve is always **1**.
- Probability is defined as the area under the curve between two points  $a$  and  $b$ .

## II. The Uniform Distribution

This is the simplest distribution, used when the probability is "evenly spread" over an interval from  $a$  to  $b$ . The "curve" is actually a flat rectangular line.

$$f(x) = \frac{1}{b-a} \quad \text{for } a \leq x \leq b$$

**Calculation:** To find  $P(x_1 < x < x_2)$ , simply calculate the area of the rectangle:

$$\text{Area} = \text{Base} \times \text{Height} = (x_2 - x_1) \times \frac{1}{b-a}$$

## III. The Normal Distribution

The most important distribution in all of statistics. It is symmetrical and bell-shaped, defined by its mean ( $\mu$ ) and standard deviation ( $\sigma$ ).

### 1. Characteristics

- The highest point is at the mean, which is also the median and mode.
- The curve extends to infinity in both directions but never touches the x-axis.

### 2. The Standard Normal ( $Z$ )

To find probabilities, we convert any normal variable  $x$  into a **Standard Normal Variable ( $z$ )** with  $\mu = 0$  and  $\sigma = 1$ :

$$z = \frac{x - \mu}{\sigma}$$

We then use the **Z-Table** to find the cumulative probability  $P(Z \leq z)$ .

### 3. Using the Z-Table

- **Left-tail:**  $P(X < a) \rightarrow$  Look up  $z$  directly.
- **Right-tail:**  $P(X > a) \rightarrow 1 - P(Z < z)$ .
- **Between:**  $P(a < X < b) \rightarrow P(Z < z_b) - P(Z < z_a)$ .

## IV. The Exponential Distribution

Often used to model the **time or distance between arrivals** (e.g., time between customers entering a bank).

$$f(x) = \frac{1}{\mu} e^{-x/\mu}$$

**Cumulative Probability Function:**

$$P(X \leq x) = 1 - e^{-x/\mu}$$

Where  $\mu$  is the mean time between events.

## V. Teacher's Strategy: Visualizing the Area

Students often struggle with Z-tables because they forget which "side" they are looking at.

- **Always sketch the curve.** Shade the area you are looking for.
- If your  $z$  value is positive, the area to the left must be greater than 0.50.
- If you are looking for a "between" probability, you are essentially subtracting a small tail from a larger tail.

## VI. Step-by-Step Example

**Problem:** Monthly utility bills are normally distributed with  $\mu = 100$  and  $\sigma = 20$ . What is the probability a bill is *more* than \$130?

**Logic:** 1. **Standardize:**  $z = (130 - 100)/20 = 30/20 = 1.5$ . 2. **Table Look-up:** Find  $z = 1.50$  in the Z-table. The area to the left is 0.9332. 3. **Adjust for Tail:** Since we want "more than," we need the area to the *right*. 4. **Result:**  $1 - 0.9332 = 0.0668$  (or 6.68%).

## VII. Practice Set

1. For a uniform distribution from  $a = 10$  to  $b = 50$ , what is the height of the density function? What is  $P(x < 20)$ ?
2. A process has a mean of 500 and std dev of 50. Find the z-score for  $x = 425$ .
3. The time between arrivals at a car wash follows an exponential distribution with a mean of 10 minutes ( $\mu = 10$ ). What is the probability the next car arrives within 5 minutes?
4. If  $P(Z < z) = 0.9750$ , what is the value of  $z$ ? (Work backwards using the table).

## VIII. Answer Key

1. **Height:**  $1/(50-10) = 0.025$ . **Prob:**  $(20-10) \times 0.025 = 0.25$ .
2.  $z = (425 - 500)/50 = -75/50 = -1.5$ .
3.  $P(X \leq 5) = 1 - e^{-5/10} = 1 - e^{-0.5} \approx 0.3935$ .
4.  $z = 1.96$ .