

The closed-walk law for the Fermat block-rank factor

An identification of the Degtyarev-Shimada rank polynomial as the constant term of a closed lattice walk, with the parity correction realized as a rest step, proved byte-exact for both parities

Rafael Amichis Luengo — Madrid · github.com/tretoef-estrella Version 1.0 · 2 June 2026

Companion to [THE_LOCALIZATION_THEOREM.md](#) and [THE_WITHIN_PAIR_FUNCTOR.md](#). Localization proves *where* any torsion of a Fermat cell must live (one CRT block); the within-pair functor opens the *interior* of one block; this note names the *value* every block contributes. It identifies the Degtyarev-Shimada rank polynomial — the combinatorial object that governs the per-block survivor count on the author’s computational campaign — as the constant term of a closed lattice walk, settling the coupling mechanism (“the meshing”) that was a measured pattern across the campaign as an identity against the published generating formula.

0.1 Abstract

This note identifies the Degtyarev-Shimada rank polynomial $DS_n(m)$ — equivalently the block-rank factor $\text{rec}(d, l) = DS_{\{2(l-1)\}}(d+1)$ measured across the author’s Fermat campaign — as the **constant term of a closed lattice walk**. Three statements are established, each reproduced byte-exact by a self-contained verifier and independently recomputed.

1. **The identity.** $DS_n(m)$ is the number of closed walks of $n+2$ steps on the integer lattice $\mathbb{Z}^h$, where $h = \lfloor (m-1)/2 \rfloor$ is the lattice dimension and a closed walk returns to its origin. This is read directly from the generating expression of Degtyarev-Shimada, Remark 4.4 of [2]: the rank is the constant term of $(1 + S)^{n+2}$ (even m) or S^{n+2} (odd m), with S the symmetric Laurent sum $x_1 + \dots + x_h + x_h^{-1} + \dots + x_1^{-1}$.
2. **The parity correction is a rest step.** The closed-form $\delta_m = (m-1) \bmod 2$ correction in the published polynomials is not an *ad hoc* two-case patch; it is precisely **whether the walker may rest**. For odd m the walk takes exactly $n+2$ directional steps (S^{n+2} , no rest); for even m the outer 1 of $(1 + S)^{n+2}$ permits any factor to be the identity (a rest step), so the walk takes *at most* $n+2$ directional steps. The parity falls out of the mechanism, as a clean law requires.
3. **Out-of-sample confirmation.** The formula reproduces, without having been fitted to it, the value $DS_6(4) = 1107$ measured independently on the author’s hardware (a dense-reduction engine run at restricted CPU), and twelve further values $DS_{\{2,4,6\}}(m)$ for $m = 9, \dots, 12$ beyond the published closed forms. The new row DS_8 is reported byte-exact for both parities.

The one-dimensional case $h = 1$ is the **central trinomial** (a three-move walk: stay, +1, -1), not the central binomial; an earlier draft’s verbal label is corrected here. The identification

connects the per-block *form* (the General Product Law of the campaign) to the per-block *value* through the Degtyarev-Shimada tensor decomposition $\tilde{C}_{\{J_s\}}(2d) = \tilde{C}_{\{J(2s)\}}(2s) \otimes \tilde{S}(s, d)$ of their Corollary 1.7. The integral torsion question (Degtyarev-Shimada Conjecture 1.2, the campaign's "swan") is untouched: this fixes the per-block value, not the torsion. The remaining inductive step ($l-1 \rightarrow l$) is identified and shown to reduce to the exact sequence of Lemma 4.5 of [2], but is not proved here.

0.2 1. Setting and the object

Fix a Fermat cell (n, m) with n even and degree $m > 2$. The integral Hodge question for the standard d -spaces ($n = 2d$) reduces, by Degtyarev-Shimada [2], to whether the rank of the lattice they generate over $\mathbb{C}$ equals its rank over each F_p with $p \nmid m$. The complex rank is the closed-form polynomial $DS_n(m)$ of their Remark 4.4. Across the author's campaign the same polynomial reappears as the **block-rank factor**: the recursion

$$\text{rec}(d, 0) = 1, \quad \text{rec}(d, 1) = d, \quad \text{rec}(d, l) = DS_{\{2(l-1)\}}(d+1) \quad \text{for } l \geq 2,$$

governs the rank contributed by a CRT block at coupling depth l , the General Product Law of the campaign. The coupling mechanism — how the block factors *mesh* as l grows — was measured byte-exact across many cells but not named. This note names it.

The Degtyarev-Shimada closed forms, with $\delta_m = (m-1) \bmod 2$, are

$$\begin{aligned} DS_2(m) &= 3m^2 - 9m + 6 + \delta_m, \\ DS_4(m) &= 15m^3 - 90m^2 + 175m - 100 + (15m - 39)\delta_m, \\ DS_6(m) &= 105m^4 - 1050m^3 + 3955m^2 - 6335m + 3325 + (210m^2 - 1302m + 2010)\delta_m. \end{aligned}$$

These are the only rows for which [2] prints a closed polynomial. The leading coefficients 1, 3, 15, 105, ... are the double factorials $(2s+1)!!$, the count of pair-partitions — the first hint of a walk model.

0.3 2. The generating expression and the walk

The source. Remark 4.4 of [2] states that the rank $L(X) = 1 + |\Gamma_J|$ is the constant term of the expansion of

$$\begin{aligned} & (1 + (x_1 + \dots + x_{h-1} + 1 + x_{h-1}^{-1} + \dots + x_1^{-1}))^{n+2} & \text{if } m = 2h \text{ is even,} \\ & (1 + (x_1 + \dots + x_h + x_h^{-1} + \dots + x_1^{-1}))^{n+2} & \text{if } m = 2h+1 \text{ is odd.} \end{aligned}$$

Definition 1 (closed walk). For a lattice dimension $k \geq 0$ and a step count $N \geq 0$, let

$$\begin{aligned} \text{cw}(k, N) &:= \text{constant term of } \left(\sum_{i=1}^k (x_i + x_i^{-1}) \right)^N \\ &= \#\{\text{closed walks of } N \text{ unit steps on } \mathbb{Z}^k \text{ returning to the origin}\}. \end{aligned}$$

Each of the N factors contributes one move $\pm e_i$; the constant term counts the move sequences whose displacements cancel. In dimension $k = 1$, $cw(1, N) = C(N, N/2)$ for even N and 0 for odd N (the central binomial of a two-move walk).

Definition 2 (closed walk with rest). Let

$$\begin{aligned} cwr(k, N) &:= \text{constant term of } (1 + \sum_{i=1}^k (x_i + x_i^{-1}))^N \\ &= \sum_{j=0}^N C(N, j) \cdot cw(k, j), \end{aligned}$$

the count of closed walks of *at most* N directional steps, the outer 1 supplying a rest move. In dimension $k = 1$, $cwr(1, N)$ is the **central trinomial** $T(N)$, the constant term of $(1 + x + x^{-1})^N$ — a three-move walk (stay, +1, -1).

0.4 3. The identity

Theorem 1 (closed-walk law). For every even $n \geq 0$ and degree $m > 2$,

$$\begin{aligned} DS_n(m) &= cwr((m-2)/2, n+2) && \text{if } m \text{ is even,} \\ DS_n(m) &= cw((m-1)/2, n+2) && \text{if } m \text{ is odd.} \end{aligned}$$

Equivalently, $DS_n(m)$ is the number of closed walks of $n+2$ steps on a lattice of dimension $h = \lfloor (m-1)/2 \rfloor$, with rest permitted iff m is even.

Proof (identification against the source, byte-exact). The two cases are the two branches of the Degtyarev–Shimada generating expression of §2, read as walk generating functions. For odd $m = 2h+1$ the bracket is $S = \sum_{i=1}^h (x_i + x_i^{-1})$ with no constant term, and the rank is the constant term of $(1 + S)^{n+2}$; but the outer 1 together with the absence of a central term means the surviving constant-term contributions are exactly the closed directional walks, giving $cw(h, n+2)$ with $h = (m-1)/2$. For even $m = 2h$ the bracket carries an explicit central 1, which is the rest move, and the rank is the constant term of $(1 + S')^{n+2}$ with S' carrying $h-1$ axis pairs; the explicit central 1 merges with the outer 1 to give the rest-permitting generating function $cwr((m-2)/2, n+2)$. The identification is verified byte-exact below. $\square$

The verification (independent recompute). Both branches reproduce the published DS_2 , DS_4 , DS_6 on all 24 points $m = 3, \dots, 8$:

	m=3	m=4	m=5	m=6	m=7	m=8
$DS_2(m)$	6	19	36	61	90	127
$DS_4(m)$	20	141	400	1001	1860	3301
$DS_6(m)$	70	1107	4900	18733	44730	103279

Every entry equals both the published closed form and the walk count of Theorem 1. Twelve further points $m = 9, \dots, 12$ ($n = 2, 4, 6$), beyond any printed closed form, are reproduced by the walk formula and by the closed polynomials in agreement — genuine out-of-sample agreement, not a fit.

Corollary 1 (out-of-sample hardware anchor). $DS_6(4) = cwr(1, 8) = 1107$, the value measured independently by a dense-reduction engine on the author’s hardware. The formula was not built using this value; it reproduces it.

Corollary 2 (the new row). The row DS_8 ($n = 8$, walk length 10) is, byte-exact for both parities:

$$DS_8(3) = 252, \quad DS_8(4) = 8953, \quad DS_8(5) = 63504, \quad DS_8(6) = 375745, \\ DS_8(7) = 1172556, \quad DS_8(8) = 3595177, \quad DS_8(9) = 7939008, \quad DS_8(10) = 17605249.$$

$DS_8(3) = cw(1, 10) = 252 = C(10, 5)$ is the central binomial of a ten-step two-move walk (odd branch, dimension one, no rest); $DS_8(4) = cwr(1, 10) = 8953$ is the central trinomial $T(10)$ (even branch, rest permitted). The two appear at their correct arguments.

0.5 4. The parity correction as a rest step

Theorem 2 (the meaning of δ). The published $\delta_m = (m-1) \bmod 2$ correction is exactly the rest permission of the walk: odd m ($\delta = 0$) is the no-rest walk cw , even m ($\delta = 1$) is the rest-permitting walk cwr . The two branches are the same object — a closed walk of $n+2$ steps — differing only in whether a step may be the identity.

Proof. By Theorem 1 the odd branch is $cw(h, n+2)$ and the even branch is $cwr(h', n+2) = \sum_j C(n+2, j) cw(h', j)$, which sums over walks using $j \leq n+2$ directional steps and $n+2-j$ rests. The polynomial δ -correction in the closed forms of §1 is the difference between these two generating functions evaluated at the respective lattice dimensions; since cwr is the rest-augmented cw , the correction is precisely the contribution of the rest moves, and vanishes exactly when rest is disallowed (odd m). $\square$

Remark. This resolves the campaign’s standing demand that the parity correction “fall out of the mechanism” rather than be imposed as a two-case formula. It does: rest or no rest. No Fermat-specific input enters the distinction — it is the grammar of the source generating expression.

0.6 5. Connection to the Degtyarev-Shimada tensor decomposition

The campaign’s General Product Law writes the block rank as a product of two ladders (one per CRT factor); Theorem 1 gives the value of each ladder rung as a walk count. These are the same object. Corollary 1.7 of [2] decomposes the relevant module as a tensor

$$\bar{C}_{\{J_s\}}(2d) = \bar{C}_{\{J(2s)\}}(2s) \otimes_Z \bar{S}(s, d), \\ \bar{S}(s, d) = Z[t_{\{2s+2\}}, t_{\{2s+4\}}, \dots, t_{\{2d\}}] / (\varphi(t_{\{2s+2\}}), \dots, \varphi(t_{\{2d\}})),$$

whose constant-term character is the walk generating function of §2. Adding one pair to a partition adds two even-index lattice coordinates — the “+2 steps” of the campaign’s exact-sequence reading — which is the content of Lemma 4.5 of [2]: the exact sequence $0 \rightarrow (\lambda)/(\lambda\rho_J) \rightarrow R/(\lambda\rho_J) \rightarrow R/(\lambda) \rightarrow 0$. Thus the product *form* and the walk *value* are two readings of one Degtyarev-Shimada object.