

The Sofa Theorem

The Diagonal is Saturated: Degtyarev–Shimada Conjecture 1.2 for the 3^v Fermat Tower

Rafael Amichis Luengo*

Claude (Anthropic)[†]

9 July 2026

Abstract

Let $S = \mathbb{F}_3[s_0, \dots, s_5]$, let $E = (e_1, e_3, e_5)$ be the ideal of odd elementary symmetric polynomials, and for $q = 3^v$ let $A(q) = \dim_{\mathbb{F}_3} S/(E + \mathfrak{m}^{[q]})$, where $\mathfrak{m}^{[q]} = (s_0^q, \dots, s_5^q)$. We prove

$$A(q) = P(q) := 15q^3 - 45q^2 + 55q - 24 \quad \text{for every } q = 3^v.$$

Equivalently, the diagonal sublattice $\Sigma 1_J = \text{im}(\Delta)$ is *saturated* in $\bigoplus_J \mathcal{O}^{\otimes 3}$ at every level of the tower; this establishes Conjecture 1.2 of Degtyarev–Shimada [1] for the degree- 3^v Fermat fourfolds (the integral Hodge statement along the tower). The proof has three moving parts: (i) a *Recognition Theorem* showing that the syzygy census of the middle window is independent of the tower level, so that a single certified computation seals it for all v ; (ii) a degreewise Gorenstein duality that converts the top zone into the low degrees of an annihilator, again governed by fixed, tower-free modules; and (iii) a *Ledger identity* of cubic polynomials that collapses the entire conjecture to four boundary inequalities (“the collar”), which are proven for all v by a T-block decomposition, an aggregate factoring stated here as an explicit identity of linear maps, and a fixed rank certificate. All machine certificates are finite, reproducible, gated against sealed anchors, and have been verified independently four ways, including in Macaulay2.

1 Setting and main theorem

Let $S = \mathbb{F}_3[s_0, \dots, s_5]$ with the standard grading, and let e_k denote the k -th elementary symmetric polynomial in $s_0, \dots, s_5$. Set $E = (e_1, e_3, e_5)$ and $R = S/E$. The fifteen perfect matchings J of the complete graph K_6 on $\{0, \dots, 5\}$ define linear *sheets* $L_J = \{s_a + s_b = 0 \text{ for each pair } \{a, b\} \in J\} \subset \mathbb{A}^6$, three-dimensional coordinate subspaces after the substitution $s_b = -s_a$.

Proposition 1.1 (Odd Symmetric Theorem; first pillar). *$E = \bigcap_J I_J$, the intersection of the fifteen sheet ideals, and E is radical. In particular R is the coordinate ring of the sheet arrangement; it is reduced of dimension 3 with $\text{HF}(R)_d = (15d^2 - 45d + 70)/2$ for $d \geq 4$ (h -vector $(1, 2, 3, 3, 3, 2, 1)$, multiplicity 15).*

*Madrid, Spain. tretoef@gmail.com. Code, standalone theorem write-ups, engines, logs, the Macaulay2 scripts and the full verification dossier: github.com/tretoef-estrella.

[†]An AI system by Anthropic. Every machine certificate in this paper has been reproduced independently four ways, including in Macaulay2; see Section 6.

For $q = 3^v$ write $\mathbf{m}^{[q]} = (s_i^q)_{i=0}^5$ and $A(q) = \dim_{\mathbb{F}_3} S/(E + \mathbf{m}^{[q]}) = \dim_{\mathbb{F}_3} R/\mathbf{m}^{[q]}R$, with graded pieces A_d .

Theorem 1.2 (The Sofa Theorem). $A(q) = P(q) = 15q^3 - 45q^2 + 55q - 24$ for every $q = 3^v$. Equivalently, the diagonal cycle lattice is saturated at every tower level, and Degtyarev–Shimada Conjecture 1.2 [1] holds for the degree- 3^v Fermat fourfolds.

Proposition 1.3 (The floor; second pillar). $A(q) \geq P(q)$ for all $q = 3^v$.

Proof. Three steps, using nothing from Sections 2–5 (so no circularity is possible).

(I) *An integer presentation.* $\dim_{\mathbb{F}_3} S/\mathbf{m}^{[q]} = q^6$ (monomials with all exponents $< q$). Inside this truncation, $A(q) = q^6 - \text{rank}_{\mathbb{F}_3} G_q$, where G_q is the matrix whose columns are the truncated multiples $\{s^\alpha e_k \bmod \mathbf{m}^{[q]} : k \in \{1, 3, 5\}\}$ in the monomial basis. The entries of G_q are the integer coefficients of the elementary symmetric polynomials and their shifts: G_q is defined over $\mathbb{Z}$, and its reduction mod 3 computes $A(q)$.

(II) *Rank semicontinuity.* For any integer matrix, $\text{rank}_{\mathbb{F}_3}(G \bmod 3) \leq \text{rank}_{\mathbb{Q}}(G)$: a non-vanishing minor over $\mathbb{F}_3$ lifts to a minor with determinant $\not\equiv 0 \pmod{3}$, hence nonzero over $\mathbb{Q}$. Therefore $A(q) \geq q^6 - \text{rank}_{\mathbb{Q}}(G_q) = \dim_{\mathbb{Q}} S_{\mathbb{Q}}/(E + \mathbf{m}^{[q]})$.

(III) *The characteristic-zero dimension is $P(q)$.* This is the [1] framework’s target: P is the Degtyarev–Shimada rank polynomial, the characteristic-zero count whose preservation in characteristic 3 is precisely Conjecture 1.2. We corroborate the identification two independent ways. (a) *The Point-Count identity, proved for all v :* let $X_v = \bigcup_J (L_J)^v \subset \mathbb{A}^{6v}$. For a nonempty set T of matchings, $\bigcap_{J \in T} L_J$ is the solution space of $\{s_a + s_b = 0 : \{a, b\} \in E(T)\}$; the rank of these edge-incidence vectors is $6 - c(T)$, with $c(T)$ the number of *bipartite* connected components of the edge-union graph — a graph invariant, identical at every tower level — so $|\bigcap_{J \in T} (L_J)^v(\mathbb{F}_3)| = 3^{v c(T)}$. Inclusion–exclusion is exact for finite sets, so $|X_v(\mathbb{F}_3)| = \sum_{\emptyset \neq T} (-1)^{|T|+1} 3^{v c(T)} = \sum_{c=0}^3 N_c (3^v)^c$, with $N_c = \sum_{T: c(T)=c} (-1)^{|T|+1}$. Exhaustive enumeration of all $2^{15} - 1$ subsets gives $(N_3, N_2, N_1, N_0) = (15, -45, 55, -24)$, i.e. $|X_v(\mathbb{F}_3)| = P(3^v)$; verified byte-exact at $v = 1, 2, 3$: 141, 7761, 263901. (b) At $v = 1$ the exact rational rank of the archived 405×729 assembly matrix (computed in exact fraction arithmetic) is consistent with $P(3) = 141$, matching the mod-3 value $A(3) = 141$. Combining (I)–(III): $A(q) \geq P(q)$. $\square$

Proposition 1.4 (Two-column presentation and duality; third pillar). Let ℓ_1, ℓ_2, ℓ_3 be the certified transverse frame $\ell_1 = (0, 2, 0, 1, 0, 1)$, $\ell_2 = (1, 0, 0, 1, 1, 0)$, $\ell_3 = (2, 1, 2, 2, 0, 0)$ (in the coordinates $s_0, \dots, s_5$; all fifteen 3×3 sheet restriction matrices are invertible), and let $\Lambda = R/(\ell_1^q, \ell_2^q, \ell_3^q)R$. Then Λ is a graded Artinian complete intersection (degrees $1, 3, 5, q, q, q$), hence Gorenstein with one-dimensional socle in degree $3q+3$; moreover $\Lambda/(s_1^q, s_2^q)\Lambda = R/\mathbf{m}^{[q]}R$ (Frobenius linearity collapses the six s_i^q into $(\ell^{[q]}) + (s_1^q, s_2^q)$), and the perfect socle pairing yields the degreewise duality

$$A_d = \dim \text{ann}_{\Lambda}(s_1^q, s_2^q)_{3q+3-d} \quad \text{for all } d. \quad (1)$$

2 The Recognition Theorem and the census

For a pair $p = \{a, b\}$ let $V(p) = \bigcup_{J \ni p} L_J$ (three sheets) and $I(p) = I(V(p)) = (s_a + s_b) + (\varepsilon_1^p, \varepsilon_3^p)$, where ε_k^p are the elementary symmetric polynomials of the four remaining variables. $I(p)$ is radical: modulo $\varepsilon_1, \varepsilon_3$ factors into three distinct linear forms in characteristic 3.

Theorem 2.1 (Recognition). *Let $q = 3^v$ and let $\lambda = (\lambda_0, \dots, \lambda_5)$ be homogeneous of degree $e < q$. Then $\sum_i \lambda_i s_i^q \in E$ if and only if $\lambda_a - \lambda_b \in I(\{a, b\})$ for each of the fifteen edges of K_6 . Consequently the space of window syzygies at coefficient degree e is canonically independent of v .*

Proof. $F = \sum \lambda_i s_i^q \in E$ iff F vanishes on every sheet. On L_J , with pair coordinates u_k , $F|_{L_J} = \sum_k (\lambda_{a_k} - \lambda_{b_k})|_{L_J} u_k^q$. For $e < q$ two products $u_k^q m$, $u_l^q m'$ with $k \neq l$ and $\deg m, \deg m' < q$ have disjoint monomial supports, so $F|_{L_J} = 0$ iff each coefficient vanishes on the sheet; collecting the three sheets through a fixed edge gives membership in $I(p)$. $\square$

Let $M = \{\lambda \in S^6 : \lambda_a - \lambda_b \in I(\{a, b\}) \ \forall \text{ edges}\}$, a finitely generated graded S -module, and let $\sigma_e = \dim M_e - [\text{HF}_S(e) + 5 \dim E_e]$ be the census (solutions modulo the trivial family $\mu \cdot (1, \dots, 1) + E^6$).

Theorem 2.2 (The census law; certified). $\sigma_e = 15e^2 - 90e + 145$ for all $e \geq 4$, and $\sigma_{0..3} = 0, 0, 1, 5$.

Proof (certificate). $M = \ker(\Phi : S^6 \rightarrow T = \bigoplus_{(J, \text{pair})} S/I_J)$ with the explicit 45×141 presentation. The kernel is computed by a complete Gröbner basis of the extended module in a block elimination order (*no* pair-skipping criteria; the product criterion is not valid for modules); the zero-first-block elements form a Gröbner basis of the syzygies (Schreyer) and project onto 33 generators of M . A second Gröbner basis of $M \subset S^6$ gives the exact Hilbert series with kernel-side reading $\text{HF}_M(e) = 6 \binom{e+5}{5} - \sum_i N_i \binom{e-i+5}{5}$ (numerator width 10). Gates, all byte-exact: (A) every generator of degree ≤ 4 vanishes on all fifteen sheets; (B) $\text{HF}_M(0..5) = 1, 11, 52, 171, 456, 1032$, the values of two independent prior computations; (C) the law holds for $4 \leq e \leq 100$; since both sides are single quadratic polynomials for $e \geq 13$, agreement there is an identity. All values reproduced in Macaulay2 (Section 6). $\square$

Corollary 2.3. *For every q and every $0 \leq e < q$: $\dim(\mathfrak{m}^{[q]}R)_{q+e} = 5 \text{HF}(R)_e - \sigma_e$, hence $A_{q+e} = \text{HF}(R)_{q+e} - 5 \text{HF}(R)_e + \sigma_e$ — an explicit polynomial, uniform in the tower. Also $A_d = \text{HF}(R)_d$ for $d < q$ (Zone A).*

3 The top zone by duality

By (1), the top degrees $d \geq 2q$ of A are the low degrees $c = 3q+3-d \leq q+3$ of the annihilator. For $c < q$, an element $x \in \Lambda_c = R_c$ lies in $\text{ann}(s_1^q, s_2^q)$ iff there exist $Y, Z \in S_c^3$ with $s_i^q X - \sum_k \ell_k^q Y_k \in E$ ($i = 1$ with Y ; $i = 2$ with Z). Since $(\sum c_k u_k)^q = \sum c_k u_k^q$ over $\mathbb{F}_3$ (Frobenius), the sheet conditions are *linear with fixed frame coefficients*: the annihilator in degrees $c < q$ is governed by fixed, tower-free modules $M' \subset S^7$ and $N_3 \subset S^3$, with $\text{ann}_c = \text{HF}_{M'}(c) - 2 \text{HF}_{N_3}(c) - \dim E_c$.

Theorem 3.1 (Top main body; certified). $\text{ann}_c = 15 \binom{c-4}{2}$ for all c (zero for $c \leq 5$); hence $A_d = 15 N_q(d)$ for all $d \geq 2q + 4$ and all v , where $N_q(d) = \#\{a \in [0, q-1]^3 : |a| = d\}$.

Proof (certificate). The same elimination pipeline applied to the explicit presentations of M' (90 slots, 277 columns) and N_3 (45 slots, 138 columns). Gates: the nine dual anchors $\text{ann}_{0..8} = 0, 0, 0, 0, 0, 0, 15, 45, 90$ from the sealed $v=2$ footprint; consistency $\text{ann}_{9..12} = 150, 225, 315, 420$; the law for $0 \leq c \leq 120$ (polynomial identity past width). All pass byte-exact and were reproduced in Macaulay2 with its native kernel machinery (Section 6). $\square$

4 The collar

Four degrees remain: $d \in \{2q, \dots, 2q + 3\}$, i.e. $c \in \{q, \dots, q + 3\}$, where q -freeness genuinely breaks. Write $e' = q + f$, $f \in \{0, 1, 2, 3\}$, and let $\text{Syz}(e') = \{\lambda \in (R_{e'})^6 : \sum \lambda_i s_i^q = 0 \text{ in } R\}$.

Theorem 4.1 (T-block decomposition). *Let $q \geq 9$, $f \leq 3$, and $\sum \lambda_i s_i^q \in E$ with $\deg \lambda = q + f$. On each sheet write $\Delta_k = (\lambda_{a_k} - \lambda_{b_k})|_{L_J} = \Delta_k^\emptyset + \sum_l u_l^q \delta_{k,l}$ (reduced part plus large parts, $\deg \delta = f$). Then the sheet identity splits into disjoint monomial classes and forces: (i) $\Delta_k^\emptyset = 0$; (ii) $\delta_{k,k} = 0$; (iii) $\delta_{l,k} = -\delta_{k,l}$ (Koszul antisymmetry), with cofactor degree $f \leq 3$.*

Proof. Since $\deg \Delta = q + f < 2q$, each monomial has at most one exponent $\geq q$, giving the canonical split. In the identity $\sum_k u_k^q \Delta_k = 0$ the classes {exactly one exponent in $[q, 2q - 1]$ }, {one exponent $\geq 2q$ } and {two exponents $\geq q$ } (possible as $q + f < 3q$ forbids three, $q \geq 9$) are disjoint, so each vanishes separately, yielding (i)–(iii). $\square$

We now state the aggregate factoring as an explicit identity of linear maps, so that its correctness — including for monomials heavy for several pairs at once — can be checked by inspection. Write $\lambda_i = \sum_m c_{i,m} s^m$. The monomial-wise restriction formula is exact:

$$s^m|_{L_J} = (-1)^{\sum_k m_{b_k}} \prod_k u_k^{m_{a_k} + m_{b_k}}. \quad (2)$$

Fix J , fix l , and let $p = P_l = \{a, b\}$ ($a < b$). By (2), a monomial s^m contributes to the u_l^q -part of $\lambda_i|_{L_J}$ iff its *pair-weight* $w_p(m) := m_a + m_b$ is $\geq q$. Writing such $m = (m_a, m_b, \gamma)$ with γ the restriction of m to the four variables outside p and $w = m_a + m_b = q + t$, the total degree forces

$$t = f - |\gamma| \quad (\text{determined by } \gamma). \quad (3)$$

By (2) the contribution of s^m is $(-1)^{m_b} (-1)^{\epsilon(\gamma, J)} u_l^t \mu(\gamma, J)$, where $\mu(\gamma, J) = \prod_{k \neq l} u_k^{\gamma_{a_k} + \gamma_{b_k}}$ and $\epsilon(\gamma, J) = \sum_{x \in \text{supp } \gamma, x = \text{some } b_k} \gamma_x$ depend only on (γ, J) . *The split-dependence enters only through $(-1)^{m_b}$.* Define the alternating aggregates — linear functionals on the coefficients of λ_i :

$$\tilde{c}_i(p, \gamma) := \sum_{\beta \geq 0} (-1)^\beta c_{i, (w - \beta, \beta, \gamma)}, \quad w = q + f - |\gamma|. \quad (4)$$

Then, coordinate by coordinate,

$$[u_l^q\text{-part of } \lambda_i|_{L_J}] = \sum_{\gamma} (-1)^{\epsilon(\gamma, J)} \tilde{c}_i(p, \gamma) u_l^{f - |\gamma|} \mu(\gamma, J). \quad (5)$$

Lemma 4.2 (Aggregate factoring). *Let $W_f := \bigoplus_{(i,p,\gamma)} \mathbb{F}_3$ (i among the 6 slots, p among the 15 pairs, γ of degree $\leq f$ on the complement of p), and define $\text{agg} : (S_{q+f})^6 \rightarrow W_f$ by (4) and $A_f : W_f \rightarrow \bigoplus_{J, k \neq l} \mathbb{F}_3[u]_f$ by $A_f(e_{(i,p,\gamma)}) = \sum_{J \ni p} \sum_{k \neq l(J,p)} \epsilon_i^{(J,k)} (-1)^{\epsilon(\gamma, J)} u_l^{f - |\gamma|} \mu(\gamma, J)$ in coordinate (J, k, l) , where $\epsilon_i^{(J,k)} = +1$ if $i = a_k$, -1 if $i = b_k$, 0 otherwise. Both maps have fixed coefficients, independent of q . Then*

$$\Phi_{\text{collar}} = A_f \circ \text{agg}$$

as linear maps. Consequently $\text{Im}(\Phi_{\text{collar}}|_{\text{Syz}}) \subseteq D_f := \text{Im}(A_f) \cap \{\text{antisymmetric}\}$, a fixed space.

Proof. Evaluate both sides on λ at each coordinate (J, k, l) : the left side is $\delta_{k,l}^{(J)} = [u_l^q\text{-part of } (\lambda_{a_k} - \lambda_{b_k})|_{L_J}]$; by (5) applied to $i = a_k$ (sign +1) and $i = b_k$ (sign -1) this equals the (J, k, l) -coordinate of $A_f(\text{agg}(\lambda))$. The two sides agree on every λ and every coordinate. Antisymmetry of the image on syzygies is Theorem 4.1(iii). $\square$

Remark 4.3 (Multi-heavy monomials). A monomial may have $w_p(m) \geq q$ for several pairs p simultaneously. This creates no ambiguity: agg is a tuple of linear functionals — the same coefficient $c_{i,m}$ simply enters the aggregates of every pair for which m is heavy, each time with the sign dictated by (4). No disjoint decomposition of λ is used anywhere; the factoring identity is verified coefficient-wise above. Each output coordinate (J, k, l) reads only the $p = P_l$ aggregates, once, so no sign conflict can arise.

Theorem 4.4 (Collar ceilings). *For $f = 0, 1, 2, 3$: $\dim D_f = 10 + 45\binom{f+1}{2} = 10, 55, 145, 280$ (a fixed rank certificate: $\text{rank } A_f = 45, 180, 405, 720$; symmetric parts $35, 125, 260, 440$; reproduced in Macaulay2). Moreover $\ker(\Phi_{\text{collar}}|_{\text{Syz}}) = \pi(M)$, of dimension $\text{HF}(R)_{e'} + \sigma_{e'}$, and the required bound satisfies the q -cancellation identity*

$$5 \text{HF}(R)_{q+f} - \text{HF}(R)_{2q+f} + 15\binom{q-1-f}{2} - \sigma(q+f) = 10 + 45\binom{f+1}{2} \quad (\text{all } q\text{-terms cancel}).$$

Therefore $\text{Syz}(q+f) \leq [\text{HF}(R) + \sigma](q+f) + \dim D_f$, which gives $A_d \leq 15 N_q(d)$ at $d = 2q, \dots, 2q+3$ for all $v \geq 2$.

Proof. The rank certificate is a finite computation on the fixed matrices of Lemma 4.2, verified four ways (Section 6). *Kernel identification (by radicality, not by Recognition out of its range)*: for a syzygy in the kernel, Theorem 4.1(i) kills the reduced parts and $\delta = 0$ kills the large parts, so $\Delta_k = 0$ identically on every sheet; then $\lambda_a - \lambda_b$ vanishes on all three sheets through each edge p , i.e. on $V(p)$, and since $I(p)$ is radical, $\lambda_a - \lambda_b \in I(V(p)) = I(p)$, i.e. $\lambda \in M$; conversely $M \subseteq \ker$ trivially. Its dimension is $\text{HF}(R)_{e'} + \sigma_{e'}$ by the census law (valid for all e). The q -cancellation is a direct expansion with the exact $\text{HF}(R)$ and the certified σ : the coefficients of q^2, qf and q vanish identically ($45-60+15$; $90-60-30$; $-45+90-45$), leaving $10+45\binom{f+1}{2}$; re-verified numerically at $q = 9, 27, 81, 243$. The inequality chain is rank-nullity on $\Phi_{\text{collar}}|_{\text{Syz}}$, and $\text{Syz}(q+f) \leq B(q+f)$ translates through $\dim(\mathfrak{m}^{[q]}R)_{q+e'} = 6 \text{HF}(R)_{e'} - \text{Syz}(e')$ into $A_d \leq 15 N_q(d)$. $\square$

5 The Ledger identity and the proof of Theorem 1.2

Theorem 5.1 (Ledger). *With $ZA(q) = \sum_{d < q} \text{HF}(R)_d$ and $W(q) = \sum_{e < q} [\text{HF}(R)_{q+e} - 5 \text{HF}(R)_e + \sigma_e]$:*

$$ZA(q) + W(q) + 15\binom{q}{3} = P(q) \quad \text{identically for } q \geq 5,$$

both sides being cubic polynomials in q (checkpoints $q = 9, 27, 81, 243$ agree byte-exact; $15\binom{q}{3} = 15 \# \{a \in [0, q-1]^3 : |a| \geq 2q\}$ via the bijection $b = (q-1) - a$). Each summand is a genuine closed form in q as a free variable — $\text{HF}(R)_d$ exactly for $d \geq 4$, σ_e the Hilbert function of one fixed module, the count purely combinatorial — so no hidden v -dependence can hide behind the four power-of-three checkpoints.

Proof of Theorem 1.2. For $v \geq 2$: Zone A and the window are exact (Cor. 2.3); the top main body is exact (Thm. 3.1); the Ledger then gives $A(q) - P(q) = \sum_{d=2q}^{2q+3} [A_d - 15 N_q(d)]$. By Theorem 4.4 each summand is ≤ 0 , so $A(q) \leq P(q)$; with the floor $A(q) \geq P(q)$ (Prop. 1.3), $A(q) = P(q)$. For $v = 1$, $A(3) = 141 = P(3)$ is the sealed direct computation. $\square$

6 Verification record, reproducibility, and a referee’s guide

All machine steps are finite computations over $\mathbb{F}_3$ on fixed (q -free) objects, each gated before reading. Every load-bearing certificate has been reproduced **four independent ways**: (1) the authors’ engines (Gröbner-kernel and rank computations, Python and portable C++, with byte-exact gates against the sealed anchors $A(3) = 141$, $A(9) = 7761$ and their full degree-wise footprints, cross-checked on independent hardware); (2) re-runs of all engines on a second machine; (3) from-scratch reconstructions by an independent implementation using *different methods* (direct linear algebra in place of Gröbner bases), agreeing byte-exact; (4) **Macaulay2** (v1.26.06): all three certificates re-implemented using M2’s native `kernel`, `hilbertFunction` and `rank` machinery — census $\mathrm{HF}_M(0..5) = 1, 11, 52, 171, 456, 1032$ and the σ -law; collar $\dim D_f = 10, 55, 145, 280$; annihilator HF_{N_3} , $\mathrm{HF}_{M'}$ and $\mathrm{ann}_c = 0, \dots, 0, 15, 45, 90, 150, 225, 315, 420 = 15\binom{c-4}{2}$ through $c = 12$ — all byte-exact on the first run.

A referee’s guide. In our own judgment the four load-bearing joints of this proof, in decreasing order of the attention they deserve, are: (i) Lemma 4.2, now stated as a coordinate-wise identity precisely so that Remark 4.3 can be checked by inspection; (ii) the range hygiene of every theorem (the kernel identification in Theorem 4.4 deliberately uses radicality, not Recognition, at $e \geq q$); (iii) Proposition 1.3, whose only imported ingredient is the [1] identification of P as the characteristic-zero target; and (iv) the geometric dictionary between the algebraic statement proved here and Conjecture 1.2 as formulated in [1], which belongs to the experts of that framework and which we cannot strengthen from inside. An expanded version of this guide, with attack suggestions for each joint, accompanies the repository (`SOFA_WHERE_TO_ATTACK.md`).

Status. Every link of the proof is either a pencil argument valid for all v or a finite gated certificate on a q -free object; the computational layer has been verified as described above. The record has not yet been refereed by a human expert; it is offered to the community precisely for that examination. The complete verification dossier (including seven independent logical reviews across four AI model families), every engine with its logs, the Macaulay2 scripts, and standalone write-ups of each theorem are available at github.com/tretoef-estrella.

References

- [1] A. Degtyarev, I. Shimada, *On the topology of projective subspaces in complex Fermat varieties*, J. Math. Soc. Japan 68:3 (2016), 975–996, arXiv:1405.4683.