

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Contents

1	Introduction	1
1.1	Competition	1
1.2	Substrates	1
1.3	Growth Model	1
1.4	Environmental Influence on Hyphal Tip Extension (ν)	2
1.5	Decomposition kinetics	2
1.6	Enzyme interactions	2
2	Model Formulation	3
2.1	Growth Model	3
2.2	Defining Representative Enzymes	4
2.3	Estimating ν For Various Environments	4
2.4	Other Responses to Temperature and Water Potential	5
2.5	Assumptions	6
2.5.1	Growth	6
2.5.2	Decomposition Kinetics	6
2.5.3	Temperature/Moisture effects	6
3	Parameter selection and representative result	6
3.1	Growth model parameters	7
3.2	Decomposition parameters	7
3.3	Representative results	7
4	Analysis	7
4.1	Distribution of relative concentration of fungi	7
4.2	Environmental fluctuations	8
4.3	ν vs Average Contribution to Substrate Decay	9
4.4	Total Carbon vs Time for Specific Environments	9
4.5	Average Contribution vs ν with Competitive Rankings	9
A	Model Parameters	10
B	Appendix Tables	10
C	Appendix Figures	11

List of Tables

1	Specific Environments	5
2	Temperature & Water Potentials	5
3	Model Parameters	10
4	Environment Temperature Data	10
5	Environment Water Potential Data	11

6	Environmental & Parameter Table	11
---	---	----

1 Introduction

Decomposition by fungal and microbial communities is second only to photosynthesis in driving carbon cycling in the Earth's ecological communities [18]. While this significance is rarely understated, inner mechanisms and interactions within decomposition communities have been poorly understood and treated as a 'black box' in the past by scientists studying ecosystem processes [9].

Fungi are made up of hyphae: long winding filaments enmeshed together to form a thick mat of mycelium. At the frontline of battle, hyphal tips are extending forward and branching to cover new area at a velocity dictated by the hyphal extension rate, and concurrently other hyphae are dying off or merging together [4]. Like machinery, each fungus has ideal operating conditions. Fungi native to dry or arid biomes will grow efficiently in a moisture-lacking environment when other fungi will falter. Maynard et al. (2019) defines these operating conditions by assigning temperature and moisture niche-widths to 37 studied species of fungi [28].

1.1 Competition

When fungi come into contact with each other, competitive interactions are most often observed. Tactics studied in Boddy, L. (2000) include deploying diffusable toxins, obtaining nutrients parasitically, destruction of hyphae by intentional interference, and establishing wall-like barricades to thwart attackers [11]. Most researchers, including Maynard et al. (2019) construct a competitive ranking scale based on experimental observations to encapsulate the ramifications of these complex interactions [28].

Under a constant temperature regime, previous research concludes that dominant fast-growing suppress other species which negatively impacts the overall community's efficiency. However fluctuating environmental conditions change the outcome and promote complementary growth [15]. An explanation for this is presented by Lusten-

houwer et al. (2020) suggesting a spectrum between slow-growing, stress-tolerant fungi operating in a wider environmental range than fast-growing, highly competitive fungi [31].

1.2 Substrates

Determining which fungi are present in a community depends on the organic material, or 'substrate', being broken down. In the case of fungal-mediated wood decomposition, the substrate is made up of lignocellulose polysaccharides, and the fungi involved are equipped with specialized enzymes capable of splitting linkages in the polymers to harvest usable carbohydrates. Long-term wood decomposition follows a well-studied three-phase process: the first phase is dominated by fast and opportunistic fungus claiming easily-consumable soluble sugars, followed in the second phase by consistent and dependable decomposers that degrade the holocellulose constituting 65%-85% of the material [21]. Slow and highly specialized fungi trail behind in the third phase to clean up rigid lignin polymers (15%-35%) with powerful oxidative enzymes [14]. While phase progression is an important agent governing community changes on the long-term scale, this paper will focus on only the middle phase of cellulose decomposition and will not consider lignin decomposition. [31].

1.3 Growth Model

An important step in modeling fungal communities is acknowledging separation between the growth rate and the decomposition rate of the fungus, which are influenced by different factors. Fungal growth models typically focus on the increase in overall biomass due to hyphal extension. Fungi with a greater density of hyphal tips per unit volume will grow and accrue biomass faster, as hyphae only grow from their tips. New tips are created when hyphae branch off, while others die or fuse together via anastomosis. Edelstein (1982) models these dynamics with the following:

$$\frac{\partial \rho}{\partial t} = nv - d \quad (1)$$

$$\frac{\partial n}{\partial t} = -\frac{\partial nv}{\partial x} + \sigma \quad (2)$$

Where ***** These equations will operate standing alone, without any other influences by enzymes or growth limitations. This allows many models to treat it as a baseplate to expand upon. Edelstein (1983) and Davidson et al. (1997) do this by considering concentrations of nutrients inside the fungi and outside in the surrounding substrate [5, 20]. Our model will likewise follow in eliminating the assumption that fungi grow over an unlimited nutrient supply. A more contemporary approach by Du et al. (2019) describes the growth of a fungus with a 3-dimensional reaction-diffusion equation based on microscopic growth mechanisms, also focusing on tip density and extension rate but with an added term to identify the proportion of tips which are actively growing [26].

1.4 Environmental Influence on Hyphal Tip Extension (ν)

Environmental impacts are introduced to the decomposition model through ν , the hyphal tip elongation rate. The mechanism of elongation is relatively unknown; as described in Gervais et al. (1999), "cell turgor pressure corresponds to an overpressure which allows the cell morphology, elongation, division and hence the biomass evolution" [10, 16]. In fungi, cell turgor comes from diffusion of water dictated by the water potential¹ gradients rather active transport within the cell, making it very environmentally dependent [10]. This fact justifies determining ν based on environmental factors on fungal decomposition rates by varying ν outside of baselines for given species in set locations in our model. Experimental data from Maynard et al. (2019) shows the relationship between ν and the environmental parameters of water potential (ψ) and temperature (T) [28].

¹Water potential (ψ) describes the content of availability of moisture surrounding a fungi as dictated by the soil composition.

1.5 Decomposition kinetics

Moore et al. (2015), Edelstein (1983), Moorhead et al. (2006), Parnas (1974), and Schimel et al. (2003) describe the decomposition of a certain substrate by a microbial decomposer as a function of the substrate's concentration with the Michaelis–Menton equation [3, 5, 13, 14, 22]. Thus, decomposition of carbon in a substrate is described by the following equation.

$$\frac{dC}{dt} = \frac{KBC}{K_e + C} \quad (3)$$

Where ***** Schimel et al. (2003) and Moorhead et al. (2006) assert that decomposition levels off at a maximum rate proportional to the concentration of enzymes acting on it [13, 14]. To determine this enzyme concentration, we will make the assumption that enzyme concentration is proportional to the concentration of biomass in our system represented by r multiplied by two environmental limiting factors in $K = rS_M S_T$ [13]. This environmental factors are discussed at greater length in **section that they're discussed**

1.6 Enzyme interactions

Enzymes produced by fungi are expelled into the surrounding substrate to autonomously break down macromolecules. Relative enzyme concentrations can be quantified by the production rates of each enzyme by a fungus. Production of enzymes is a metabolically taxing process, therefore fungi that produce enzymes which return much more usable energy and C than was put into making the enzyme, a high "return on investment", will be able to synthesize even more enzymes and grow rapidly [13]. Lustenhouwer et al (2020) strengthens this theory by showing through statistical analysis of 1582 various fungal isolates that fungi containing larger proportions of enzymes like acid phosphatase, which are associated with

slow lignin decomposition, are demonstrated to strongly correlate with slower decomposition rates [31]. Lignin exemplifies a low return on investment, requiring powerful enzymes and greater energy input per unit C than holocellulose. But it's ultimately the fungi that have a subset of enzymes that best match their surrounding substrate which successfully grow and establish a niche. As an example, in a substrate of 70%+ lignin, the slower growing fungi with lignin enzymes normally at a disadvantage will win out over other fungi competing for a fraction of available holocellulose.

Communities with similar fungi of similar enzymatic makeup often see a few dominant species overwhelm the rest through competition for similar resources in the substrate, but communities of fungi with greatly differing enzymes will instead form distinct niches that partition different resources in which they are separately able to grow. In these conditions, one dominant species faces difficulty in wiping out competitors who are individually masters of their own substrates. This effect is known as niche complementarity [15]. Research by Toljander et al. (2006) contends that environmental fluctuations further robustify a diverse community as faster growing highly competitive fungi are hindered by conditions that lapse out of their moisture and temperature niche, and slower growing stress-tolerant fungi benefit from decreased pressure by competitors. The resulting dampened and stable diverse community will decompose the most efficiently across a range of conditions [15].

2 Model Formulation

2.1 Growth Model

In the review of various fungal growth models described in Lin et al. (2016), a specific method of modeling branching and hyphal death was found to match experimental observations in the widest variety of settings. Based on these results, for σ and d in (1) and (2) we will choose to imple-

ment the following relationships, comprising dichotomous branching, tip-hypha anastomosis, and hyphal death (or YHD).

$$\sigma = \alpha_1 * n - \mu n \rho \quad (4)$$

$$d = \gamma_1 \rho \quad (5)$$

Enzymes conducting decomposition make up a fraction of the total biomass, and the total biomass changes according to the growth rate. The total biomass B defined in (6) multiplies by maximum rate of the Michaelis-Menton equation (3) to couple growth rate with decomposition rate.

$$B = \int_{allx} \rho dx \quad (6)$$

The rate of decomposition of carbon will then be dynamically effected by the growth of the fungus contrary to the typical assumption of a static B as described by Schimel et al. (2003) [13]. Our formulation of K which accounts for the proportion of biomass in the fungal growth containing the necessary enzyme for the carbon decomposition will change to account for this.

$$K = r * S_M * S_T * 0.04 * G \quad (7)$$

Where r is a dimensionless parameter representing the relative abundance of one enzyme out of the total enzyme biomass of that specific fungus. The multiplication by 0.02 is a generalization that for any given species of fungus, 2% of the biomass will be relevant enzymes [14]. G is an empirically derived rate constant relating the mass of relevant enzymes to the maximum rate of the decomposition.

Limited carbon availability will also impact a fungi's growth rate. To account for this, we include a term multiplying ν that can be interpreted as the available carbon for the fungi to consume for growth. The first equation in the growth model then becomes

$$\frac{\partial \rho}{\partial t} = (1 - LCI) * (10^9) * C * n * \nu - d \quad (8)$$

Where LCI is the lignocellulose index² of a certain material and $(1 - LCI)$ is then the proportion of the substrate carbon stored in holocellulose. Multiplying by $(1 * 10^9)$ is approximation to convert grams of carbon to hyphae length in mm. [citation needed for g to mm] An investigation of the effects of including the carbon term in the growth model are addressed in a later section.

So our coupled growth and decomposition models can be summarized by equations (2), (3), (6), and (8).

2.2 Defining Representative Enzymes

Maynard et al. (2019) collected data for 8 relevant enzymes and utilized a standard clustering approach to create a subset of four enzymes which best represented the effects by the fungi's enzymatic makeup on competitive interactions [28]. Our model utilizes this open-source data by normalizing the production rates of each enzyme and calculating their relative abundance as a fraction of the total enzyme biomass for that specific fungus. Interactions caused by differing rates of decomposition among different enzymes in varying environmental conditions is then represented in our model by taking the weighted sum of all contributions to the decomposition rate from any one of four enzymes in a given fungi species, represented in the following equation

$$\frac{dC_1}{dt} = c_{1,a} \frac{de_{1,a}}{dt} + c_{1,b} \frac{de_{1,d}}{dt} + c_{1,c} \frac{de_{1,c}}{dt} + c_{1,d} \frac{de_{1,d}}{dt} \quad (9)$$

Where $c_{1,a}$ is a enzyme breakdown efficiency coefficient for the fungal species 1 and enzyme a . Here the rate $\frac{dC_1}{dt}$ is representative of the rate of decay of the substrate by the fungi species 1. Breakdown efficiencies $c_{1,a}$ for each enzyme were derived from correlation coefficients drawn from experimentally

observed data in Lustenhouwer et al. (2020) [31].

The total decomposition rate is then given by the sum of all rates over the various species of fungi.

$$\frac{dC_{tot}}{dt} = \sum_{i=1}^n \frac{dC_i}{dt} \quad (10)$$

Where n is the total number of fungi being simulated. **address the fact that all the 35 fungi will be growing/decomposing together at the same time**

2.3 Estimating ν For Various Environments

When considering a sampling of decomposition rates in various environment types, we must determine how to estimate temperate and water potentials that ground fungi would experience. In the case of determining ν , this requires finding projected temperature and water potential.

Amongst existing biome classification models, Whitaker's scheme [2] is notably simple, providing a layout of biomes based on mean annual temperature and mean annual precipitation. However, modern classification of biomes has drifted away from using these traits as definitive biome identifiers [24]. These traits alone do not define all features of concern to fungal growth, such as soil composition for example. In addition, by Whitaker's scheme, we find that a given biome can have a wide range of average annual temperatures (such as arid desserts ranging from about -10 to 30 °C) [2]. We can sample a range of temperature and moisture values to output various ν and then select certain regions to profile in order to gauge potential environments where fungi can decompose and interact.

The following are the specific environments selected to represent various biomes:

²The lignocellulose index describes the ratio of non-hydrolysable lignin to hydrolysable holocellulose in litter

Biome	Specific Environment Selected
Dessert (Arid)	Sonoran Desert, USA
Grasslands/shrublands	(Semi-Arid), central Argentina
Temperate Forest	Sal Forests, central Himalayas
Boreal Forest (Arboreal)	Pine Forests, central Himalayas
Tropical Rain Forest	Tropical Forests, Barro Colorado Island, Panama

Table 1: Specific Environments

Although water potentials can be approximated based on predictive models, the measure is best found experimentally from soil samples [25]. Single water potential and annual temperature values were selected from ranges of values experimentally determined by a variety of field studies on these environments (see: Table 2). This decision was made since we're aiming to compare discrete environmental conditions rather than create a complete span of environmental conditions.

Biome/ Environment	Temperature [$^{\circ}\text{C}$]	Water Potential [MPa]
Dessert (Arid)	15	-4.5
Grasslands/shrublands (Semi-Arid)	15.3	-3.2
Temperate Forest	12.49	-1.09
Boreal Forest (Arboreal)	12.49	-1.51
Tropical Rain Forest	27.5	-0.79

Table 2: Temperature & Water Potentials

Note that these are not wholly representative configurations, but rather examples to provide insight into how fungal species with specific traits may respond in discrete and distinct conditions likely to exist. We can then find ν using experimental data from Maynard et al 2019 [28]. **talk about continuous data technique thingy in the extra methods info sheet**

2.4 Other Responses to Temperature and Water Potential

Our growth model takes into account temperature (T) and water potential (ψ) in two more parameters: the soil temperature coefficient (S_T) and the soil moisture coefficient (S_M). Moorhead et al. (1991) provides a simple relationship between S_T and T using the rate of increase (Q):

$$\log_{10}(S_T) = \frac{T - 25}{10} \log_{10}(Q) \quad (11)$$

Although this equation does not take into account specific fungal response to temperature change, more recent evidence supports that this relationship is not direct, with most of the direct impact

coming from moisture [Petraglia2018]. As the ratio of rates of decomposition given a temperature change, Q as a parameter should represent the effective output of various mechanisms influenced by temperature rather than focusing on specific mechanisms. However, for the sake of simplicity, Q has been set standard constant to a value of 2.5 [7].

Water potential also relates to a constant, S_M , in a simple equation described in Moorhead et al. (1991) using α_2 and λ :

$$S_M = \alpha_2 - \lambda \log_{10}(-\psi) \quad (12)$$

These two parameters help calculate the max-

imum growth rate by the relationship:

$$\beta = S_T S_M r \quad (13)$$

why is this β and not K ?

2.5 Assumptions

Further assumptions made in the formulation of our model that have not yet been discussed are largely simplifications that are not relevant to any significant or interesting discussion. An itemized list of these assumptions is provided below.

2.5.1 Growth

- Neglecting effects from direct interactions between fungi
- Neglecting effects from N limitation
- Litter is not added to or removed from the carbon pool by forces external of the fungi
- LCI is xxx and remains constant over time
- Dynamics regarding the rate of nutrient absorption and transport of metabolites within the mycelium are ignored or simplified
*more specific
- Ignoring other growth traits: LIST
- Hyphal death rate γ_1 remains constant regardless of environmental fluctuations
- 100% of living hyphal tips are active
- No biomass created by a fungus will be geometrically isolated from the substrate
- Neglecting effects due to spacial limitations

2.5.2 Decomposition Kinetics

- Neglecting effects due to enzymes not included in the model

- Production of enzymes increases proportionally with biomass
- For any species of fungus, 2% of the biomass will be relevant enzymes
- Relative proportions of enzymes will stay constant regardless of environmental fluctuations
- Neglecting effects on decomposition rate due to changing metabolic rate of the fungi

2.5.3 Temperature/Moisture effects

- Turgor is a function of water potential
- Fungal decomposition occurs in surface soil (0-30 cm)
- Experimental data for water potentials in specific biomes/environments are representative of the conditions of those environments
- Rate of increase of temperature Q remains constant regardless of environmental fluctuations

3 Parameter selection and representative result

To select realistic values for the parameters, we created a representative run of the coupled growth and decomposition model, consulting the literature to find parameter values. An overview of the parameter selection for this representative run can be seen in (parameter table), however several parameters are worth some discussion.

3.1 Growth model parameters

The effect of two main parameters were studied for their effect on the growth model's result: the branching rate and the hyphal death rate. These parameters were not agreed upon in the literature so their effect on the model output was evaluated and they were selected from a pool of possible values based on the most likely realistic output. Most results of the growth model were comprised of traveling wave solutions, converging to a uniform distribution in both space and time. This can be thought of as the maximum growth of the fungus into its total space. The branching rate (α_1) was found to increase the hyphal density in the end behavior of the solution, with the ending density increasing as α_1 increases. The hyphal death rate (γ_1) was found to increase the oscillations in time and thus larger γ_1 values would increase the oscillations and the time to reach a given end-behavior. Many of the papers discussing the values of these parameters [Edelstein 1982] [Lin 2016] [Schnepf 2007] [Du 2019] are concerned more with short term dynamics in perfectly ideal conditions. The purposes of this paper is to assess the long term decomposition rates under different conditions, so parameters that can predict long-term behavior, comparable to long term decomposition dynamics described in [Moorhead 2000] [Moorhead 2006] [Moorhead 1991] were selected.

3.2 Decomposition parameters

The value for G is interpreted as a rate constant, relating a concentration of relative enzymes to the maximum rate of carbon decomposition. formulation for S_M and S_T come from [Moorhead 1991] and in said model, a simpler equation is used for the reaction dynamics for decomposition of carbon. Because of this, we cannot use the same rate constant and must base this parameter of agreement with other models or experimental data. The most important aspect we examined for the section of G was the timescale of decomposition to be comparable with that of [Lustenhauer 2019].

3.3 Representative results

As a representative result of the coupled growth and decomposition model, the two were run using the parameters summarized in (parameter table). In the growth model, ρ was found to exhibit traveling wave dispersion throughout space, as similarly shown in [Mimura 2000] and [Edelstein 1982]. Due to the simulation only running for a timespan of 3 years, the decrease in carbon over time looks fairly linear, with slight fluctuations likely due to the fluctuating fungal biomass densities over time. The first initial runs were done without the coupling term in the growth equations, so the growth model was assuming an abundance of nutrients and not affected by the decreasing concentration of carbon. Depending on the timescale being simulated, that could be a perfectly valid assumption, as short time scales tended to show little overall decrease in the concentration of holocellulose carbon. However the differences seen in including the nutrient availability term in the growth equations become more apparent as longer time scales are considered. The effects of this can be seen in our results. The overall biomass was also mostly unaffected other than replacing decaying oscillations towards a steady state with a somewhat constantly increasing biomass overtime with a decaying slope. Overall, including the nutrient availability term in the growth equation captures more of the fundamental interaction that this paper is concerned with (that of decomposing wood and ground litter rather than free-growing fungi), so it was included in further analysis.

4 Analysis

4.1 Distribution of relative concentration of fungi

The model allows us to adjust the relative concentrations of each fungi species being simulated. In the model, this is represented with a set of $0 < c_i \leq 1$ such that $\sum c_i = 1$. Each one of these c_i

multiplies the rates of hyphal density increase and boundary/initial conditions in the growth model. With this metric, the relative concentrations of different fungi species are forced to satisfy the conditions $B_i/B_{tot} = c_i$ where B_i is the biomass of the i 'th fungi species. This is obviously an unrealistic constraint on the system as the relative concentrations of different fungal species should fluctuate with time as the different species of fungi grow at varying rates. But for the purpose of assessing the effect of biodiversity on the overall decomposition of the substraight, concentration fluctuation would add unneeded complexity to the analysis.

Also considered for this analysis are the competitive rankings of the different fungi species (CITATION NEEDED). The ranking-distribution coefficient, a is then defined to be

$$a = \sum_{i=1}^n c_i * R_i \quad (14)$$

Where R_i is the competitive ranking for the i 'th fungi species. The ranking-distribution coefficient can be thought of as a dot-product in n -dimensional space (where n is the number of various fungi species being simulated). So a will be highest when the n -dimensional vectors "line-up" in the sense that the distribution of fungi species is closest to the distribution of competitive rankings.

To study the effects of a on the system overall, sample c values had to be chosen. Due to a lack of literature results on simple, constant fungi distributions, arbitrary Gaussian distributions spread out over the different species of fungi were chosen. Because we are mainly concerned with the relation of any given distribution to the competitive rankings, the functions can be viewed as arbitrary or random distributions (because the ordering of fungal species in our Gaussian distribution has no physical meaning). The results of model runs using these various distributions is shown in (figure carbon-decrease-fungi-distribution). To assess the overall effectiveness of a single species distribution, the average carbon decrease over the whole interval was found via the equation

$C_{avg} = \int (C_0 - C)dt/(\delta t)$ where C_0 is our initial amount of carbon and δt is the total time change over which the carbon decreases. The results of comparing this average carbon decomposed to the a value for the species distribution in question is seen in (figure a-competitive-distribution-coef-to-avg-Clost). Several interesting qualitative behaviors arise from this comparison. Firstly, there is a clear smooth trend between a and the average carbon decomposed up to approximately $a = 0.65$. At this point the smooth trend completely vanishes and the data points appear to be randomly distributed in the area. We hypothesize that this critical value of a could be representative of a bifurcation point; however further analysis on the dynamics of these coupled systems interacting would need to be done to investigate this hypothesis further. Additionally we notice that at this critical point, the average carbon decomposed tends to grow significantly. The physical interpretation of this critical point is deeply related to the effectiveness of the competitive rankings as they relate to a biologically diverse system. The results show that beyond a certain point, distributing the biodiversity more closely to the competitive rankings does not have any noticeable effect on the overall effectiveness of the system.

This leads to two possibilities beyond the critical point for the system. One possibility is that optimum biodiversity is based on more complex dynamics than the competitive rankings can assess. The other possibility is that the system behaves in a chaotic or random nature beyond this critical point and no clear metric of biodiversity could be correlated to the effectiveness of a given distribution.

4.2 Environmental fluctuations

Realistically, the external environment dictates environmental conditions of the system both in short time scales (year fluctuations for example) and long time scales. To study the effects of these short time scale fluctuations, we choose to implement a yearly moisture cycle. The cycle dictates a repeating, yearly pattern of ψ and thus v . To Incorporate

porated this into the model, ν and ψ were dynamically allocated over time based on the oscillatory behavior shown in (figure oscillating-psi-nu). Due to time limitations and scarce data on the parameters of these yearly oscillations, only two species of fungi were simulated. A comparison of their respective decomposition contributions with that of similar environmental conditions but static ν and ψ is shown in (figure oscillating2nonoscillating-psi-contribution). There is no note-worthy new oscillatory motion seen in these results. There is also seen to be a slight increase in the effectiveness of both fungi in the oscillatory case.

4.3 ν vs Average Contribution to Substrate Decay

Using the decomposition model, average contribution to substrate decay and the tip elongation rate ν can be found for each fungal species for a given environmental configuration. Aside from indicating an expected positive correlation between ν and contribution, we find that in general, environments with a higher moisture potential lead to greater contributions overall and a greater contribution sensitivity to ν in our model. Our model agrees with the notion of water potential serving as an important limiting factor to the decomposition ability of individual fungi. This corroborates with various findings, that find soil moisture as a particularly influential environmental factor, influencing elongation through pressure gradients and testing of tolerance [28] [31].

4.4 Total Carbon vs Time for Specific Environments

Similarly, a sampling of different environmental parameters within the decomposition model demonstrates the decrease of total carbon in a fungal-waste system over time. Here, we see more moist soil environments lead to clearly more expedient rates of carbon total carbon decrease and reach total decomposition quicker in our model. Again in the model, the rain forest environment exceeds the other environmental profiles, effectively completing decomposition in under 1640 days (roughly 4.5 years).

4.5 Average Contribution vs ν with Competitive Rankings

With one set of environmental parameters, we can gain insight into how different species of fungi potentially interact relative to each other. In addition to an expected loose positive correlation between ν and the average contribution of a given fungi, there is a loose correlation of higher fungi competitive ranking with greater ν and greater average contribution. This suggests that the fungi that are more 'active' in our model, with the largest growth rates and proportions, are also the most competitive. Competitive ranking has a higher correlation with ν , indicating that in our model growth rate may be more significant than proportion when it comes to competitive circumstances. This aligns with established trends of the primary 2 fungal niches [28].

A Model Parameters

Here are the model parameters for the coupled growth and decomposition model for *Armillaria gallica* located at 30.465247 degrees latitude and -89.040298 degree longitude secreting cellobiohydrolase (Cel7A) to decompose hardwood holocellulose [Kari2014, 28]:

Parameter	Symbol	Value	Units	Source and Specification
Half-Saturation constant ³	K_e	7	$\frac{g_{enzyme}}{L_{litter}}$	[Kari2014] Enzyme
Holocellulose carbon ⁴	$1 - LCI^5$	0.709	N/A	[21]
Hyphal tip elongation rate	ν	0.250	$\frac{mm}{day}$	[28] Species, ψ , T
Temperature	T	25	$^{\circ}C$	[28] Specie's habitat
Water potential	ψ	-0.5	MPa	[28]
Enzyme biomass ratio ⁶	r	0.437	$\frac{g}{g}$	[28] Species
Hyphal death rate	γ_1	0.15	day^{-1}	[17]
Anastomosis coefficient	μ	0.3	$\frac{mm}{day}$	[Lyn2016]
Branching rate	α_1	1.2372	day^{-1}	[26]
Intercept of S_M function ⁷	α_2	0.311	N/A	[7]
Slope of S_M function	λ	0.345	N/A	[7]
Soil moisture coefficient	S_M	0.4149	N/A	[7] ψ
Soil temperature coefficient	S_T	1	N/A	[7] T
Rate of Increase	Q	2.5	$^{\circ}C$	[7] T
Rate constant ⁸	G	10	$g * mm^{-1} * day^{-1}$	[31] ⁹

Table 3: Model Parameters

B Appendix Tables

Biome/ Environment	Average Annual Temperature [$^{\circ}C$]	Selected Value [$^{\circ}C$]
Dessert (Arid)	10 to 20 [Davey2007]	15
Grasslands/shrublands (Semi-Arid)	15.3 [8]	15.3
Temperate Forest	-1.42 to 26.39 [30]	12.49
Boreal Forest (Arboreal)	-1.42 to 26.39 [30]	12.49
Tropical Rain Forest	23 to 32 [29]	27.5

Table 4: Environment Temperature Data

³Also called Michaelis Constant.

⁴We assume that all carbon compounds excluding lignin are holocellulose.

⁵Where LCI is the lignocellulose index.

⁶Proportion of specific enzyme biomass to total enzyme biomass.

⁷The intercept of soil moisture effect on decay rate.

⁸Proportionality constant between maximum rate of decomposition and enzyme biomass.

⁹Empirically derived.

Biome/ Environment	Water Potential Range [MPa]	Selected Value [MPa]
Dessert (Arid)	-4.0 to -5.0 MPa [6]	-4.5 MPa
Grasslands/shrublands (Semi-Arid)	-1.4 to -5.0 ¹⁰ [8]	-3.2 MPa
Temperate Forest	-0.44 (Fall), -1.19 (Winter), -0.58 (Spring), -1.42 (Early Summer), -1.81 (Summer) [12]	-1.09 MPa
Boreal Forest (Arboreal)	-0.83 (Fall), -1.20 (Winter), -0.55 (Spring), -1.61 (Early Summer), -3.36 (Summer) [12]	-1.51 MPa
Tropical Rain Forest	-1.57 MPa to 0.00 MPa [27]	-0.79

Table 5: Environment Water Potential Data

Parameter	Symbol	Value	Units	Source and Specification
Half-Saturation constants ¹¹	K_e	-	mM	Enzyme
	K_e - Phenol oxidase	0.89	mM	[20]
	K_e - Phosphatase	0.94	mM	[19]
	K_e - Peroxidase	0.7475	mM	[1]
	K_e - Cellobiohydrolase	13.90	mM	[23]
Hyphal tip elongation rate	$v(T, \psi)$	Various	$\frac{mm}{day}$	[28] Species, ψ , T
Soil moisture coefficient	S_M	$S_M = \alpha_2 - \lambda \log_{10}(-\psi)$	N/A	[7] ψ
Soil temperature coefficient	S_T	$\log_{10}(S_T) = \frac{T-25}{10} \log_{10}(Q)$	N/A	[7] T

Table 6: Environmental & Parameter Table

C Appendix Figures

Figures here

¹⁰Measurements taken November through January at 100 cm soil depth.

¹¹Also called Michaelis Constant.

References

- [1] B. Chance, *Journal of Biological Chemistry* **1943**, 151, 553–577.
- [2] R. Whittaker, *Communities and ecosystems*, Macmillan, **1970**.
- [3] H. Parnas, *Soil Biology and Biochemistry* **1975**, 7, 161–169.
- [4] L. Edelstein, *Journal of Theoretical Biology* **1982**, 98, 679–701.
- [5] L. Edelstein, L. Segel, *Journal of Theoretical Biology* **1983**, 104, 187–210.
- [6] E. T. Nilsen, M. R. Sharifi, P. W. Rundel, W. M. Jarrell, R. A. Virginia, *Ecology* **1983**, 64, 1381–1393.
- [7] D. L. Moorhead, J. F. Reynolds, *Ecological Modelling* **1991**, 56, 197–219.
- [8] D. Peláez, R. Distel, R. Bóo, O. Elia, M. Mayor, *Journal of Arid Environments* **1994**, 27, 71–78.
- [9] O. AndreÂna, J. Balandreaub in, **1999**.
- [10] P. Gervais, C. Abadie, P. Molin, *Journal of Scientific & Industrial Research* **1999**, 58, 671–677.
- [11] L. Boddy, *FEMS Microbiology Ecology* **2000**, 31, 185–194.
- [12] D. B. Zobel, S. C. Garkoti, S. P. Singh, A. Tewari, C. M. S. Negi, *Current Science* **2001**, 80, 774–779.
- [13] J. P. Schimel, M. N. Weintraub, *Soil Biology and Biochemistry* **2003**, 35, 549–563.
- [14] D. L. Moorhead, R. L. Sinsabaugh, *Ecological Monographs* **2006**, 76, 151–174.
- [15] Y. Toljander, B. Lindahl, L. Holmer, N. Högborg, *Oecologia* **2006**, 148, 625–31.
- [16] G. Steinberg, *Eukaryotic Cell* **2007**, 6, 351–360.
- [17] A. Schnepf, T. Roose, P. Schweiger, *Journal of The Royal Society Interface* **2008**, 5, 773–784.
- [18] K. L. McGuire, K. K. Treseder, *Soil Biology and Biochemistry* **2010**, 42, 529–535.
- [19] P. Nannipieri, L. Giagnoni, L. Landi, G. Renella in, **2011**, pp. 215–243.
- [20] E. A. Davidson, S. Samanta, S. S. Caramori, K. Savage, *Global Change Biology* **2012**, 18, 371–384.
- [21] F. Segato, A. R. L. Damásio, R. C. de Lucas, F. M. Squina, R. A. Prade, *Microbiology and Molecular Biology Reviews* **2014**, 78, 588–613.
- [22] J. A. M. Moore, J. Jiang, W. M. Post, A. T. Classen, *Ecosphere* **2015**, 6, art29.
- [23] B. S. Razavi, E. Blagodatskaya, Y. Kuzyakov, *Frontiers in microbiology* **2015**, 6, 1126–1126.
- [24] L. Mucina, *Biome: evolution of a crucial ecological and biogeographical concept*, New Phytologist, **2018**.
- [25] F. Z. Abkenar, A. Rasoulzadeh, A. Asghari, *Bragantia* **2019**, 78, 119–130.
- [26] H. Du, T.-B.-T. Tran, P. Perré, *Journal of Theoretical Biology* **2019**, 470, 90–100.
- [27] S. J. Kupers, B. M. J. Engelbrecht, A. Hernández, S. J. Wright, C. Wirth, N. Rüger, *Journal of Ecology* **2019**, 107, 860–874.

- [28] D. S. Maynard, M. A. Bradford, K. R. Covey, D. Lindner, J. Glaeser, D. A. Talbert, P. J. Tinker, D. M. Walker, T. W. Crowther, *Nature Microbiology* **2019**, 4, 846–853.
- [29] S. Paton, Barro Colorado Island, Clearing *temperature, manual*, **2019**.
- [30] S. N. Zaz, S. A. Romshoo, R. T. Krishnamoorthy, Y. Viswanadhapalli, *Atmospheric Chemistry and Physics* **2019**, 19, 15–37.
- [31] N. Lustenhouwer, D. S. Maynard, M. A. Bradford, D. L. Lindner, B. Oberle, A. E. Zanne, T. W. Crowther, *Proceedings of the National Academy of Sciences* **2020**, 117, 11551–11558.