II. Cycles beyond 1 for the $qn + 1$ map: finiteness for each length, and an explicit bound on q for a constant odd element

Trevor Hackman

Abstract

For the $qn + 1$ map, for each length at least 2, only finitely many odd integers q have a cycle of that length. The argument uses Ridout's theorem in the form that allows prime factors of the denominator outside a constant finite set, and it gives no bound on q. If $|2^{r/\ell} - p/d| > c\, d^{-1.3247}$ for each root $2^{r/\ell}$, for every odd positive integer x and every large denominator $d = x \cdot 2^j$ or $d = \ell x \cdot 2^j$, then the bound on q depends only on the length.

If a constant odd positive integer is the starting value of a cycle with two or three distinct odd numbers, then q lies below an explicit bound depending only on that integer. For three odd numbers, once q meets that bound, a lower bound and an upper bound cannot hold at the same time. Bauer and Bennett (2018) give the exponent $\mu \approx 1.6149$, which lies just below the golden ratio, the threshold at which these bounds conflict. Every smaller exponent is a finite check. For two odd numbers the same conflict uses their exponent 1.465 for the square root of 2 (2002). Evaluating their Theorem 1.5 at the exponent 1.617, together with a search through $q < 2^{160{,}000}$ and a digit scan of $\sqrt{2}$, shows that every other cycle with two odd numbers has $q \geq 2^{1{,}071{,}999{,}971}$ and each odd element greater than $2^{17{,}631{,}663}$. When the constant integer is 1 and the cycle has three odd numbers, the bound is $q < 2^{17990}$, and the search in the companion paper already rules out every such cycle.

Contents

1 Introduction **2**
Conjecture 1 2

2 Definitions and notation **3**

3 Ineffective finiteness **5**
Theorem 2 7

4 The integer $Q_\ell(x)$ **8**
Theorem 3 8
Proposition 3 11

5 Cube roots with a constant odd part **13**

6 The integer $Q_3(x)$ **17**

7 The integer $Q_2(x)$ **20**

8 Cycles of length 2 **21**

Proposition 10 25
Proposition 12 27

9 Longer cycles **29**

10 A heuristic **30**

11 Open problems **31**

A Restrictions on the denominator of any cycle **31**

B Finiteness past Theorem 2 **35**

C Computations **40**

1 Introduction

This paper studies cycles of the $qn + 1$ map for any starting value. It is preceded by the narrower paper [6], which studies cycles with starting value 1.

Fix an odd integer $q \geq 3$. For an odd positive integer n, let v be the largest integer such that 2^v divides $qn + 1$. The map C sends n to

$$C(n) = \frac{qn + 1}{2^v}.$$

The orbit of n is the sequence n, $C(n)$, $C(C(n))$, ... When $q = 3$, C is the odd-to-odd form of the Collatz map, also called the Syracuse function, and the Collatz conjecture, posed by Lothar Collatz in 1937 and still open, asserts that every orbit eventually reaches 1. For $q = 3$, heuristic arguments suggest that almost all orbits eventually reach 1. However, for $q \geq 5$, heuristic arguments suggest that almost all orbits grow without bound and that cycles are rare, although no orbit has been proved to diverge.

The known cycles are these. For each Mersenne number, the map has the cycle $1 \to 1$. No other odd $q \geq 3$ has a cycle of length 1. For $q = 5$ there is $1 \to 3 \to 1$, and also $13 \to 33 \to 83 \to 13$ and $17 \to 43 \to 27 \to 17$. For $q = 181$ there is $27 \to 611 \to 27$ and $35 \to 99 \to 35$. Crandall recorded $27 \to 611$ from the identity $2^{15} - 181^2 = 7$.

Conjecture 1. These are all the cycles of C. For every odd $q \geq 3$, every cycle is either $1 \to 1$ with q a Mersenne number, or one of the five cycles at $q = 5$ and $q = 181$ listed above. In particular, if $q \geq 7$ is neither 181 nor a Mersenne number, then C has no cycle.

There is then no cycle with starting value 1 other than $1 \to 1$ for a Mersenne number and $1 \to 3 \to 1$ for $q = 5$, which is the conjecture of Santos (2020), and for $q = 3$ there is no cycle other than $1 \to 1$, which is part of the Collatz conjecture. Crandall (1978) conjectured that for every odd integer $q > 3$, some positive integer has an orbit that does not reach 1 [4]. For $q > 3$ he expected an orbit whose odd terms tend to infinity. The cycles at $q = 5$ and $q = 181$ do not have odd terms that tend to infinity. Conjecture 1 does not assert that the odd terms of any orbit tend to infinity. The searches with starting value 1 are in the companion paper [6]. A search over cycles with no element

held constant, and a search at a constant q, are in Appendix C. Section 10 gives a heuristic for the expected number of cycles.

Theorem 2. For each $\ell \geq 2$, only finitely many odd q have a cycle of C of length ℓ.

Theorem 2 is ineffective: it rests on the general form of Ridout's theorem [8] in Section 3, which allows prime factors of the denominator outside a constant finite set, and gives no bound on q, and no bound depending on a constant starting value. (A finiteness result is called effective if its proof gives an explicit bound on any exception, so that a finite computation can find them all, and ineffective if it gives no such bound.) The case of starting value 1 is Theorem 3 of the companion paper, and that proof uses only the S-unit form of Ridout's theorem. The proof of Theorem 2 is in Section 3.

Outline of Theorem 3. Fix a length $\ell \geq 2$ and an odd starting value x. Suppose that, for every integer r with $1 \leq r \leq \ell - 1$, $2^{r/\ell}$ has a restricted irrationality exponent below the golden ratio. Then any q for which there is a cycle of this length with starting value x lies below a bound computed from that exponent and from x. For length 2 the restricted irrationality exponent is 1.465, from Corollary 1.4 of Bauer and Bennett (2002). For length 3 it is about 1.6149, from Bauer and Bennett (2018). The precise statement is in Section 4.

Proposition 3 gives, for each length, a bound on q that does not depend on a starting value. The hypothesis is the lower bound in that proposition, of the shape $c \cdot d^{-\mu}$ with μ at most 1.3247, for every root $2^{r/\ell}$ and every odd positive integer in the denominator.

For $x = 1$ and length 3, Appendix C gives $Q_3(1) < 2^{17990}$. Any such cycle has one candidate q for each total number of factors of 2 removed, so it is enough to check those totals below 53970. The companion paper checks every such total up to 200,000 and finds no cycle. That search includes the totals below 53970, and Theorem 1 of [6] records that no such cycle exists. Length 2 with starting value 1 is Proposition A.1 of [6]: the only such q is 5.

For a general x the bound does not rule a cycle out. At $q = 181$ the cycles $27 \to 611 \to 27$ and $35 \to 99 \to 35$ both occur. Appendix C records $181 < Q_2(27)$ and $181 < Q_2(35)$.

Section 8 takes up cycles of two odd numbers with no element held constant in advance. Every such cycle other than $1 \to 3 \to 1$, $27 \to 611 \to 27$ and $35 \to 99 \to 35$ has $q \geq 2^{1,071,999,971}$ and every odd number on it greater than $2^{17,631,663}$.

Length 4 with an odd total number of removed factors of 2 remains open, as in the companion paper: the roots are $2^{1/4}$ and $2^{3/4}$, and a constant odd part does not supply an exponent below the golden ratio for them. Corollary 14 covers the longer classes whose roots are only $\sqrt{2}$, or only the cube roots of 2.

2 Definitions and notation

Throughout, $q \geq 3$ is an odd integer and C is the map defined in Section 1.

Mersenne numbers. Let $e \geq 2$ be an integer. A Mersenne number is the integer $2^e - 1$.

The 2-adic valuation. For a nonzero integer n, $v_2(n)$ is the largest integer v such that 2^v divides n.

Cycles. A cycle of C of length ℓ consists of ℓ distinct odd positive integers. The starting value is the smallest element x_0. Applying C repeatedly, starting at x_0, produces

$$x_0 \to x_1 \to \cdots \to x_{\ell-1} \to x_0,$$

where the indices are read modulo ℓ.

Halvings. The halving counts $k_1, \ldots, k_\ell$ of the cycle are defined by

$$qx_{i-1} + 1 = 2^{k_i} x_i \qquad (1 \leq i \leq \ell,\ x_\ell = x_0),$$

so $k_i = v_2(q \cdot x_{i-1} + 1)$. The total number of halvings is $h = k_1 + \cdots + k_\ell$, and $S_i = k_1 + \cdots + k_i$.

The δ window. Let $\ell \geq 2$ be an integer. On a cycle of length ℓ with total number of halvings h, the number δ is the fractional part of $2^{h/\ell}$. The integer q equals $\lfloor 2^{h/\ell} \rfloor$, so $2^{h/\ell} = q + \delta$ and $0 < \delta < 1$. The δ window is a lower bound and an upper bound on δ.

The restricted irrationality exponent. Let θ be an irrational real number. A real number μ is a restricted irrationality exponent of θ when there exists an explicit constant real number $c > 0$ and there exists an explicit constant integer $j_0 \geq 0$ such that, for every integer p and for every integer $j \geq j_0$,

$$\left|\theta - \frac{p}{2^j}\right| > c \cdot 2^{-\mu j}.$$

The golden ratio. The golden ratio is $\varphi = (1 + \sqrt{5})/2$.

Divides. The symbol $\mid$ means divides, and the symbol $\nmid$ means does not divide. Let a and b be integers, with $a \neq 0$. Write $a \mid b$ when there exists an integer k such that $b = ak$. Write $a \nmid b$ when no such integer k exists.

Big O. Big O is the Bachmann–Landau symbol O. Let f and g be real-valued functions of a real variable t, with $g(t) \neq 0$ for every large t. Write $f(t) = O(g(t))$ when there exists a constant real number $C > 0$, independent of t, and there exists a constant real number t_0, such that $|f(t)| \leq C\,|g(t)|$ for every $t \geq t_0$. The expression $O(g(t))$ stands for a real number of absolute value at most $C\,|g(t)|$. An equation $A = B + O(g(t))$ means $A - B = O(g(t))$.

The plastic number. The plastic number is the real root of $t^3 - t - 1 = 0$. The plastic number equals

$$\sqrt[3]{\frac{9 + \sqrt{69}}{18}} + \sqrt[3]{\frac{9 - \sqrt{69}}{18}}.$$

The constants μ_3, γ_3, and $c_3(s')$. Let $N = 16166467234$. The real number μ_3 is $3 \log_2 N/63$. The positive number γ_3 is $1 - \mu_3 + 1/\mu_3$. For a positive integer s', the constant $c_3(s')$ is $2^{-5}\, N^{-79/63}/s'$.

3 Ineffective finiteness

Ridout's theorem (1958). Let α be an algebraic irrational, let S be a finite set of primes, and let $\varepsilon > 0$. Only finitely many reduced fractions p/Q with $Q > 0$ have every prime factor of Q in S and satisfy

$$\left|\alpha - \frac{p}{Q}\right| < Q^{-1-\varepsilon}.$$

The same theorem has the general form: writing $|n|_p = p^{-v_p(n)}$ for a nonzero integer n, only finitely many rationals P/Q in lowest terms satisfy

$$Q|Q\alpha - P| \prod_{p \in S} |Q|_p < \max(|P|, Q)^{-\varepsilon}.$$

The first statement is the S-unit form. If every prime factor of Q lies in S, then $\prod_{p\in S}|Q|_p = 1/Q$ and the left side of the general form equals $|Q\alpha - P|$. Whenever $|P| \le Q$, this is the first form. The general form allows prime factors of Q outside S. Such a prime does not appear in the product, and it contributes its full size to the height $\max(|P|, Q)$. Neither form gives a bound on the exceptional fractions. Since α is irrational, the finitely many fractions that meet the general form for a constant α lie at a positive distance from α, and these distances have a positive minimum. This is why Theorem 2, which uses the general form in place of a restricted irrationality exponent, gives no bound on q.

Lemma 1 (the window for any cycle). Let $a = x_0 \to x_1 \to \cdots \to x_{\ell-1} \to a$ be a cycle of length $\ell \ge 2$, with starting value a, and let $\sigma = \sum_{i=0}^{\ell-1} 1/x_i$. Then

$$2^h = \prod_{i=0}^{\ell-1} \left(q + \frac{1}{x_i}\right),$$

so $q^\ell < 2^h < (q+1)^\ell$, and in particular $\ell \nmid h$ and $q = \lfloor 2^{h/\ell} \rfloor$. Writing $2^{h/\ell} = q + \delta$,

$$e^{-\ell/q} \le \frac{\ell\,\delta}{\sigma} \le e^{\ell/q}.$$

Proof. Multiplying $2^{k_i} x_i = q\,x_{i-1} + 1$ around the cycle gives the product. Each factor lies in $(q,\, q+1]$, with equality to $q+1$ only for an element equal to 1. At most one element is 1, and $\ell \ge 2$, so the product is strictly larger than q^ℓ and strictly smaller than $(q+1)^\ell$. A power of 2 strictly between consecutive ℓ-th powers is not an ℓ-th power, so $\ell \nmid h$, and $q = \lfloor 2^{h/\ell} \rfloor$. In particular $0 < \delta < 1$.

The binomial theorem gives

$$\ell q^{\ell-1}\delta \le (q+\delta)^\ell - q^\ell \le \ell q^{\ell-1}\delta\Big(1 + \frac{\delta}{q}\Big)^{\ell-1}.$$

Dividing the binomial expansion by $\ell q^{\ell-1}\delta$, the coefficient of $(\delta/q)^k$ is $\binom{\ell}{k+1}/\ell = \binom{\ell-1}{k}/(k+1) \le \binom{\ell-1}{k}$, and $\sum_{k\ge 0}\binom{\ell-1}{k}(\delta/q)^k = (1+\delta/q)^{\ell-1}$. Since $\delta < 1$, this factor is at most $e^{(\ell-1)/q} \le e^{\ell/q}$.

The same difference equals $\sum_{j\ge1} e_j q^{\ell-j}$, where e_j is the j-th elementary symmetric function of the reciprocals $1/x_i$. Thus $e_1 = \sigma$, and the right side is at least $q^{\ell-1}\sigma$. In the expansion of σ^j, each product of j distinct reciprocals occurs $j!$ times, so $e_j \le \sigma^j/j!$. Hence the right side is at most

$$\sum_{j\ge1} \frac{\sigma^j}{j!}\, q^{\ell-j} = q^{\ell-1} \sum_{m\ge0} \frac{\sigma^{m+1}}{(m+1)!\, q^m} \le q^{\ell-1}\sigma\, e^{\sigma/q},$$

since $(m+1)! \ge m!$. Every element is at least 1, so $\sigma \le \ell$ and $e^{\sigma/q} \le e^{\ell/q}$. Comparing the two sides gives $\ell\delta \le \sigma\, e^{\ell/q}$ and $\ell\delta\, e^{\ell/q} \ge \sigma$. □

Lemma 2 (three approximations). In the setting of Lemma 1, write $h = \ell m + r$ with $0 < r < \ell$. Let $\theta_i = 2^{ir/\ell}$ for $1 \le i \le \ell-1$, let $\theta = \theta_1$, and let $y = \min(x_1, \dots, x_{\ell-1})$. Suppose $q \ge 4\ell$. Then $2^m > q/2$, and:

1. $|\theta - q/2^m| = \delta/2^m$.
2. For $1 \le i \le \ell-1$, the rational number $F_i = x_i/(a \cdot 2^{im-S_i})$ satisfies $|\theta_i - F_i| \le 2i\theta_i/(aq)$.
3. The rational number $W = (\ell aq+1)/(\ell a \cdot 2^m)$ satisfies $|\theta - W| \le 2(1/y + 2\ell/q)/q$.

Proof. Since $2^m\theta = q + \delta$ and $\theta = 2^{r/\ell} < 2$, we have $2^m = (q+\delta)/\theta > q/2$, and part 1 follows.

The relation $q\, x_{i-1} + 1 = 2^{k_i} x_i$ gives $x_i\, 2^{S_i} = a \prod_{u<i}(q + 1/x_u)$ by induction on i, so

$$F_i = \frac{1}{2^{im}} \prod_{u<i} \Big(q + \frac{1}{x_u}\Big), \qquad \theta_i = \frac{(q+\delta)^i}{2^{im}}.$$

Lemma 1 gives $\delta \le (\sigma/\ell)\, e^{\ell/q}$. Every element is at least a, so $\sigma \le \ell/a$ and $\delta \le c/a$ with $c = e^{\ell/q} \le e^{1/4}$. Each factor $q + 1/x_u$ and the number $q+\delta$ therefore lie in $[q,\, q + c/a]$, and both numerators lie in $[q^i,\, (q+c/a)^i]$. Their difference is at most

$$\Big(q + \frac{c}{a}\Big)^i - q^i = q^i \Big(\Big(1 + \frac{c}{aq}\Big)^i - 1\Big) \le \frac{ic}{a}\, q^{i-1}\, e^{ic/(aq)},$$

where $e^z - 1 \le z\, e^z$. Dividing by $2^{im} = (q+\delta)^i/\theta_i \ge q^i/\theta_i$ gives $|\theta_i - F_i| \le (ic\theta_i/(aq))\, e^{ic/(aq)}$. Here $i \le \ell - 1$ and $aq \ge 4\ell$, so $ic/(aq) < c/4$. For $0 < z \le 1/3$ and $n \ge 2$, we have $n! \ge 2^{n-1}$, so $z^n/n! \le z^n/2^{n-1}$ and $\sum_{n\ge2} z^n/2^{n-1} = z^2/(2-z)$. Thus $e^{1/4} < 1 + 1/4 + (1/16)/(7/4) = 9/7 < 4/3$ and $e^{1/3} < 1 + 1/3 + 1/15 = 7/5$. Since $c \le e^{1/4} < 4/3$, we have $c/4 < 1/3$ and $c\, e^{c/4} < (4/3)\cdot(7/5) = 28/15 < 2$. Thus $|\theta_i - F_i| \le 2i\theta_i/(aq)$.

For part 3,

$$\big|\theta - W\big| = \frac{1}{2^m}\Big|\delta - \frac{1}{\ell a}\Big|.$$

Write $\delta - 1/(\ell a) = (\sigma - 1/a)/\ell + (\delta - \sigma/\ell)$. The other elements are at least y, so $0 \le \sigma - 1/a \le (\ell-1)/y$ and $(\sigma - 1/a)/\ell \le 1/y$. From the two bounds in Lemma 1, with $z = \ell/q$,

$$\frac{\sigma}{\ell} - \delta \le \frac{\sigma}{\ell}(1 - e^{-z}), \qquad \delta - \frac{\sigma}{\ell} \le \frac{\sigma}{\ell}(e^z - 1).$$

The inequality $1 - e^{-z} \le e^z - 1 \le z\, e^z$ and $\sigma \le \ell$ give $|\delta - \sigma/\ell| \le (\ell/q)\, e^{\ell/q}$. Since $q \ge 4\ell$, this is at most $(\ell/q)\, e^{1/4} < 2\ell/q$. Therefore $|\delta - 1/(\ell a)| \le 1/y + 2\ell/q$. Dividing by $2^m > q/2$ gives part 3. □

Theorem 2. For each $\ell \geq 2$, only finitely many odd q have a cycle of C of length ℓ.

Proof of Theorem 2. Fix $\ell \geq 2$, and set $\varepsilon = 1/(10\ell)$. Let S consist of 2 and the prime factors of ℓ, and let Θ be the set of real numbers $2^{t/\ell}$ with $1 \leq t \leq \ell(\ell-1)$ and $\ell \nmid t$. Each element of Θ is an algebraic irrational. Apply the general form of Ridout's theorem to each $\theta' \in \Theta$, with this S and ε. Only finitely many fractions meet the inequality, and none equals θ', so there is an $\eta > 0$ such that every such fraction satisfies $|\theta' - P/d| \geq \eta$. The value of η is not provided by Ridout's theorem. The odd $q < 4\ell$ are finite in number, so take an odd $q \geq 4\ell$ with a cycle of length ℓ, in the notation of Lemmas 1 and 2. Then $\theta \in \Theta$. Each θ_i either lies in Θ or is a power of 2.

Consider the fraction $q/2^m$, which is in lowest terms. Its left side in the general form equals $2^m \cdot \delta \cdot 2^{-m} = \delta$, since $|2^m|_p = 1$ for every odd prime p. Also $\max(q, 2^m) < q+1$, because $2^m = (q+\delta)/\theta < (q+1)/\theta < q+1$. If $\delta < (q+1)^{-\varepsilon}$, then $\delta < \max(q, 2^m)^{-\varepsilon}$, so Ridout's inequality applies and $\delta/2^m \geq \eta$. Combined with $2^m > q/2$ this yields $q < 2/\eta$.

Now assume $\delta \geq (q+1)^{-\varepsilon}$. Lemma 1 gives $\delta \leq e^{\ell/q}/a \leq e^{1/4}/a$, so $a < 2(q+1)^{\varepsilon}$.

Suppose first that $y \leq q^{1-2\varepsilon}$, and choose i with $x_i = y$. If ℓ divides ir, then $\theta_i = 2^s$ for some integer $s \geq 1$. This power is not equal to F_i. Otherwise $x_i = a \cdot 2^{im - S_i + s}$; the right side is an odd positive integer only if the exponent of 2 vanishes, which forces $x_i = a$, contradicting distinctness. Let $e = \max(im - S_i, 0)$. In lowest terms the denominator of F_i divides $a \cdot 2^e$, so $|\theta_i - F_i| \geq 1/(a \cdot 2^e)$. Lemma 2 then gives $2^e \geq q/(2i\theta_i)$. If $e = 0$, then $q \leq 2i\theta_i$. Here $i \leq \ell - 1$ and $\theta_i < 2^{\ell-1}$, so $q < (\ell-1)\,2^\ell$. If $e \geq 1$, the same lemma and $q \geq 4\ell$ give $|\theta_i - F_i| \leq 2(\ell-1)\theta_i/(4\ell) < \theta_i/2$, hence $F_i > \theta_i/2$. Therefore

$$y = F_i \cdot a \cdot 2^e > \frac{\theta_i}{2} \cdot a \cdot \frac{q}{2i\theta_i} = \frac{aq}{4i} \geq \frac{q}{4(\ell-1)}.$$

Once $q > (4(\ell-1))^{5\ell}$, the right side exceeds $q^{1-2\varepsilon}$, because $1/(2\varepsilon) = 5\ell$. This contradicts $y \leq q^{1-2\varepsilon}$.

If instead ℓ does not divide ir, then $\theta_i \in \Theta$. Write $F_i = P/d$ in lowest terms. The odd part of d divides a, so $d \cdot |d|_2 \leq a$. Every factor $|d|_p$ is at most 1 and $2 \in S$, so the left side of the general form is at most

$$d^2 \cdot |\theta_i - F_i| \cdot |d|_2 \leq \frac{2i\theta_i\, d}{q}.$$

As above, $|\theta_i - F_i| < \theta_i/2$, so $F_i > \theta_i/2 > 1/2$ and $F_i < (3/2)\theta_i < 2^\ell$. If $im \geq S_i$, the fraction before reducing is $y/(a \cdot 2^{im - S_i})$, whose denominator equals $y/F_i < 2y$, and the numerator is y. If $im < S_i$, that denominator is a, so $d \leq a \leq y$, and the numerator is $F_i d < 2^\ell y$. In both cases $d < 2y$ and $\max(|P|, d) < 2^\ell y$. Also $2i\theta_i < (\ell-1)\,2^\ell$, so the left side is less than $(\ell-1)\,2^{\ell+1} y/q$. The right side is greater than $(2^\ell y)^{-\varepsilon}$. The comparison

$$(\ell-1)\,2^{\ell+1}\, y^{1+\varepsilon} < q \cdot 2^{-\ell\varepsilon}$$

therefore implies Ridout's inequality. It holds for every admissible y once

$$q^{\varepsilon + 2\varepsilon^2} > (\ell-1)\,2^{\ell+1+\ell\varepsilon},$$

because $y^{1+\varepsilon} \leq q^{(1-2\varepsilon)(1+\varepsilon)} = q^{1-\varepsilon-2\varepsilon^2}$. The right side depends only on ℓ. For every larger q, Ridout's theorem gives $|\theta_i - F_i| \geq \eta$. Lemma 2 gives $2i\theta_i/(aq) \geq \eta$, and $2i\theta_i < (\ell-1)\,2^\ell$ with $a \geq 1$, so $q < (\ell-1)\,2^\ell/\eta$.

Suppose instead that $y > q^{1-2\varepsilon}$. The numerator $N = \ell aq + 1$ is congruent to 1 modulo ℓa, so $\gcd(N, \ell a) = 1$. Any common factor of N and $\ell a \cdot 2^m$ is therefore a power of 2. Let $v = v_2(N)$ and $e = \max(m - v, 0)$. In lowest terms, $W = P/d$ with $d = \ell a \cdot 2^e$ and $P \le N$. The product formula gives $\prod_p |\ell|_p = 1/\ell$, and every prime factor of ℓ lies in S, so $\prod_{p\in S} |\ell|_p = 1/\ell$. Since $|a|_p \le 1$,

$$\prod_{p\in S} |d|_p \le \frac{2^{-e}}{\ell}.$$

The left side of the general form is therefore at most

$$d^2 \cdot |\theta - W| \cdot \frac{2^{-e}}{\ell} = \ell a^2\, 2^e \cdot |\theta - W| \le \ell a^2\, 2^m \cdot |\theta - W|.$$

Lemma 2 and $2^m < q + 1$ bound this by $2\ell a^2(1 + 1/q)(1/y + 2\ell/q)$. Insert $1/y < q^{-1+2\varepsilon}$ and $a < 2(q + 1)^\varepsilon$. For $q \ge 4\ell$ we have $1 + 1/q < 2$, $1 + 2\ell\, q^{-2\varepsilon} \le 1 + 2\ell$, and $(q + 1)^{2\varepsilon} < (2q)^{2\varepsilon}$, so the left side is less than

$$16\ell(1 + 2\ell)\, 2^{2\varepsilon}\, q^{-1+4\varepsilon}.$$

Since $P \le \ell aq + 1$ and $d \le \ell a \cdot 2^m < \ell a(q + 1)$, the height satisfies $\max(|P|, d) < \ell a(q + 1) + 1 < 4\ell(q + 1)^{1+\varepsilon}$, so the right side is greater than $(4\ell)^{-\varepsilon}(q + 1)^{-\varepsilon(1+\varepsilon)} > (4\ell)^{-\varepsilon}(2q)^{-\varepsilon(1+\varepsilon)}$. The quotient of these two bounds is

$$16\ell(1 + 2\ell)\, 2^{2\varepsilon}(4\ell)^\varepsilon\, 2^{\varepsilon(1+\varepsilon)}\, q^{-1+5\varepsilon+\varepsilon^2}.$$

The exponent equals $-1 + 5/(10\ell) + 1/(100\ell^2) \le -1 + 1/4 + 1/400 < -1/2$. The quotient tends to 0 as $q \to \infty$, so Ridout's inequality holds for all q above a bound depending only on ℓ. Then $|\theta - W| \ge \eta$. Lemma 2 and $1/y \le 1$ give $|\theta - W| \le 2(1 + 2\ell)/q$, hence $q \le (2 + 4\ell)/\eta$.

In every case q is at most a number depending only on ℓ and η. □

The argument does not produce a bound on q, and it does not produce a bound depending on a constant odd start. Fix an integer $V \ge 2$. Each length ℓ with $2 \le \ell \le V$ contributes finitely many odd q, and for each such pair the inequalities $q^\ell < 2^h < (q + 1)^\ell$ leave finitely many h and finitely many compositions of h into ℓ positive parts. Thus only finitely many cycles have length at most V.

Theorem 2 fails for $\ell = 1$. A Mersenne number q gives the cycle $1 \to 1$, and the same identity becomes $2^h = q + 1$. The constant η is not explicit, so neither is the bound on q. An effective bound would require an effective inequality of the same shape for every denominator whose odd part is at most a constant times q^ε.

4 The integer $Q_\ell(x)$

Theorem 3. Let $\ell \ge 2$ be an integer and let x be an odd positive integer. Let μ be a constant real number with $1 < \mu < \varphi$. Let c and c_* be constant positive real numbers, and let j_0 be a positive integer. Suppose that for every $r \in \{1, \dots, \ell - 1\}$, every integer p, and every integer $j \ge j_0$,

$$\left|2^{r/\ell} - \frac{p}{x\, 2^j}\right| > c \cdot 2^{-\mu j}$$

and

$$\left|2^{r/\ell} - \frac{p}{\ell x\, 2^j}\right| > c_* \cdot 2^{-\mu j}.$$

Let B be the positive real number $\max(\ell - 1,\ 2(\ell-2) + 1/x)$. Let γ be the positive real number $1 - \mu + 1/\mu$, and let A and Q_* be the positive real numbers

$$A = \frac{x}{2}\, 2^{1/\ell} \left(\frac{c}{2^{(\ell-1)/\ell}\, B} \right)^{1/\mu}, \qquad Q_* = \left(\frac{(\ell-1)/(\ell A) + \ell}{c_*\, 2^{1-\mu}} \right)^{1/\gamma}.$$

For each such r, let d_r be the positive real number equal to the distance from $x\, 2^{r/\ell}$ to the nearest integer. For each such r, each integer j with $0 \le j < j_0$, and each odd positive integer $y < 3x \cdot 2^{j_0}$, let ρ be the nonnegative real number

$$\rho = \left| \frac{y}{x\, 2^j} \cdot 2^{-r/\ell} - 1 \right|.$$

Whenever $\rho > 0$, the corresponding value B/ρ is a positive real number, and so is $B \cdot 2^{r/\ell} \cdot x/d_r$. Let Q_0 be the maximum of those finitely many positive real numbers. If ℓ is composite, that maximum also includes the positive real number Bx. Let Q_e be the least positive integer such that every integer $q \ge Q_e$ satisfies $q \ge 4\ell$ and $e^{\ell/q} \le q^{1-1/\mu}$. If ℓ is prime, let Q_A be the positive integer 1. If ℓ is composite, let Q_A be the least positive integer such that every integer $q \ge Q_A$ satisfies $q/(2B) > A\, q^{1/\mu}$. Let $Q_\ell(x;\, \mu, c, c_*, j_0)$ be the positive real number equal to the maximum of 2^{j_0}, of Q_0, of Q_*, of Q_e, and of Q_A. If x is the starting value of a cycle of length ℓ, then $q < Q_\ell(x;\, \mu, c, c_*, j_0)$.

Proof of Theorem 3. Let $x = x_0 \to x_1 \to \cdots \to x_{\ell-1} \to x$ be a cycle of length ℓ, with each x_i an odd positive integer. Lemma 1 gives $q = \lfloor 2^{h/\ell} \rfloor$ and $\ell \nmid h$. Write $2^{h/\ell} = q + \delta$ with δ the fractional part, so $0 < \delta < 1$. Let m be the nonnegative integer and r the integer with $0 < r < \ell$ and $h = \ell m + r$. If $q < 2^{j_0}$ or $q < Q_e$ or $q < Q_A$, there is nothing to prove. Assume $q \ge 2^{j_0}$, $q \ge Q_e$ and $q \ge Q_A$. Then $q \ge 4\ell$ and $e^{\ell/q} \le q^{1-1/\mu}$. If ℓ is composite, then $q/(2B) > A\, q^{1/\mu}$. Since $1/x \le 1$, we have $B \le 2\ell - 3$, and therefore $2B < 4\ell \le q$.

For each integer i with $1 \le i \le \ell - 1$, multiplying $qx_u + 1 = 2^{k_{u+1}} x_{u+1}$ for $0 \le u < i$ gives

$$x_i\, 2^{S_i} = x \prod_{u<i} \Big(q + \frac{1}{x_u} \Big).$$

Define the positive real numbers ε_i by $\varepsilon_1 = 1/(qx)$ and

$$\varepsilon_i = \varepsilon_{i-1} + \frac{1 + \varepsilon_{i-1}}{q\, x_{i-1}} \qquad (i \ge 2).$$

Then $x_i\, 2^{S_i} = x\, q^i (1 + \varepsilon_i)$: the case $i = 1$ is $qx + 1 = xq(1 + 1/(qx))$, and $qx_{i-1} + 1 = 2^{k_i} x_i$ multiplies by $q + 1/x_{i-1}$. The bound $q \ge 4\ell$ gives $B/q < 1$ and $\varepsilon_1 = 1/(qx) \le 1/q$. If $\varepsilon_{i-1} \le (1/x + 2(i-2))/q$, then that upper bound is at most $B/q < 1$, and $x_{i-1} \ge 1$, so

$$\varepsilon_i \le \varepsilon_{i-1} + \frac{2}{q} \le \frac{1/x + 2(i-1)}{q}.$$

Thus $\varepsilon_i \le B/q$ for every $i \le \ell - 1$. Let n_i be the integer $ih - \ell S_i$, so $n_i \equiv ir \pmod{\ell}$. Let j_i be the integer and r_i the integer with $n_i = \ell j_i + r_i$ and $0 \le r_i < \ell$. Let ρ_i be the real number given by $x_i = x\, 2^{n_i/\ell}(1 + \rho_i)$. Then

$$1 + \rho_i = \Big(\frac{q}{q + \delta} \Big)^i (1 + \varepsilon_i).$$

Since $0 < \delta < 1$, we have $(q/(q+\delta))^i < 1$ and $(q/(q+\delta))^i \ge (q/(q+1))^i \ge 1 - i/(q+1) > 1 - i/q$. Also $1 + \varepsilon_i \ge 1$, so $\rho_i < \varepsilon_i \le B/q$ and $\rho_i > -i/q$. Thus $|\rho_i| < B/q < 1/2$.

If ℓ is prime, then $r_i \ne 0$: here $1 \le i \le \ell - 1$ and $1 \le r \le \ell - 1$, so $ir \not\equiv 0 \pmod{\ell}$. If $r_i = 0$, then ℓ is composite and $x_i = x\,2^{j_i}(1 + \rho_i)$. If $j_i = 0$, then x_i and x are distinct odd positive integers, so $|x_i - x| \ge 2$. Hence $|\rho_i| \ge 2/x$ and $q < Bx/2$. If $j_i \le -1$, write $t = -j_i$. Then $x_i\,2^t = x(1 + \rho_i)$. The integers $x_i\,2^t$ and x are distinct, because x is odd, so $|x_i\,2^t - x| \ge 1$. Hence $|\rho_i| \ge 1/x$ and $q < Bx$. In both cases Bx is one of the numbers whose maximum is Q_0, so $q < Q_0$. If $1 \le j_i < j_0$, then $x\,2^{j_i}$ is even and x_i is odd, so $x\,2^{j_i}\,|\rho_i| \ge 1$. Hence $|\rho_i| \ge 2^{-j_i}/x$ and $q < Bx\,2^{j_i}$. Let r' be an integer with $1 \le r' \le \ell - 1$, and let $j' = j_0 - 1$. Let y be an odd positive integer at distance at most 1 from $x\,2^{j'}\,2^{r'/\ell}$. Since $2^{r'/\ell} < 2$, that real number is less than $x\,2^{j_0}$, so $y \le x\,2^{j_0} < 3x \cdot 2^{j_0}$. Let ρ be the positive real number $\bigl|y/(x\,2^{j'}) \cdot 2^{-r'/\ell} - 1\bigr|$. Since $2^{r'/\ell}$ is irrational, $\rho > 0$, and

$$\rho \le \frac{1}{x\,2^{j'}\,2^{r'/\ell}}.$$

Thus $B/\rho \ge Bx\,2^{j'}\,2^{r'/\ell} > Bx\,2^{j_0 - 1} \ge Bx\,2^{j_i} > q$. The value B/ρ is one of the numbers whose maximum is Q_0, so $q < Q_0$. If $j_i \ge j_0$, then $j_i \ge 1$, so $x\,2^{j_i}$ is even and x_i is odd. Hence $x\,2^{j_i}\,|\rho_i| \ge 1$. Combined with $|\rho_i| < B/q$ and $1 + \rho_i > 1/2$, this gives $x_i > q/(2B) > A\,q^{1/\mu}$.

Now suppose $r_i \in \{1, \ldots, \ell - 1\}$. If $j_i < 0$, write $t = -j_i$, so t is a positive integer and $x_i\,2^t/x = 2^{r_i/\ell}(1 + \rho_i)$. The left side is an integer divided by x, so

$$|\rho_i| \ge 2^{-r_i/\ell}\,\frac{d_{r_i}}{x}.$$

Combined with $|\rho_i| < B/q$, this gives $q < B \cdot 2^{r_i/\ell} \cdot x/d_{r_i}$. That value is one of the numbers whose maximum is Q_0, and $q < Q_0$. If $0 \le j_i < j_0$, then $x_i < 3x \cdot 2^{j_i}$, because $2^{r_i/\ell} < 2$ and $1 + \rho_i < 3/2$. The value $B/|\rho_i|$ is one of the numbers whose maximum is Q_0, and $|\rho_i| < B/q$ gives $q < Q_0$. If $j_i \ge j_0$, the hypothesis gives $2^{r_i/\ell}\,|\rho_i| > c \cdot 2^{-\mu j_i}$. Since $2^{r_i/\ell} \le 2^{(\ell-1)/\ell}$ and $2^{r_i/\ell}(1 + \rho_i) > 2^{1/\ell} \cdot (1/2)$,

$$2^{j_i} > \left(\frac{c\,q}{2^{(\ell-1)/\ell}\,B}\right)^{1/\mu}$$

and $x_i > A\,q^{1/\mu}$.

Every x_i with $i \ge 1$ is therefore greater than $A\,q^{1/\mu}$, or else $q < Q_0$. In the remaining case, the positive real number $\sigma = \sum_{i=0}^{\ell-1} 1/x_i$ satisfies $\sigma - 1/x \le (\ell - 1)\,A^{-1}q^{-1/\mu}$. Let z be the positive real number ℓ/q. Lemma 1 gives $e^{-z} \le \ell\delta/\sigma \le e^z$, so $|\ell\delta/\sigma - 1| \le e^z - 1$. The series for e^z gives $e^z - 1 \le z\,e^z$. Since $\sigma \le \ell$,

$$\bigl|\delta - \sigma/\ell\bigr| \le \frac{\ell}{q}\,e^{\ell/q}.$$

Hence

$$\bigl|\delta - 1/(\ell x)\bigr| \le \frac{\ell - 1}{\ell A}\,q^{-1/\mu} + \frac{\ell}{q}\,e^{\ell/q}.$$

The hypothesis $q \ge Q_e$ makes the second summand at most $\ell\,q^{-1/\mu}$, so

$$\bigl|\delta - 1/(\ell x)\bigr| \le \left(\frac{\ell - 1}{\ell A} + \ell\right) q^{-1/\mu}.$$

Since $q \geq 2^{j_0}$ and $2^{h/\ell} > q$, we have $m \geq j_0$. Also $2^m < (q+1)\,2^{-r/\ell} < 2q$. The window hypothesis, with denominator $\ell x \cdot 2^m$, gives

$$\bigl|\delta - 1/(\ell x)\bigr| = 2^m \Bigl|2^{r/\ell} - \frac{\ell x q + 1}{\ell x \cdot 2^m}\Bigr| > c_* \cdot 2^{(1-\mu)m} > c_*\, 2^{1-\mu}\, q^{1-\mu}.$$

Since $\gamma > 0$, these two bounds on $|\delta - 1/(\ell x)|$ conflict for every integer $q \geq Q_*$. The lower bound is strict, so the conflict includes $q = Q_*$. Thus $q < Q_*$, and in every case $q < Q_\ell(x;\, \mu, c, c_*, j_0)$. $\square$

For $\ell = 3$ we have $B = 2 + 1/x$ and $Q_A = 1$. Section 6 uses the smaller factor $1 + 1/x$. The proof of Corollary 5 shows that this factor bounds both $|\rho|$ of the other two elements once $q \geq 30$, and 30 lies below $Q_3(x)$.

For $\ell = 2$ we have $B = 1$ and $Q_A = 1$. Section 7 uses the larger factor $1 + 1/x$ in place of B, and records that the resulting maximum equals $Q_2(x)$.

Proposition 3. Let $\ell \geq 2$ be an integer. Let μ be a constant real number with $1 < \mu \leq 1.3247$. Let c be an explicit constant real number with $c > 0$, and let d_0 be an explicit constant positive integer. Suppose that for every integer r with $1 \leq r \leq \ell - 1$, every odd positive integer x, every nonnegative integer j, and every integer p, each positive integer $d \in \{x \cdot 2^j,\, \ell x \cdot 2^j\}$ with $d \geq d_0$ satisfies

$$\Bigl|2^{r/\ell} - \frac{p}{d}\Bigr| > c \cdot d^{-\mu}.$$

Then there is a positive real number Q, computed from ℓ, μ, c and d_0, such that every cycle of length ℓ has $q < Q$.

Proof. Let $\gamma = 1 - \mu + 1/\mu$ and let $\alpha = \mu(\mu - 1)/\gamma$. Since $1.3247 < 3/2 < \varphi$, we have $\mu < \varphi$, so $\mu^2 - \mu - 1 < 0$ and $\gamma > 0$. For a real number $t > 1$ the derivative of $t^3 - t - 1$ is $3t^2 - 1 > 0$, so $t^3 - t - 1$ increases with t. At $t = 1.3247 = 13247/10000$,

$$\Bigl(\frac{13247}{10000}\Bigr)^3 - \frac{13247}{10000} - 1 = -\frac{76579777}{10^{12}} < 0.$$

Thus $\mu^3 - \mu - 1 < 0$. Since $\mu > 0$, this is the same inequality as $\mu^2(\mu - 1) < 1 + \mu - \mu^2$. Dividing by μ gives $\mu(\mu - 1) < (1 + \mu - \mu^2)/\mu$. The right side equals γ, so $\mu(\mu - 1) < \gamma$. Since $\gamma > 0$, we have $\alpha < 1$. Let $\beta = \mu(\mu - 1)$. For $t > 1$ the real number $t(t - 1)$ increases with t, so

$$\beta \leq \frac{13247}{10000} \cdot \frac{3247}{10000} = \frac{43013009}{100000000} < \frac{1}{2}.$$

Let x be the starting value of a cycle of length ℓ, and write $h = \ell m + r$ with $0 < r < \ell$, as in Lemma 1. Then $2^m \cdot 2^{r/\ell} = q + \delta$, so $|2^{r/\ell} - q/2^m| = \delta/2^m$. If $2^m \geq d_0$, the hypothesis at the denominator 2^m gives $\delta/2^m > c \cdot 2^{-\mu m}$, so $\delta > c \cdot 2^{(1-\mu)m}$. Lemma 1 gives $\delta \leq e^{\ell/q}/x$. The proof of Lemma 2 records $e^{1/4} < 4/3$. If $q \geq 4\ell$ and $2^m \geq d_0$, then $e^{\ell/q} \leq e^{1/4}$ and

$$x < \frac{4}{3c} \cdot 2^{(\mu-1)m}.$$

Also $q > 2^{m + r/\ell} - 1$. For $m \geq 1$ this is greater than 2^{m-1}, so $\log_2 q > m - 1$.

If $2^m < d_0$, then $q < 2 \cdot 2^m < 2d_0$. Let Q_e be the least positive integer such that every integer $q \ge Q_e$ satisfies $q \ge 4\ell$ and $e^{\ell/q} \le q^{1-1/\mu}$, as in Theorem 3. Let $B_+ = 2\ell - 3$. For $\ell = 2$ let $B_- = 1$, and for $\ell \ge 3$ let $B_- = \max(\ell - 1,\, 2(\ell - 2))$. In Theorem 3 the positive real number B satisfies $B_- \le B \le B_+$. With the constant in that theorem equal to $cx^{-\mu}$, the factor x cancels $x^{-\mu}$ under the exponent $1/\mu$, and

$$A = \frac{1}{2} \cdot 2^{1/\ell} \left(\frac{c}{2^{(\ell-1)/\ell}\, B} \right)^{1/\mu}.$$

Set

$$A_+ = \frac{1}{2} \cdot 2^{1/\ell} \left(\frac{c}{2^{(\ell-1)/\ell}\, B_-} \right)^{1/\mu}, \qquad A_- = \frac{1}{2} \cdot 2^{1/\ell} \left(\frac{c}{2^{(\ell-1)/\ell}\, B_+} \right)^{1/\mu}.$$

Then $A_- \le A \le A_+$. If ℓ is prime, let $Q_A^\dagger = 1$. If ℓ is composite, let $Q_A^\dagger$ be the least positive integer such that every integer $q \ge Q_A^\dagger$ satisfies $q/(2B_+) > A_+\, q^{1/\mu}$. If $q/(2B_+) > A_+\, q^{1/\mu}$, then $q/(2B) > A\, q^{1/\mu}$. Thus the positive integer Q_A in Theorem 3 is at most $Q_A^\dagger$.

If $d_0 \le 1$, let $Q_1 = 1$. If $d_0 > 1$, then for each odd positive integer $z < d_0$ let $j_0(z)$ be the least integer $j \ge 1$ with $z \cdot 2^j \ge d_0$. Theorem 3, with constants $c\, z^{-\mu}$ and $c\, (\ell z)^{-\mu}$ and with this $j_0(z)$, gives $q < Q_\ell(z;\, \mu,\, cz^{-\mu},\, c(\ell z)^{-\mu},\, j_0(z))$ whenever z is the starting value of a cycle of length ℓ. Let Q_1 be the maximum of those finitely many values.

Now suppose $x \ge d_0$, $2^m \ge d_0$ and $q \ge \max(Q_e,\, Q_A^\dagger)$. Then $q \ge 4\ell$. For every integer $j \ge 1$ the denominators $x \cdot 2^j$ and $\ell x \cdot 2^j$ are at least d_0, so

$$\left| 2^{r/\ell} - \frac{p}{x \cdot 2^j} \right| > c\, x^{-\mu} \cdot 2^{-\mu j}$$

and

$$\left| 2^{r/\ell} - \frac{p}{\ell x \cdot 2^j} \right| > c\, (\ell x)^{-\mu} \cdot 2^{-\mu j}.$$

Theorem 3, with these constants and with $j_0 = 1$, gives $q < Q_\ell(x;\, \mu,\, cx^{-\mu},\, c(\ell x)^{-\mu},\, 1)$. In that maximum, $2^{j_0} = 2$. Since $A \ge A_-$ and $c_* = c(\ell x)^{-\mu}$,

$$Q_* \le \left(\frac{K\, (\ell x)^\mu}{c \cdot 2^{1-\mu}} \right)^{1/\gamma}, \qquad K = \frac{\ell - 1}{\ell A_-} + \ell.$$

Let

$$\Lambda = \frac{1}{\gamma} \log_2 \left(\frac{K\, \ell^\mu}{c \cdot 2^{1-\mu}} \right) + \frac{\mu}{\gamma} \log_2 \left(\frac{4}{3c} \right).$$

From $x < (4/(3c)) \cdot 2^{(\mu-1)m}$ we have $\log_2 Q_* < \Lambda + \alpha m$.

For $0 \le j < 1$ the only such integer is $j = 0$. The hypothesis at denominator x gives, for the distance d_r from $x \cdot 2^{r/\ell}$ to the nearest integer, $d_r > c\, x^{1-\mu}$. The value $B \cdot 2^{r/\ell} \cdot x/d_r$ in the maximum Q_0 is then less than $(2B_+/c)\, x^\mu$. For an odd positive integer y and for $\rho = |yx^{-1}2^{-r/\ell} - 1|$, the same hypothesis gives $B/\rho < (2B_+/c)\, x^\mu$ whenever $\rho > 0$. If ℓ is composite, the value Bx in that maximum satisfies $Bx \le B_+ x \le B_+ x^\mu$. Let $K_0 = \max(B_+,\, 2B_+/c)$. Thus $Q_0 \le K_0 x^\mu$, and

$$\log_2(K_0 x^\mu) < \log_2 K_0 + \mu \log_2 \left(\frac{4}{3c} \right) + \beta m.$$

Let m_0 be the least integer $m \geq \max(1, \lceil \log_2 d_0 \rceil)$ such that

$$\Lambda + \alpha m < m - 1$$

and

$$\log_2 K_0 + \mu \log_2 \left(\frac{4}{3c} \right) + \beta m < m - 1.$$

Such an integer exists because $\alpha < 1$ and $\beta < 1/2$. If $m \geq m_0$, both $\log_2 Q_*$ and $\log_2(K_0 x^\mu)$ are less than $m-1$, hence less than $\log_2 q$. Then $Q_* < q$ and $K_0 x^\mu < q$. Together with $q \geq Q_e$, $q \geq Q_A^\dagger$ and $q > 2$, this contradicts $q < Q_\ell(x;\, \mu,\, cx^{-\mu},\, c(\ell x)^{-\mu},\, 1)$.

Therefore $m < m_0$, or $q < 2d_0$, or $q < Q_1$, or $q < \max(Q_e, Q_A^\dagger)$. In the first of these, $q < 2^{m+1} \leq 2^{m_0}$. Let Q be the maximum of $2d_0$, of Q_1, of Q_e, of $Q_A^\dagger$ and of 2^{m_0}. Every cycle of length ℓ has $q < Q$. □

The positive root of $t^3 - t - 1 = 0$ is the plastic number, and the plastic number is greater than 1.3247. For a constant real number μ below the plastic number we have $\mu^3 - \mu - 1 < 0$, and the same argument produces a bound on q.

5 Cube roots with a constant odd part

Bauer and Bennett [2] prove the inequality for $s = 1$ and for $s = 3$. The inequality used here is the one obtained in the proof of their Theorem 2.1 from their inequality (5.3), Lemma 4.1 and Proposition 5.1, before they round μ_3 to 1.62. The companion paper records it for $s = 1$ and for $s = 3$. Lemma 4 gives the same exponent μ_3 for every positive integer s. The factor from s is carried by the constant $c_3(s')$ and by the integer $j_0(s)$.

Lemma 4 (constant odd part). Let s be a positive integer. Let $s' = s/\gcd(s,3)$, and let $v_3(s)$ be the exponent of 3 in s. Let $j_0(s)$ be the integer

$$j_0(s) = \max\Big(11000,\ \lceil 21 \log(66s')/\log(5/4) \rceil + 5,\ 14\, v_3(s) + 20\Big).$$

For $r \in \{1, 2\}$, every integer p, and every integer $j \geq j_0(s)$,

$$\left| 2^{r/3} - \frac{p}{s\, 2^j} \right| > c_3(s')\, 2^{-\mu_3 j}.$$

If $s' \leq 16$ and $v_3(s) \leq 700$, then $j_0(s) = 11000$.

Proof. Write $m = 3j + r$. For each $\delta \in \{0, 1\}$, the auxiliary positive integer of Bauer and Bennett is

$$n_1 = 1 + \left\lfloor \frac{m - 7r + 21\delta}{63} \right\rfloor.$$

This choice gives $7r/3 + 7(3n_1 - \delta) > m/3$. Since $r \geq 1$ and $\delta \leq 1$,

$$n_1 \leq 1 + \frac{m - 7r + 21\delta}{63} = \frac{m + 63 - 7r + 21\delta}{63} \leq \frac{m + 77}{63}.$$

Once $j \geq 11000$, we have $m \geq 33001$ and $n_1 \geq 497$. Also $m - 7r + 21\delta \geq 3j - 12$, so

$$n_1 > \frac{m - 7r + 21\delta}{63} \geq \frac{3j - 12}{63} > \frac{3j - 14}{63}.$$

Let A and B be the integers in their linear form for this δ, with $A \neq 0$, and set

$$e = 2n_1 - \lfloor n_1/2 \rfloor - \delta.$$

Let M be the integer $7(3n_1 - \delta) + 2r$, and let $L = M - j$. The inequality $7r/3 + 7(3n_1 - \delta) > m/3$ is $M + r/3 > j + r/3$, so $M > j$. Both M and j are integers, and therefore L is a positive integer. The second term of the linear form is $5^r B/(A\, 3^e\, 2^{M+r/3})$, and

$$\Lambda = \frac{|p\, A\, 3^e\, 2^L - s\, 5^r B|}{s\, A\, 3^e\, 2^{M+r/3}}.$$

Write D for the positive real number $A\, 3^e\, 2^{M+r/3}$, and write N for the integer $p\, A\, 3^e\, 2^L - s\, 5^r B$. Then $\Lambda = |N|/(sD)$.

Let $v = v_3(s)$. For $n_1 \geq 497$ and $\delta \in \{0, 1\}$, the exponent e increases with n_1, and

$$e \geq 2 \cdot 497 - \lfloor 497/2 \rfloor - 1 = 994 - 248 - 1 = 745.$$

The bounds $\lfloor n_1/2 \rfloor \leq n_1/2$ and $\delta \leq 1$ give $e \geq (3/2)n_1 - 1$. Combined with $n_1 > (3j - 14)/63$,

$$e > \frac{3j - 14}{42} - 1.$$

At $j = 14v + 20$ the right side equals $v + 2/21$, so $e > v$. Thus $v \leq e$ for every $j \geq 14v + 20$. In that range $N = 3^v N'$ with the integer

$$N' = p\, A\, 3^{e-v}\, 2^L - s'\, 5^r B,$$

and $\Lambda = |N'|/(s'D)$. If $N' \neq 0$, then $|\Lambda| \geq 1/(s'D)$.

Bauer and Bennett set

$$A = \frac{3^{\lfloor n_1/2 \rfloor} 2^{7n_1}}{\Pi}\, P, \qquad B = \frac{3^{2n_1 - \delta} 2^{28n_1 - 7\delta}}{\Pi}\, Q,$$

where P and Q are the values at $3/128$ of their polynomials (13) and (14) for this n_1 and δ, and Π is the positive integer gcd in their Section 5. Thus

$$\frac{B}{A} = 3^e\, 2^{7(3n_1 - \delta)}\, \frac{Q}{P}.$$

Substituting this quotient into $5^r B/(A\, 3^e\, 2^{M+r/3})$ cancels $3^e\, 2^{7(3n_1-\delta)}$ and leaves $(5/2^{7/3})^r Q/P$. Since A is a nonzero multiple of P, we have $P \neq 0$. Thus $N' = 0$ if and only if $p/(s\, 2^{m/3})$ equals that value. The two choices of δ give distinct values, by the comparison of the ratios Q/P below. For every integer p, some choice therefore has $N' \neq 0$. Choose that δ, and keep its n_1.

Let ν be the rational $r/3$. The integral for $I_{n_1,n_2}(x)$ in Section 4 of Bauer and Bennett equals $P_{n_1,n_2}(x) - (1-x)^\nu Q_{n_1,n_2}(x)$. The coefficient of x^m in that integral is a constant times the sum of the

residues at 0 and at 1 of $z^{m-n_1-1}(1-z)^{-n_2-1}$. For $m \le n_1+n_2$ the degree of the denominator exceeds the degree of the numerator by at least 2, so the residue at infinity vanishes. The sum of all residues of a rational function is zero, and the coefficient of x^m vanishes. Hence $P_{n_1,n_2}(x)-(1-x)^\nu Q_{n_1,n_2}(x)$ has a zero of order n_1+n_2+1 at $x=0$. The coefficient of x^{n_2} in Q_{n_1,n_2} is $\binom{n_1-\nu}{n_2}(-1)^{n_2}$. No factor in that product is zero, so the degree of Q_{n_1,n_2} is n_2. The same count gives degree n_1 for P_{n_1,n_2}.

Suppose the two choices of δ give the same n_1. The corresponding degrees n_2 are $4n_1$ and $4n_1-1$. Write P, Q for the polynomials of degree pair $(n_1, 4n_1)$ and P_*, Q_* for the pair $(n_1, 4n_1-1)$, and write I, I_* for the corresponding remainders. Then $PQ_* - P_*Q = IQ_* - I_*Q$. The orders of the zeros give a zero of order $5n_1+1$ for IQ_* and a zero of order $5n_1$ for I_*Q, so every coefficient of x^m in $PQ_* - P_*Q$ with $m < 5n_1$ is zero. The degree is $5n_1$, because $\deg(P_*Q) = 5n_1$ and $\deg(PQ_*) \le 5n_1-1$, and the leading coefficient is nonzero. A polynomial of degree $5n_1$ with no term of degree less than $5n_1$ equals $c\,x^{5n_1}$ with a nonzero rational c. At $x = 3/128$ the value is nonzero, so the two ratios differ.

Suppose instead that the auxiliary integers differ. The floor increases by at most 1 when its argument increases by $1/3$, so the two values are a positive integer n and $n+1$, with degree pairs $(n, 4n)$ and $(n+1, 4n+3)$. Write P, Q, I for the first pair and P_b, Q_b, I_b for the second. The orders of the zeros are $5n+1$ and $5n+5$, and

$$\deg(PQ_b) \le 5n+3, \qquad \deg(P_bQ) \le 5n+1.$$

The leading term of PQ_b has degree $5n+3$ and does not cancel, so $PQ_b - P_bQ$ has degree $5n+3$. From $IQ_b - I_bQ$, the zero at $x=0$ has order at least $5n+1$, and I_b does not meet the coefficients of x^{5n+1}, x^{5n+2}, and x^{5n+3}. Thus

$$PQ_b - P_bQ = x^{5n+1}(c_0 + c_1x + c_2x^2)$$

with rational numbers c_0, c_1, and c_2. The coefficient of x^{5n+1} in I is nonzero. The function $z^{4n}(1-z)^{-4n-1}$ is holomorphic at 0, and its expansion at infinity begins with a nonzero multiple of z^{-1}, so the residue at 1 is nonzero. The binomial prefactor $\binom{4n+\nu}{5n+1}$ is nonzero because ν is not an integer. The constant term of Q_b is the positive integer $\binom{5n+4}{n+1}$, so $c_0 \ne 0$.

The next coefficient of I, divided by the coefficient of x^{5n+1}, equals $(n+1-\nu)(4n+1)/(5n+2)$. The coefficient of x in Q_b, divided by the constant term, equals $-(n+1-\nu)(4n+3)/(5n+4)$. Their sum is

$$(n+1-\nu)\left(-\frac{4n+3}{5n+4} + \frac{4n+1}{5n+2}\right).$$

The numerator of the difference in parentheses is $-2(n+1)$, and $n+1-\nu = (3n+3-r)/3$, so

$$\frac{c_1}{c_0} = -\frac{2(n+1)(3n+3-r)}{3(5n+4)(5n+2)}.$$

For $n \ge 1$ and $r \in \{1,2\}$ we have $3n+3-r \le 3n+2$, and

$$10(5n+4)(5n+2) - 63(n+1)(3n+2) = 61n^2 - 15n - 46.$$

The roots of the right side are $(15 \pm 107)/122$, so the positive root is 1, and the quadratic is positive for every integer $n \ge 2$. The absolute value of c_1/c_0 is therefore at most $20/189$, which is less than $1/9$.

The coefficient c_2/c_0 is the sum of

$$\frac{(n+1-\nu)(n-\nu)}{2}\cdot\frac{(4n+3)(4n+2)}{(5n+4)(5n+3)},$$
$$-\frac{(n+1-\nu)^2(4n+1)(4n+3)}{(5n+2)(5n+4)},$$
$$\frac{(n+1-\nu)(n+2-\nu)(2n+1)(4n+1)}{(5n+2)(5n+3)}.$$

Clearing the denominator $(5n+2)(5n+3)(5n+4)$, together with the factors of 3 in ν, writes the sum as a quotient of two polynomials of degree 6. For both $r=1$ and $r=2$ the denominator coefficients, from degree 0 through degree 6, are

$$576,\ 6240,\ 27700,\ 64500,\ 83125,\ 56250,\ 15625.$$

For $r=1$ the numerator coefficients are

$$-\frac{64}{3},\ -\frac{1952}{9},\ -\frac{8108}{9},\ -\frac{17608}{9},\ -\frac{21085}{9},\ -\frac{4400}{3},\ -375.$$

For $r=2$ they are

$$-\frac{40}{3},\ -\frac{1346}{9},\ -\frac{6119}{9},\ -\frac{14428}{9},\ -\frac{18610}{9},\ -\frac{4150}{3},\ -375.$$

Each absolute numerator coefficient is at most $1/27$ of the matching denominator coefficient, so $|c_2/c_0|\le 1/27$. At $x=3/128$,

$$\Big|\frac{c_1}{c_0}\Big|x+\Big|\frac{c_2}{c_0}\Big|x^2\le\frac{1}{9}\cdot\frac{3}{128}+\frac{1}{27}\Big(\frac{3}{128}\Big)^2=\frac{1}{384}+\frac{1}{49152}=\frac{129}{49152}<1.$$

The factor $c_0+c_1x+c_2x^2$ is nonzero at $x=3/128$, and the two ratios differ in this case as well.

For the chosen δ, $|\Lambda|\ge 1/(s'D)$. Bauer and Bennett pass from $\Lambda\ge 1/D$ to their inequality (21). The same steps with $1/(s'D)$ in place of $1/D$ multiply the subtracted term in the numerator by s' and multiply the denominator by s'. Let F be the positive real number in their (17), so $F<11979/1000$, and let G be the positive real number $3^2\cdot 2^{28}F$. Their Proposition 8 gives $\Pi>1.8^{n_1}$. Before the factor s', the subtracted term is at most $33\cdot(1.427)^{n_1}$. Here $\pi>22/7$ and $(8/5)^3=512/125>4$, so $2^{2/3}<8/5$ and

$$\frac{4}{\pi}\,2^{14/3}<\frac{14}{11}\cdot 16\cdot\frac{8}{5}=\frac{1792}{55}<33,$$

and Bauer and Bennett record that $3^7/(2^7F)<1.427$. Since $1.427\cdot 5/4=1.78375<1.8$, the ratio of that subtracted term to 1.8^{n_1} is less than $33\cdot(4/5)^{n_1}$. Once $(5/4)^{n_1}>66s'$, the product of this ratio with s' is less than $1/2$. The numerator is then greater than $\Pi/2$, and

$$\left|1-\frac{p}{s\,2^{m/3}}\right|>\frac{1.8^{n_1}}{2s'\cdot 1.26\cdot 2^{14/3}\cdot G^{n_1}}.$$

For $s'\le 16$ and $n_1\ge 497$,

$$\Big(\frac{5}{4}\Big)^{496}=\Big(\Big(\frac{5}{4}\Big)^4\Big)^{124}>2^{124}>66\cdot 16,$$

so the comparison holds at $j = 11000$. Also $(5/4)^4 > 2$, so

$$\frac{\log(66 \cdot 16)}{\log(5/4)} < \frac{\log(2^{11})}{\log(5/4)} = 44 \cdot \frac{\log 2}{\log((5/4)^4)} < 44.$$

Thus $21 \log(66s')/\log(5/4) + 5 < 21 \cdot 44 + 5 = 929 < 11000$ for every $s' \le 16$, and $14 \cdot 700 + 20 = 9820 < 11000$. The formula for $j_0(s)$ therefore equals 11000 when $s' \le 16$ and $v_3(s) \le 700$. For a larger s' the comparison holds once $n_1 \ge \log(66s')/\log(5/4)$. Combined with $n_1 > (3j - 14)/63$, this is true for every $j \ge \lceil 21 \log(66s')/\log(5/4) \rceil + 5$. Thus $v \le e$ and $(5/4)^{n_1} > 66s'$ both hold for every $j \ge j_0(s)$.

Multiplying by $2^{r/3}$ gives the distance from $2^{r/3}$ to $p/(s\,2^j)$. Since $G < 9 \cdot 2^{28} \cdot 11979/1000$ and $360000 \cdot N > 201 \cdot 9 \cdot 2^{28} \cdot 11979$,

$$\frac{1.8\,N}{G} > \frac{201}{200}.$$

For $n_1 \ge 200$,

$$\Big(\frac{201}{200}\Big)^{n_1} > \Big(\frac{201}{200}\Big)^{200} > 2,$$

because $(1 + 1/200)^{200} > 1 + 200 \cdot (1/200)$. Divide the lower bound on the distance by $c_3(s')\,2^{-\mu_3 j}$. Using $m = 3j + r$ and $n_1 \le (m + 77)/63$, the quotient is at least

$$\frac{2^{r/3+4}}{1.26 \cdot 2^{14/3}} \Big(\frac{1.8\,N}{G}\Big)^{n_1} N^{(2-r)/63}.$$

For $r = 2$ the powers of 2 cancel and the quotient is greater than $(1.8\,N/G)^{n_1}/1.26$, which is greater than $2/1.26 > 1$. For $r = 1$ the same expression is larger by $N^{1/63}/2^{1/3}$. That factor exceeds 1 because $N > 2^{21}$. The resulting constant is $c_3(s')$. □

Per unit of j the restricted irrationality exponent is $\mu_3 \approx 1.61487$, which lies below φ.

6 The integer $Q_3(x)$

Let x be an odd positive integer. Let s' be the positive integer $x/\gcd(x, 3)$. These formulas are the length-3 case of Theorem 3, with the factor $1 + 1/x$ in place of $B = 2 + 1/x$, the constant real number $\mu = \mu_3$, the constant positive real number $c = c_3(s')$, the constant positive real number $c_* = c_3(x)$, and the positive integer $j_0 = J_3(x)$.

Set

$$A_3 = \frac{x}{2}\,2^{1/3} \left(\frac{c_3(s')}{2^{2/3}\,(1 + 1/x)} \right)^{1/\mu_3}$$

and

$$Q_{3,*}(x) = \left(\frac{2/(3A_3) + 3}{c_3(x)\,2^{1-\mu_3}} \right)^{1/\gamma_3}.$$

Let $J_3(x) = \max(j_0(x),\, j_0(3x))$. For each $r \in \{1, 2\}$ and each integer j with $0 \le j < J_3(x)$, and for each odd positive integer y with $y < 3x \cdot 2^{J_3(x)}$, set

$$\rho = \left| \frac{y}{x\,2^j} \cdot 2^{-r/3} - 1 \right|.$$

Whenever $\rho > 0$, set the corresponding value equal to $(1 + 1/x)/\rho$. The integer $Q_{3,0}(x)$ is the maximum of those finitely many values. Then

$$Q_3(x) = \max(Q_{3,0}(x),\, Q_{3,*}(x),\, 2^{J_3(x)+1}).$$

The factor $x/2$ in A_3 is the condition $|\rho| < 1/2$. The power $2^{1/3}$ is the smaller cube root in the lower bound for another element of the cycle, and $2^{2/3}$ is the larger cube root in the error of that approximation. The numerator $2/(3A_3) + 3$ is two other elements, divided by the length, plus the length.

Corollary 5. Let x be an odd positive integer. If x is the starting value of a cycle of length 3, then $q < Q_3(x)$.

Proof. The statement of Theorem 2 of [6] does not apply: the cycle may have any starting value, and the odd part is x or $3x$ rather than 1 or ℓ. The comparison in its proof does apply, with that odd part.

Consider a cycle of length 3 with starting value x, written $x \to y \to z \to x$. Lemma 1 gives $2^h = (q + 1/x)(q + 1/y)(q + 1/z)$, so $q = \lfloor 2^{h/3} \rfloor$ and $3 \nmid h$. Write $2^{h/3} = q + \delta$ with $0 < \delta < 1$, and write $h = 3m + r$ with $r \in \{1, 2\}$. The fraction $(3xq + 1)/(3x \cdot 2^m)$ then satisfies

$$\left| 2^{r/3} - \frac{3xq + 1}{3x \cdot 2^m} \right| = \frac{|\delta - 1/(3x)|}{2^m}.$$

The denominator $3x \cdot 2^m$ has odd part $3x$, so Lemma 4 applies with constant $c_3(3x/\gcd(3x, 3)) = c_3(x)$.

Lemma 1 gives $e^{-3/q} \le 3\delta/\sigma \le e^{3/q}$ with $\sigma = 1/x + 1/y + 1/z$. Therefore $|\delta - \sigma/3| \le (\sigma/q)\, e^{3/q}$, and

$$\big|\delta - 1/(3x)\big| \le (1/y + 1/z)/3 + (\sigma/q)\, e^{3/q}.$$

From $qx + 1 = 2^{k_1} y$ and $2^{h/3} = q + \delta$,

$$\frac{y}{x} = 2^{h/3 - k_1} \cdot \frac{q + 1/x}{q + \delta}.$$

Since $3 \nmid h$, the exponent $h - 3k_1$ is congruent to 1 or 2 modulo 3. Write it as $3j + r'$ with $r' \in \{1, 2\}$ and

$$\frac{y}{x\, 2^j} = 2^{r'/3}(1 + \rho), \qquad \rho = \frac{1/x - \delta}{q + \delta}.$$

Then $|\rho| < (1 + 1/x)/q$. If $q < 30$, then $q < Q_3(x)$, because Appendix C records $Q_{3,*}(x) > 2^{17906}$. Assume $q \ge 30$. Let $\sigma = 1/x + 1/y + 1/z$. From Lemma 1, $e^{-3/q} \le 3\delta/\sigma \le e^{3/q}$. Hence $|3\delta/\sigma - 1| \le e^{3/q} - 1 \le (3/q)\, e^{3/q}$, and $|\delta - \sigma/3| \le (\sigma/q)\, e^{3/q}$. From $qx + 1 = 2^{k_1} y$ and $qy + 1 = 2^{k_2} z$ there is a real number ρ_z with

$$\rho_z = \frac{q(1/x + 1/y - 2\delta) + 1/(xy) - \delta^2}{(q + \delta)^2}.$$

Since $1/x$, $1/y$ and $1/z$ lie in $(0,1]$, we have $|1/x + 1/y - 2/z| \le 2$, so

$$\left|1/x + 1/y - 2\delta\right| \le \frac{2}{3} + \frac{6\,e^{3/q}}{q}.$$

Hence $|\rho_z| \le (2/3)/q + (6e^{3/q}+2)/q^2$. For $q \ge 30$ we have $e^{3/q} < 1.1052$ and $6e^{3/q}+2 < 8.64$, while $8.64/q \le 8.64/30 < 1/3$. Thus $|\rho_z| < (1+1/x)/q$. For $q \ge 30$ both $|\rho|$ and $|\rho_z|$ are less than $1/2$. Write $2h - 3(k_1+k_2) = 3j_z + r_z$ with an integer j_z and with $r_z \in \{1,2\}$, so

$$\frac{z}{x\,2^{j_z}} = 2^{r_z/3}(1+\rho_z).$$

If $j < 0$, let $t = -j$ and let d be the distance from $x\,2^{r'/3}$ to the nearest integer. Then $y\,2^t/x = 2^{r'/3}(1+\rho)$, so $|\rho| \ge 2^{-r'/3}\,d/x$ and

$$q < (1+1/x)\,2^{r'/3}\,x/d.$$

Let p be the nearest integer to $x\,2^{r'/3}$. Since $2^{r'/3}$ is irrational, $|x^3\,2^{r'} - p^3| \ge 1$, and

$$d = \frac{|x^3\,2^{r'} - p^3|}{x^2\,2^{2r'/3} + x\,2^{r'/3}\,p + p^2} \ge \frac{1}{3(x\,2^{r'/3}+1)^2}.$$

Since $r' \le 2$ and $1 + 1/x \le 2$,

$$q < 12x\,(2x+1)^2.$$

The same bound holds when $j_z < 0$, with r_z in place of r'. Since $\mu_3 > 3/2$, we have $\gamma_3 < 1/6$ and $2^{1-\mu_3} < 2^{-1/2}$. Since $N > 2^{33}$ and $79/63 > 1$, we have $N^{79/63} > 2^{33\cdot 79/63}$. Here $33 \cdot 79 = 2607$ and $2607/63 > 41$, so $N^{79/63} > 2^{41}$ and $c_3(x) < 2^{-46}/x$. The product $c_3(x)\,2^{1-\mu_3}$ is therefore less than $2^{-46}/x$. Also $2/(3A_3) + 3 > 3$ and $1/\gamma_3 > 6$, so

$$Q_{3,*}(x) > (3x \cdot 2^{46})^6.$$

For every integer $x \ge 1$ the right side is greater than $12x(2x+1)^2$, because $(2x+1) \le 3x$ and $3^6 \cdot 2^{276} > 108$. Thus a negative exponent forces $q < Q_{3,*}(x)$.

Now suppose $j \ge 0$ and $j_z \ge 0$. If $0 \le j < J_3(x)$, then $|\rho| < (1+1/x)/q$, so $q < (1+1/x)/|\rho|$. The value $(1+1/x)/|\rho|$ is one of the numbers whose maximum is $Q_{3,0}(x)$, and $q < Q_{3,0}(x)$. The same bound holds when $0 \le j_z < J_3(x)$. If $j \ge J_3(x)$ and $j_z \ge J_3(x)$, Lemma 4 and $|2^{r'/3} - y/(x\,2^j)| = 2^{r'/3}|\rho| \le 2^{2/3}|\rho|$ give

$$2^j > \left(\frac{c_3(s')\,q}{2^{2/3}\,(1+1/x)}\right)^{1/\mu_3},$$

and therefore $y > A_3\,q^{1/\mu_3}$. The same lower bound holds for z. Then $\sigma \le 1/x + 2A_3^{-1}q^{-1/\mu_3}$, so

$$(\sigma/q)\,e^{3/q} \le \frac{e^{3/q}}{xq} + 2A_3^{-1}e^{3/q}\,q^{-1-1/\mu_3}.$$

Here $N < 2^{34}$ and $79/63 < 4/3$, so $N^{79/63} < 2^{46}$ and $c_3(s') > 2^{-51}/x$. Also $N > 2^{33}$, so $\mu_3 > 3/2$, hence $1/\mu_3 < 2/3$, and $2^{2/3}(1+1/x) \le 4$. Hence $A_3 > 2^{-37}\,x^{1/3}$ and $4A_3^{-1} < 2^{39}$. Appendix C records $Q_{3,*}(x) > 2^{17906}$, so $q > 4A_3^{-1}$ whenever $q > Q_{3,*}(x)$. For such q we have $e^{3/q} < 2$ and

$q^{-1} \le q^{-1/\mu_3}$, because $\mu_3 > 1$. The first summand is then at most $2\,q^{-1/\mu_3}$, and the second is at most q^{-1/μ_3}. Therefore

$$\bigl|\delta - 1/(3x)\bigr| \le (2/(3A_3) + 3)\, q^{-1/\mu_3}.$$

If $m \ge J_3(x)$, the denominator $3x \cdot 2^m$ has odd part $3x$, so Lemma 4 applies. Together with $2^m < 2q$ and $1 - \mu_3 < 0$, it gives

$$\bigl|\delta - 1/(3x)\bigr| > c_3(x)\, 2^{1-\mu_3}\, q^{1-\mu_3}.$$

These two bounds on $|\delta - 1/(3x)|$ conflict for every integer $q \ge Q_{3,*}(x)$. The lower bound is strict, so the conflict includes $q = Q_{3,*}(x)$. If instead $m < J_3(x)$, then $2^m > q/2$ gives $q < 2^{m+1} < 2^{J_3(x)+1}$. In every case $q < Q_3(x)$. □

7 The integer $Q_2(x)$

These formulas are the length-2 case of Theorem 3, with the factor $1 + 1/x$ in place of $B = 1$, the constant real number $\mu = 1.465$, the constant positive real numbers $c = c_* = 1$, and the positive integer $j_0 = j_0(x)$.

Let x be an odd positive integer. Let $\mu_2 = 1.465$ and $\gamma_2 = 1 - \mu_2 + 1/\mu_2$. The value $\mu_2 = 1.465$ is the exponent printed to three decimals in Corollary 1.4 of Bauer and Bennett [1], for the square root of 2. For each constant positive integer s that corollary gives an integer $j_0(s)$ such that for every integer p and every integer $j \ge j_0(s)$,

$$\left|\sqrt{2} - \frac{p}{s\,2^j}\right| > 2^{-\mu_2 j}.$$

The constant in front is 1. The integer $j_0(x)$ is that exponent for $s = x$. Set

$$A_2 = \frac{x}{2}\, 2^{1/2} \left(\frac{1}{2^{1/2}\,(1 + 1/x)} \right)^{1/\mu_2}$$

and

$$Q_{2,*}(x) = \left(\frac{1/(2A_2) + 2}{2^{1-\mu_2}} \right)^{1/\gamma_2}.$$

For each integer j with $0 \le j < j_0(x)$, and for each odd positive integer y with $y < 3x \cdot 2^{j_0(x)}$, set

$$\rho = \left| \frac{y}{x\,2^j} \cdot 2^{-1/2} - 1 \right|.$$

Whenever $\rho > 0$, set the corresponding value equal to $(1 + 1/x)/\rho$. The integer $Q_{2,0}(x)$ is the maximum of those finitely many values. The length-2 bound is this integer:

$$Q_2(x) = Q_{2,0}(x).$$

The quantities $2^{j_0(x)+1}$ and $Q_{2,*}(x)$ are at most $Q_{2,0}(x)$ for every odd positive integer x. The proof below records that comparison.

Corollary 6. Let x be an odd positive integer. If x is the starting value of a cycle of length 2, then $q < Q_2(x)$.

Proof. Consider a cycle of length 2 with starting value x, written $x \to y \to x$. Here h is odd. Write $2^{h/2} = q + \delta$ and $h = 2m + 1$. The argument for $Q_3(x)$ in Section 6, with one other element and with Corollary 1.4 of Bauer and Bennett (2002) [1] in place of Lemma 4, produces A_2 and $Q_{2,*}(x)$. Both the product around the cycle and the approximation for y have odd part x, and the constant in front of $2^{-\mu_2 j}$ is 1. The numerator $1/(2A_2) + 2$ is that one other element, divided by the length, plus the length.

The exponent $j = 0$ fails the inequality of Corollary 1.4 for every $x \ge 1$, so $j_0(x) \ge 1$. The distance from $x\,2^{j_0(x)-1}\sqrt{2}$ to an odd integer is at most 1, and the resulting value of $(1 + 1/x)/\rho$ is at least $(x + 1)\sqrt{2}\,2^{j_0(x)-1}$. For $x \ge 3$ this is at least $2^{j_0(x)+1}$, because $(x + 1)\sqrt{2} \ge 4\sqrt{2} > 4$. For $x = 1$ the same lower bound is $2\sqrt{2}\,2^{j_0(1)-1} > 2^{j_0(1)}$. The exponent $j = 7$ fails the inequality for $x = 1$: the value of $(1 + 1)/\rho$ there has logarithm $14.192568 > 10$, so $j_0(1) > 7$ and that value lies in the maximum $Q_{2,0}(1)$. Thus $2^{j_0(x)} \le Q_{2,0}(x)$.

The quantity A_2 increases with x, so $Q_{2,*}(x)$ decreases, and $Q_{2,*}(1) < 2^{11}$. Every exponent j with $x \cdot 2^{-0.465j} \ge 1/2$ fails the inequality, and the largest such j contributes at least $(1 + 1/x)\,x\sqrt{2}\,2^j$ even when the distance to an odd integer is 1. For every odd $x \ge 5$ that lower bound already exceeds $Q_{2,*}(x)$. For $x = 1$ and $x = 3$ the values $Q_{2,0}(1) > 2^{14}$ and $Q_{2,0}(3) > 2^{17}$ do the same. Thus $Q_{2,*}(x) \le Q_{2,0}(x)$, and $Q_2(x) = Q_{2,0}(x)$. For $\ell = 2$, Theorem 3 has $B = 1$ and $Q_A = 1$. The factor $1 + 1/x$ is at least B, so each value B/ρ is at most the corresponding value $(1 + 1/x)/\rho$. At $q = 8$ we have $q \ge 4 \cdot 2$ and $e^{2/8} < 8^{1-1/1.465}$, so $Q_e \le 8$. The exponent $j = 2$ satisfies $x \cdot 2^{-0.465 \cdot 2} > 1/2$ for every $x \ge 1$, so it fails the inequality, and the value $(1 + 1/x)/\rho$ is at least $4(x + 1)\sqrt{2} \ge 8\sqrt{2} > 8$. Thus $2^{j_0(x)}$ and Q_e are at most $Q_{2,0}(x)$. The remaining values in Q_0 are $\sqrt{2}\,x/d$, where d is the distance from $x\sqrt{2}$ to the nearest integer. If $d \ge 1$, that value is at most $\sqrt{2}\,x$, which is less than $8\sqrt{2} \le Q_{2,0}(x)$ for $x = 1$ and less than $4(x + 1)\sqrt{2}$ for $x \ge 3$. If $d < 1$ and p is the nearest integer, then $|p^2 - 2x^2| \ge 1$ and $p + x\sqrt{2} < 2x\sqrt{2} + 1 < 4x$, so $d > 1/(4x)$ and the value is less than $4\sqrt{2}\,x^2$. For $x = 1$ we have $4\sqrt{2} < 2^{14} < Q_{2,0}(1)$. For $x \ge 3$, the largest integer j with $x \cdot 2^{-0.465j} \ge 1/2$ satisfies $2^j \ge (2x)^{1/0.465}/2$, and

$$(x + 1)\sqrt{2} \cdot 2^j \ge \frac{\sqrt{2}\,x\,(2x)^{1/0.465}}{2}.$$

The quotient of the right side by $4\sqrt{2}\,x^2$ is $2^{1/0.465}\,x^{1/0.465-1}/8 > 2^{2.15}\,x/8$. Since $2^{2.15} > 4.4$, the quotient is greater than $0.55x \ge 0.55 \cdot 3 > 1$. Thus the maximum in Theorem 3 is at most $Q_2(x)$, and a cycle of length 2 with starting value x satisfies $q < Q_2(x)$.

For $x = 1$ and length 2, Proposition A.1 of [6] gives the only such q, namely $q = 5$. That conclusion does not change the formula for $Q_2(1)$. □

8 Cycles of length 2

The aim of the research in this section was to prove that the only cycles of length 2 are $1 \to 3$, $35 \to 99$, and $27 \to 611$. That is more ambitious than Theorem 1 of the companion paper, which only rules out a cycle of length 3 with starting value 1. Here the starting value may be any odd

positive integer. A cycle of length 2 was selected because it's the smallest non-trivial length, making it look like the easiest problem of this kind.

For a constant starting value, Theorem 3 bounds q, and a computation can finish the search. The bound grows with the starting value, so the same search does not cover every starting value at once. Proposition 3 shows that its lower bound, with exponent 1.3247, gives a bound on q for every starting value and every length at least 2. Bauer and Bennett (2002) [1] give the exponent 1.465 for $\sqrt{2}$.

The aim was not achieved, but this section records the properties of a cycle of length 2, and large bounds that follow from those properties by a modest computation.

Throughout this section, $x \to y \to x$ is a cycle of length 2, and x is its starting value. Here $qx + 1 = 2^{k_1} y$ and $qy + 1 = 2^{k_2} x$, and $h = k_1 + k_2$. Lemma 1 gives $q = \lfloor 2^{h/2} \rfloor$ and h odd. Let $D = 2^h - q^2$.

Lemma 7 (the length-2 equation). Then $k_1 < k_2$. Let $t = k_2 - k_1$. Then t is positive and odd, $h = 2k_1 + t$, and

$$x = \frac{q + 2^{k_1}}{D}, \qquad y = \frac{q + 2^{k_2}}{D}.$$

The integer D divides $2^t - 1$, and $D \equiv 7 \pmod 8$. Every prime divisor of D is congruent to 1 or 7 modulo 8. Let $u = (2^t - 1)/D$. Then u is a positive odd integer,

$$y = x + u\, 2^{k_1}, \qquad y^2 - 2^t x^2 = u,$$

$2^{k_1} \le q - 1$, $t > \log_2(q/x)$, and $y^2 > qx + 1$.

Proof. Since h is odd, $k_1 \ne k_2$. From $qy + 1 = 2^{k_2} x$ and $y = (qx + 1)/2^{k_1}$,

$$2^h x = q(qx + 1) + 2^{k_1},$$

so $x(2^h - q^2) = q + 2^{k_1}$. Lemma 1 gives $q^2 < 2^h$, so $D > 0$ and $x = (q + 2^{k_1})/D$. The same elimination with the roles of x and y exchanged gives $y = (q + 2^{k_2})/D$. Since $x < y$ and $D > 0$, we have $k_1 < k_2$. Thus $t > 0$. From $h = 2k_1 + t$ and h odd, t is odd.

Now D divides $q + 2^{k_1}$ and $q + 2^{k_2}$, so D divides

$$(q + 2^{k_2}) - (q + 2^{k_1}) = 2^{k_1}(2^t - 1).$$

Here q is odd and $h \ge 3$, so 2^h is divisible by 8 and $q^2 \equiv 1 \pmod 8$. Thus $D \equiv 7 \pmod 8$, and in particular D is odd. Hence D divides $2^t - 1$. Let p be a prime divisor of D. Then $q^2 = 2^h - D \equiv 2^h \pmod p$. If p divides q, then p divides 2^h. But D is odd, so $p \ne 2$, and p does not divide 2^h. Thus 2^h is a nonzero square modulo p. Since h is odd, 2 is a quadratic residue modulo p, and therefore $p \equiv 1 \pmod 8$ or $p \equiv 7 \pmod 8$.

The quotient $u = (2^t - 1)/D$ is a positive integer. Both $2^t - 1$ and D are odd, so u is odd. Then

$$y - x = \frac{2^{k_2} - 2^{k_1}}{D} = u\, 2^{k_1},$$

so $y = x + u\,2^{k_1}$. Substitute $q = Dx - 2^{k_1}$ into $qx + 1 = 2^{k_1} y$:

$$Dx^2 - 2^{k_1+1}x + 1 = u\,4^{k_1}.$$

Therefore

$$\begin{aligned} y^2 - 2^t x^2 &= \left(x + u\,2^{k_1}\right)^2 - 2^t x^2 \\ &= u\,2^{k_1+1}x + u^2 4^{k_1} - x^2(2^t - 1) \\ &= u(2^{k_1+1}x + u\,4^{k_1} - Dx^2) \\ &= u. \end{aligned}$$

Since x and y are distinct and odd, $y \geq x + 2$. Then

$$2^{k_1} = \frac{qx+1}{y} \leq \frac{qx+1}{x+2}.$$

The inequality $(qx + 1)/(x + 2) < q$ is $1 < 2q$. Thus $2^{k_1} < q$. Both are integers, so $2^{k_1} \leq q - 1$.

From $y - x \geq 2^{k_1}$ and $2^{k_1} = (qx + 1)/y$,

$$2^{k_1}(x + 2^{k_1}) \leq qx + 1,$$

so $2^{2k_1} \leq x(q - 2^{k_1}) + 1$. For $k_1 \geq 1$ the right side is at most $x(q - 2) + 1$, and $x(q - 2) + 1 < xq$ because $1 < 2x$. Thus $2^{2k_1} < xq$. Lemma 1 gives $2^{h/2} = q + \delta$ with $\delta > 0$, so $h > 2\log_2 q$. Therefore

$$t = h - 2k_1 > 2\log_2 q - \log_2(xq) = \log_2(q/x).$$

Finally $y(y - x) \geq y \cdot 2^{k_1} = qx + 1$. Since $x \geq 1$, we have $y > y - x$, and therefore $y^2 > qx + 1$. □

Lemma 8 (the case $D = 7$). If $D = 7$, the cycle is $1 \to 3$ with $q = 5$, or $35 \to 99$ with $q = 181$, or $27 \to 611$ with $q = 181$.

Proof. Here $q^2 + 7 = 2^h$. By Theorem 1.8 of Bauer and Bennett (2002), the positive solutions (q, h) are $(1, 3)$, $(3, 4)$, $(5, 5)$, $(11, 7)$ and $(181, 15)$. Lemma 7 requires $q \geq 3$ and h odd, which leaves $(5, 5)$, $(11, 7)$ and $(181, 15)$. The order of 2 modulo 7 is 3, so 3 divides t, and Lemma 7 requires t odd and $1 \leq t \leq h - 2$.

For $(q, h) = (5, 5)$ the only such t is 3, so $k_1 = 1$ and $k_2 = 4$. Then $(5+2^1)/7 = 1$ and $(5+2^4)/7 = 3$, which is $1 \to 3$.

For $(q, h) = (11, 7)$ we have $-11 \equiv 3 \pmod 7$. The powers of 2 modulo 7 are 2, 4 and 1, so $q + 2^{k_1}$ is never divisible by 7.

For $(q, h) = (181, 15)$ the admissible values are $t = 3$ and $t = 9$. If $t = 3$, then $k_1 = 6$ and $k_2 = 9$, and

$$\frac{181 + 2^6}{7} = 35, \qquad \frac{181 + 2^9}{7} = 99,$$

which is $35 \to 99$. If $t = 9$, then $k_1 = 3$ and $k_2 = 12$, and

$$\frac{181 + 2^3}{7} = 27, \qquad \frac{181 + 2^{12}}{7} = 611,$$

which is $27 \to 611$. □

Lemma 9 (the case $u = 1$). If $u = 1$, the cycle is $1 \to 3$ with $q = 5$, or $35 \to 99$ with $q = 181$.

Proof. Here $D = 2^t - 1$, so $q^2 = 2^{2k_1+t} - 2^t + 1$ and

$$(q-1)(q+1) = 2^t(2^{2k_1} - 1).$$

The factor $2^{2k_1} - 1$ is odd, and $\gcd(q-1, q+1) = 2$, so

$$v_2(q-1) + v_2(q+1) = t$$

with the smaller valuation equal to 1. Lemma 7 gives t odd and $D \equiv 7 \pmod 8$, so $t \geq 3$. Let $c = t - 2$, which is odd and at least 1.

If $v_2(q+1) = 1$ and $v_2(q-1) = t-1$, write $q + 1 = 2r$ and $q - 1 = 2^{t-1}s$ with r and s odd. Then $r = 2^{t-2}s + 1$ and $s(2^{t-2}s + 1) = 2^{2k_1} - 1$, so

$$2^c s^2 + s + 1 = 2^{2k_1}.$$

If $v_2(q-1) = 1$ and $v_2(q+1) = t-1$, write $q - 1 = 2s$ and $q + 1 = 2^{t-1}r$ with r and s odd. Then $s = 2^{t-2}r - 1$ and $r(2^{t-2}r - 1) = 2^{2k_1} - 1$, so

$$2^c r^2 - r + 1 = 2^{2k_1}.$$

In either case, for a positive odd integer X,

$$\left(2^{c+1}X \pm 1\right)^2 + (2^{c+2} - 1) = 2^{2k_1+c+2},$$

the plus sign for $X = s$ and the minus sign for $X = r$.

Suppose $c \geq 3$. Then $2^{c+2} - 1 = 2^k - 1$ with $k = c + 2 \geq 5$. Theorem 1.8 of Bauer and Bennett (2002) gives exactly two positive solutions of $Y^2 + (2^k - 1) = 2^n$, namely $Y = 1$ with $n = k$, and $Y = 2^{k-1} - 1$ with $n = 2k - 2$. The first gives $2k_1 + c + 2 = c + 2$, so $k_1 = 0$. The second gives $2k_1 + c + 2 = 2c + 2$, so $2k_1 = c$. But c is odd, so neither solution has an integer $k_1 \geq 1$.

Now $c = 1$, so $t = 3$ and $D = 7$, and the equation is $Y^2 + 7 = 2^{2k_1+3}$ with $Y = 4X \pm 1$. The same theorem gives $Y \in \{1, 3, 5, 11, 181\}$ and $2k_1 + 3 \in \{3, 4, 5, 7, 15\}$. The exponent $2k_1 + 3$ is odd and at least 5, which leaves 5, 7 and 15. If the exponent is 5, then $Y = 5$ and $k_1 = 1$. The equation $4X - 1 = 5$ has no integer solution, while $4X + 1 = 5$ gives $X = 1$. This is the plus sign, $q - 1 = 2^2 \cdot 1$, and $q = 5$, the cycle $1 \to 3$. If the exponent is 7, then $Y = 11$ and $k_1 = 2$. The equation $4X + 1 = 11$ has no integer solution, while $4X - 1 = 11$ gives $X = 3$. This is the minus sign and $q = 11$. Then $(11 + 2^2)/7 = 15/7$, which is not an integer. If the exponent is 15, then $Y = 181$ and $k_1 = 6$. The equation $4X - 1 = 181$ has no integer solution, while $4X + 1 = 181$ gives $X = 45$. This is the plus sign, $q - 1 = 2^2 \cdot 45$, and $q = 181$, the cycle $35 \to 99$. □

Proposition 10. Any cycle of length 2 other than $1 \to 3$, $35 \to 99$ and $27 \to 611$ has

$$q \geq 2^{160{,}000}, \qquad D \geq 15, \qquad u \geq 3,$$

every odd element at least 37, and, writing $h = 2m + 1$,

$$m \geq 160{,}000, \qquad D > 2^{0.52m+1}, \qquad t \geq 83{,}203.$$

In particular $k_1 \leq \lfloor \log_2 q \rfloor - 41{,}601$ and $k_2 \geq \lfloor \log_2 q \rfloor + 41{,}602$. Also $t > \log_2(3q/x)$.

Proof. Appendix C searches every cycle of length 2 with $h \leq 320{,}000$ and finds only the three cycles named above. That search excludes every cycle of length 2 with $q < 2^{160{,}000}$. Thus any other cycle has $q \geq 2^{160{,}000}$. Since $2^{h/2} > q$, we have $h > 2 \cdot 160{,}000$, so $h \geq 320{,}001$. With $h = 2m + 1$ this is $m \geq 160{,}000$.

Lemma 8 gives $D \neq 7$. Lemma 7 gives $D \equiv 7 \pmod 8$ and $D > 0$, so $D \geq 15$. Lemma 9 gives $u \neq 1$. Lemma 7 gives that u is a positive odd integer, so $u \geq 3$.

Let z be an odd element of the cycle with $z \leq 35$. Theorem 3 gives $q < Q_2(z)$. Appendix C records $Q_2(z) \leq 2^{33}$ for every odd $z \leq 35$. Then $q < 2^{33} < 2^{160{,}000}$, so the cycle is one of the three found by the search. Each of those has its odd elements in $\{1, 3, 27, 35, 99, 611\}$. An element $z \leq 35$ on one of them is 1, 3, 27 or 35. Therefore every odd element of any other cycle is at least 37.

Corollary 1.6 of Bauer and Bennett (2002) gives

$$\left| \sqrt{2} - \frac{q}{2^m} \right| > 2^{-1.48m}$$

for every integer $m > 2$ other than 3, 7 and 8. Here $m \geq 160{,}000$. Writing $2^{h/2} = q + \delta$ as in Lemma 1,

$$\delta > 2^{-0.48m}.$$

Then $D = \delta(2q + \delta) > 2q\delta$, so

$$\log_2 D > 1 + \log_2 q - 0.48m.$$

Since $q > 2^m\sqrt{2} - 1$ and $m \geq 160{,}000$,

$$\log_2 q > m + \tfrac{1}{2} - 2^{-159{,}999}.$$

Hence $\log_2 D > 0.52m + 1.5 - 2^{-159{,}999} > 0.52m + 1$, and $D > 2^{0.52m+1}$. At $m = 160{,}000$ this is $D > 2^{83{,}201}$. For $m > 160{,}000$ the same lower bound is larger. Lemma 7 gives $D \leq 2^t - 1$, so $t > 83{,}201$. Since t is odd, $t \geq 83{,}203$.

Since t is odd, $(t-1)/2$ is an integer, and $k_1 = m - (t-1)/2$. Thus $k_1 \leq m - 41{,}601$. Since $m \geq 160{,}000$ and $\sqrt{2} - 1 > 2^{-m}$, we have $2^m\sqrt{2} > 2^m + 1$, so $q \geq 2^m + 1$. Also $\sqrt{2} < 2$, so $q < 2^m\sqrt{2} < 2^{m+1}$. Thus $\lfloor \log_2 q \rfloor = m$ and $k_1 \leq \lfloor \log_2 q \rfloor - 41{,}601$. Also $k_2 = m + (t+1)/2 \geq m + 41{,}602 = \lfloor \log_2 q \rfloor + 41{,}602$.

For the last claim, $u \geq 3$ gives $y - x \geq 3 \cdot 2^{k_1}$. Combined with $2^{k_1} = (qx + 1)/y$,

$$3 \cdot 4^{k_1} \leq x(q - 2^{k_1}) + 1 < xq,$$

so $2^{2k_1} < xq/3$. With $h > 2\log_2 q$,

$$t = h - 2k_1 > 2\log_2 q - \log_2(xq/3) = \log_2(3q/x).$$

□

Lemma 11. Any cycle of length 2 other than $1 \to 3$, $35 \to 99$ and $27 \to 611$ has every odd element greater than 2^{2559}.

Proof. Suppose the cycle is other than those three, and suppose $x \le 2^{2559}$.

Theorem 1.5 of Bauer and Bennett (2002) [1] applies to the solution $181^2 + 7 = 2^{15}$, with base 2, with $a = 1$, with the power of 2 equal to 15, with constant term 7 and with $\alpha = 3$, taking the odd part to be x. In that theorem log is the natural logarithm. Let $\xi = 7/2^{15}$. Formula (1.2) of that paper gives $r(3,\xi) = 0.250030044752$, and formula (1.3) gives $F(r(3,\xi),3,\xi) = 9.479962374117$. Table (1.17) gives $c_1(3) = 2.458$ and $\log d_1(3) = -17.335$. The remaining parameters in that theorem are $\Delta_0 = 1$, $\kappa(7) = 4$, $\Omega = 2^{17}$, $m_1 = 17$, $\Omega_1 = 2^{17}$ and $\Delta_1 = 1$. Formulas (1.13) and (1.16) do not depend on x. They give

$$\chi_1 = 4.968969066213, \qquad \chi_3 = 8.684694 \times 10^{15}$$

and

$$\lambda_1 = \frac{\log \chi_3}{2 \cdot 17 \cdot \log 2} = 1.5572763652.$$

Let $\lambda_2 = 1.617$ be a constant real number. Then $\lambda_2 < \varphi$.

Let $c_2(x)$ be the constant in Theorem 1.5 at odd part x, and let $m_2(x)$ be the minimum in (1.15) at the prime 2. Define the real numbers

$$k(x) = \frac{\log(1/c_2(x))}{(\lambda_2 - \lambda_1)\log 2}, \qquad k_0(x) = \max(m_2(x) - \tfrac{1}{2},\, k(x)).$$

Theorem 1.5 gives, for every integer p and every integer $j \ge m_2(x) - 1/2$,

$$\left|\sqrt{2} - \frac{p}{x\,2^j}\right| > c_2(x)\,2^{-\lambda_1 j}.$$

For every integer $j \ge k(x)$ we have $c_2(x)\,2^{-\lambda_1 j} \ge 2^{-\lambda_2 j}$. Therefore, for every integer p and every integer $j \ge k_0(x)$,

$$\left|\sqrt{2} - \frac{p}{x\,2^j}\right| > 2^{-\lambda_2 j}.$$

Formula (1.14), at $\alpha = 3$ and at odd part x, increases with x, and $c_2(x)$ decreases as x increases. Both $m_2(x)$ and $k(x)$ increase with x, and so does $k_0(x)$. At $x = 1$ we have $k_0(1) = 2840.688\ldots$, so $|\sqrt{2} - p/2^j| > 2^{-\lambda_2 j}$ for every integer $j \ge 2841$. At $x = 2^{2559}$ we have $k_0(2^{2559}) = 77426.543\ldots$. Since $x \le 2^{2559}$, we have $k_0(x) \le k_0(2^{2559})$.

Let $\gamma = 1 - \lambda_2 + 1/\lambda_2$ and

$$A(x) = \frac{x}{2}\,2^{1/2}\left(\frac{1}{2^{1/2}\,(1 + 1/x)}\right)^{1/\lambda_2}, \qquad Q_*(x) = \left(\frac{1/(2A(x)) + 2}{2^{1-\lambda_2}}\right)^{1/\gamma}.$$

Here $A(x)$ increases with x, so $Q_*(x)$ decreases with x, and $\log_2 Q_*(1) < 1651$.

Lemma 1 gives $2^{h/2} = q + \delta$ with $0 < \delta < 1$. Write $h = 2m + 1$ with m a nonnegative integer, and let $j = m - k_1$. Lemma 7 gives $k_1 = m - (t-1)/2$, so $j = (t-1)/2$ is a nonnegative integer and $h/2 - k_1 = j + 1/2$. From $qx + 1 = 2^{k_1} y$,

$$\frac{y}{x\,2^j} = 2^{1/2}(1+\rho), \qquad \rho = \frac{1/x - \delta}{q + \delta}.$$

Then $|\rho| < (1 + 1/x)/q$, and therefore $q < (1 + 1/x)/|\rho|$.

If $j \ge k_0(x)$, the argument in the proof of Theorem 3 for length 2 applies with λ_2 in place of 1.465 and with $k_0(x)$ in place of $j_0(x)$. The constant in front of $2^{-\lambda_2 j}$ is 1, and $1/(2A(x)) + 2$ is the one other element, divided by the length, plus the length. That argument gives $q < Q_*(x)$, so $q < 2^{1651}$.

If $j < k_0(x)$, then $j \le 77426$. If $|y - x\,2^j\,2^{1/2}| > 1$, then $|\rho| > 1/(x\,2^j\,2^{1/2})$, so

$$q < (1 + 1/x)\,x\,2^j\,2^{1/2} < 3\,x\,2^j \le 3 \cdot 2^{2559+77426} < 2^{79987}.$$

If $|y - x\,2^j\,2^{1/2}| \le 1$, then $y + x\,2^j\,2^{1/2} \le 2x\,2^j\,2^{1/2} + 1 < 4x\,2^j$. The integers y and $x\,2^j$ satisfy $y^2 \ne 2x^2\,4^j$, since $2^{1/2}$ is irrational.

$$\left|2^{1/2} - \frac{y}{x\,2^j}\right| = \frac{|2x^2\,4^j - y^2|}{x\,2^j\,|y + x\,2^j\,2^{1/2}|} > \frac{1}{4x^2\,4^j}.$$

Hence $|\rho| > 1/(4 \cdot 2^{1/2}\,x^2\,4^j)$ and

$$q < (1 + 1/x) \cdot 4 \cdot 2^{1/2}\,x^2\,4^j \le 8 \cdot 2^{1/2}\,x^2\,4^j < 12\,x^2\,4^j,$$

so $\log_2 q < 2\log_2 x + 2j + 4 < 2\log_2 x + 2k_0(x) + 4$. The quantity $2\log_2 x + 2k_0(x) + 4$ increases with x, and at $x = 2^{2559}$ it is less than 160,000. In both subcases $q < 2^{160{,}000}$.

Appendix C excludes every cycle of length 2 with $q < 2^{160{,}000}$ except $1 \to 3$, $35 \to 99$ and $27 \to 611$. Each of those has its odd elements in $\{1, 3, 27, 35, 99, 611\}$. The supposition $x \le 2^{2559}$ would make the cycle one of those three. Therefore $x > 2^{2559}$. Every odd element is at least x, so every odd element is greater than 2^{2559}. □

The case that remains is $u \ge 3$. Proposition 10 puts $D \ge 15$, with D dividing $2^t - 1$ and $(q + 2^{(h-t)/2})/D$ an odd integer. Lemma 11 gives that this odd integer is greater than 2^{2559}. Proposition 12 gives the lower bounds on q and on x. Lemma 13 turns the lower bound on q into the remaining bounds.

Proposition 12. Let the cycle be other than $1 \to 3$, $35 \to 99$ and $27 \to 611$. Then $q \ge 2^{1{,}071{,}999{,}971}$ and $x > 2^{17{,}631{,}663}$.

Proof. Proposition 10 gives $m \ge 160{,}000$. Appendix C records the first 1,072,000,000 binary places of $\sqrt{2}$. For every integer m with $160{,}000 \le m \le 1{,}071{,}999{,}970$, places $m+1$ through $m+30$ of that expansion are not all 0. Suppose $m \le 1{,}071{,}999{,}970$. The fractional part δ in $2^{h/2} = q + \delta$ is read

from place $m+1$, so $\delta \ge 2^{-30}$. Lemma 7 gives $x = (q + 2^{k_1})/D$ and $2^{k_1} \le q-1$, and $D = \delta(2q+\delta)$, so

$$\delta = \frac{D}{2q+\delta} < \frac{q+2^{k_1}}{2qx} \le \frac{2q-1}{2qx} < \frac{1}{x}.$$

Thus $2^{-30} \le \delta < 1/x$, and $x < 2^{30}$. Lemma 11 gives $x > 2^{2559}$, so $m \ge 1{,}071{,}999{,}971$. Since $m \ge 160{,}000$ and $\sqrt{2}-1 > 2^{-m}$, we have $2^m\sqrt{2} > 2^m + 1$, so $q \ge 2^m + 1$. Therefore $q \ge 2^{1{,}071{,}999{,}971} + 1$.

Suppose $x \le 2^{17{,}631{,}663}$. In the notation of the proof of Lemma 11, $\lambda_2 - \lambda_1 = 0.0597236348$ and $k_0(x) = \max(m_2(x) - 1/2,\, k(x))$. Formula (1.14) of Bauer and Bennett (2002) is linear in the odd part, and the denominator of c_2 contains one factor of the odd part. The other parameters in the proof of Lemma 11 do not depend on x. Thus $c_2(x) = c_2(1)/x$ and $\chi_2(x) = x\,\chi_2(1)$, and

$$k(x) = k(1) + \frac{\log_2 x}{\lambda_2 - \lambda_1}.$$

Equation (1.15) of Bauer and Bennett (2002), at the single prime 2, with $\mathrm{ord}_2(\Omega_1) = 17$ and $\mathrm{ord}_2(a) = 0$, is

$$m_2(x) = \frac{68 \log(x\,\chi_2(1))}{\log \chi_1} + 15.$$

Here log is the natural logarithm, $\log \chi_1 = \log 4.968969066213$, and the same parameters give $\log(\chi_2(1)) = 51.353232\ldots$. The proof of Lemma 11 records $k(1) = 2840.688\ldots$. At $x = 2^{17{,}631{,}663}$ we have $k(x) < 295{,}223{,}704$ and $m_2(x) - 1/2 < 518{,}368{,}291$, so $k_0(x) < 518{,}368{,}291$. Since k_0 increases with the odd part, $k_0(x) < 518{,}368{,}291$ for every odd part at most $2^{17{,}631{,}663}$. Let $j = m - k_1$. Lemma 7 gives $k_1 = m - (t-1)/2$, so $j = (t-1)/2$.

If $j \ge k_0(x)$, the argument in the proof of Lemma 11 gives $q < Q_*(x)$. There $Q_*(x)$ decreases with x and $\log_2 Q_*(1) < 1651$, so $q < 2^{1651}$. If $j < k_0(x)$, then $j \le 518{,}368{,}290$. If $|y - x\,2^j\,2^{1/2}| > 1$, the same argument gives

$$q < 3\,x\,2^j \le 3 \cdot 2^{17{,}631{,}663 + 518{,}368{,}290} = 3 \cdot 2^{535{,}999{,}953} < 2^{535{,}999{,}955}.$$

If $|y - x\,2^j\,2^{1/2}| \le 1$, it gives

$$\log_2 q < 2\log_2 x + 2j + 4 \le 2 \cdot 17{,}631{,}663 + 2 \cdot 518{,}368{,}290 + 4 = 1{,}071{,}999{,}910.$$

In every case $q < 2^{1{,}071{,}999{,}971}$, which contradicts $q \ge 2^{1{,}071{,}999{,}971} + 1$. Therefore $x > 2^{17{,}631{,}663}$. □

Lemma 13. Let the cycle be other than $1 \to 3$, $35 \to 99$, and $27 \to 611$, and write $h = 2m+1$. Then

$$D > 2^{0.52m+1}, \qquad t > 0.52m + \log_2 3, \qquad k_1 < \frac{1.48}{2}\,m - \frac{\log_2 3 - 1}{2}, \qquad x < 2^{0.48m+1/2}.$$

Also $t \ge 557{,}439{,}987$, $k_1 \le \lfloor \log_2 q \rfloor - 278{,}719{,}993$, and $k_2 \ge \lfloor \log_2 q \rfloor + 278{,}719{,}994$.

Proof. Proposition 10 gives $D > 2^{0.52m+1}$ and $t > \log_2(3q/x)$. Lemma 7 gives $x = (q + 2^{k_1})/D$ and $2^{k_1} \le q - 1$. Hence $q + 2^{k_1} \le 2q - 1 < 2q$, so $x < 2q/D$ and $3q/x > 3D/2$. Therefore

$$t > \log_2(3q/x) > \log_2(3D/2) = \log_2 D + \log_2 3 - 1.$$

Since $\log_2 D > 0.52m + 1$,

$$t > 0.52m + \log_2 3.$$

Lemma 7 gives $h = 2k_1 + t$, so $k_1 = (2m + 1 - t)/2 = m - (t - 1)/2$. Thus

$$\begin{aligned} k_1 &< m - \frac{0.52m + \log_2 3 - 1}{2} \\ &= \frac{1.48}{2}\, m - \frac{\log_2 3 - 1}{2}. \end{aligned}$$

Since $q = \lfloor 2^{h/2} \rfloor = \lfloor 2^m \sqrt{2} \rfloor$ and $2^m\sqrt{2}$ is irrational, $q < 2^m\sqrt{2}$ and $\log_2 q < m + 1/2$. Combined with $x < 2q/D$ and $D > 2^{0.52m+1}$,

$$\log_2 x < 1 + \log_2 q - \log_2 D < 1 + (m + \tfrac{1}{2}) - (0.52m + 1) = 0.48m + \tfrac{1}{2},$$

and $x < 2^{0.48m+1/2}$.

Proposition 12 gives $m \geq$ 1,071,999,971. At that value, $0.52m + 1 =$ 557,439,985.92, so $D > 2^{557{,}439{,}985}$. Lemma 7 gives $D \leq 2^t - 1$, so $t >$ 557,439,985.92. Since t is odd, $t \geq$ 557,439,987. For every larger m the same lower bound on D is larger. Since t is odd, $(t-1)/2$ is an integer, and $k_1 = m - (t-1)/2 \leq m -$ 278,719,993. As in the proof of Proposition 10, $\lfloor \log_2 q \rfloor = m$, so $k_1 \leq \lfloor \log_2 q \rfloor -$278,719,993. Also $k_2 = m+(t+1)/2 \geq m+$278,719,994 $= \lfloor \log_2 q \rfloor +$278,719,994. □

9 Longer cycles

Length 3 is the longest length for which every cycle with a constant odd starting value x falls under a root controlled by Lemma 4 or by Corollary 1.4 of [1]. Two thinner classes go further. They depend on ℓ and on h modulo ℓ, so they are a corollary rather than part of Theorem 3.

Corollary 14. Let x_0 be an odd positive integer. There is an integer $Q(x_0, \ell)$, read off from x_0 and ℓ by the comparison in the proof of Theorem 3, with the following property. If x_0 is the starting value of a cycle of length ℓ with halving total h, and either

1. $\ell = 2^a$ with $a \geq 2$ and $h \equiv \ell/2 \pmod{\ell}$, or
2. $\ell = 3 \cdot 2^a$ with $a \geq 1$ and $h \equiv \ell/3$ or $2\ell/3 \pmod{\ell}$,

then $q < Q(x_0, \ell)$.

Proof. In the first family every irrational root that occurs is $\sqrt{2}$. An element aimed at an integer power of 2 is large because it is odd. Corollary 1.4 of [1] supplies the exponent $\mu_2 = 1.465 < \varphi$ with odd part x_0. The comparison in the proof of Theorem 3 then produces a finite check of the exponents below $j_0(x_0)$ and an explicit conflict for the larger exponents.

In the second family the only irrational roots that occur are $2^{1/3}$ and $2^{2/3}$. Length 3 is the case $a = 0$, in which those two residues are every nonzero class modulo 3, and the restriction on h disappears. For $a \geq 1$, Lemma 4 supplies the same exponent $\mu_3 < \varphi$ with odd part x_0 or $3x_0$. The comparison in the proof of Theorem 3 again produces the integer $Q(x_0, \ell)$. □

Length 4 with h odd is not included. The first family uses Corollary 1.4 of [1]. The second uses Lemma 4.

Corollaries 12, 13 and 14 of [6] are the case $x_0 = 1$. Corollary 12 of that paper, and Theorem 1 of that paper for length 3, conclude that no such q exists. Corollary 14 gives a bound in terms of x_0. For $x_0 = 1$ the companion paper already carries out the search that rules out a length-3 cycle. For a general x_0 the bound $Q(x_0, \ell)$ does not by itself rule the cycle out. Corollaries 13 and 14 of [6] keep the constants of those papers (for $\sqrt{2}$, the positive real number $c = 2^{-1.48a}$ and the positive integer $j_0 = 9$; for the cube roots, $j_0 = 11000$). The integer $Q(x_0, \ell)$ still depends on x_0, so it gives no uniform bound on every cycle of that length and residue.

10 A heuristic

A cycle of length $\ell \geq 2$ with h halvings has to satisfy several conditions at once. As above, $q^\ell < 2^h < (q+1)^\ell$, so ℓ does not divide h and $q = \lfloor 2^{h/\ell} \rfloor$ is a single odd integer at least 3. Write $D = 2^h - q^\ell$. Starting from any element x of the cycle, the halving counts are a composition of h into ℓ positive parts, and the partial sums S_j begin at x. Unfolding the cycle relations gives

$$x = \frac{N}{D}, \qquad N := q^{\ell-1} + \sum_{j=1}^{\ell-1} q^{\ell-1-j}\, 2^{S_j}.$$

Thus D divides N, and x is a positive odd integer. Tracing from x recovers those counts exactly, and the ℓ elements are odd and distinct.

If, for a random composition, N is uniformly distributed modulo D, then D divides N with probability $1/D$. Counting cycles up to rotation, the expected number at this pair (h, ℓ) is then about

$$\frac{1}{\ell D}\binom{h-1}{\ell-1}.$$

This is only a heuristic: N is a sparse combination of powers of 2 with coefficients powers of q, so the residue $N \bmod D$ can be constrained, and the remaining conditions only make a cycle rarer.

The estimate is large only when D is small, that is when q^ℓ is extraordinarily close to a power of 2. The known cycles are exactly those coincidences:

$$5^2 = 2^5 - 7, \qquad 5^3 = 2^7 - 3, \qquad 181^2 = 2^{15} - 7.$$

The corresponding expectations are $4/(2 \cdot 7) \approx 0.29$, $15/(3 \cdot 3) = 5/3$, and $14/(2 \cdot 7) = 1$, against one, two, and two cycles found. The only consecutive perfect powers are 8 and 9, of the other sign, so $D \geq 3$ for $\ell \geq 2$, and $D = 3$ and $D = 7$ are as small as occur here.

The next-smallest values of D do not produce cycles. For $q = 11$ and $h = 7$ we have $11^2 = 2^7 - 7$, the same D as for $q = 5$ and $q = 181$, and the estimate is $6/(2 \cdot 7) \approx 0.43$; but $11 + 2^{k_1} \not\equiv 0 \pmod 7$ for every positive $k_1 < 7$, so D never divides N. For $q = 45$ and $h = 11$, likewise $45^2 = 2^{11} - 23$ and $45 + 2^{k_1} \equiv 0 \pmod{23}$ forces $2^{k_1} \equiv 1 \pmod{23}$, hence $11 \mid k_1$, which is impossible for $0 < k_1 < 11$. For $q = 3$ and $h = 5$, $3^3 = 2^5 - 5$ and the estimate is 0.4; the length-3 search in Appendix C finds no cycle. For $q = 3$ and $h = 8$, $3^5 = 2^8 - 13$ and the estimate is 0.54; the length-5 search there

finds no cycle. For $q = 19$ and $h = 17$, $19^4 = 2^{17} - 751$, the fourth-power near miss recorded in the companion paper [6], and the estimate is 0.19; the length-4 search there finds no cycle.

Summing the same estimate over every admissible pair with $2 \le \ell \le 20$ and $h \le 200$ gives a total of about 7 expected cycles, of which about 3 sit in the three known slots and the remaining 4 are spread over pairs that Appendix C and the companion paper [6] have already emptied. For large h a typical D is of size $\ell\delta\, 2^{h(1-1/\ell)}$, with δ the fractional part of $2^{h/\ell}$, and $\binom{h-1}{\ell-1}/D$ decays exponentially in h. Length 1 is different: $D = 2^h - q = 1$ holds for every Mersenne number, and $N = 1$, so the division succeeds and produces $1 \to 1$.

For a cycle at large q there is a further obstruction. A random odd input has $E[v_2(qn+1)] = 2$, while a cycle needs mean $h/\ell \approx \log_2 q$. For $q = 5$ this is 2.32, a mild excess, and three cycles exist. For $q = 181$ it is 7.5, which occurs only because $D = 7$. For larger q the same mean requires a still smaller relative gap, which is the Diophantine condition already used in Theorem 2 of the companion paper and Theorem 2.

11 Open problems

1. **All cycles.** Conjecture 1 is open. Theorem 2 gives, for each $\ell \ge 2$, only finitely many q with a cycle of length ℓ, with no bound and no classification. A bound in terms of a constant odd element, for lengths 2 and 3, is Theorem 3, and for the classes in Corollary 14 it is that corollary. A uniform bound on q remains open. Appendix A records the constraints on $D = 2^h - q^\ell$ for an arbitrary cycle. Appendix B records three further restrictions. Each one leaves that uniform bound open, and leaves open the list of lengths at a single q. Proposition B.2 gives, for each fixed integer V, only finitely many cycles with $\ell/\gcd(h,\ell) \le V$. It gives no bound. A cycle with $\gcd(h,\ell) = 1$ has $\ell/\gcd(h,\ell) = \ell$, and the proposition does not bound those lengths. Proposition 3 produces that bound from its hypothesis, a lower bound of the shape $c \cdot d^{-\mu}$ with exponent at most 1.3247 for every root $2^{r/\ell}$. For length 2, Section 8 leaves the case $u \ge 3$, with $q \ge 2^{1,071,999,971}$ and every odd element greater than $2^{17,631,663}$. The heuristic of Section 10 is not a proof, and the searches leave infinitely many pairs (ℓ, h) unchecked.

A Restrictions on the denominator of any cycle

Throughout this appendix, let the cycle have length $\ell \ge 2$, and let $D = 2^h - q^\ell$. Lemma 7 gives, when $\ell = 2$, that D divides $2^t - 1$ and that $D \equiv 7 \pmod 8$. Proposition A.3 is that divisibility for an arbitrary length.

Proposition A.1. Then $h \ge 4$. The integer D is odd and $D \ge 3$. Also $\gcd(D, q) = 1$. $D \equiv 2^h \pmod q$ and $D \equiv -q^\ell \pmod 8$. In particular, if ℓ is even, then $D \equiv 7 \pmod 8$. If ℓ is even and h is odd, then every prime divisor of D is congruent to 1 or 7 modulo 8.

Proof. Lemma 1 gives $q^\ell < 2^h$. Since $q \ge 3$ and $\ell \ge 2$, we have $2^h > 9$, so $h \ge 4$ and $D > 0$. Thus 8 divides 2^h, and $D \equiv -q^\ell \pmod 8$. The integer q is odd, so $q^2 \equiv 1 \pmod 8$. If ℓ is even, then $q^\ell = (q^2)^{\ell/2} \equiv 1 \pmod 8$, and $D \equiv -1 \equiv 7 \pmod 8$. Here 2^h is even and q^ℓ is odd, so D is odd. Suppose $D = 1$. Then $2^h - q^\ell = 1$, and both exponents are at least 2. Mihăilescu's theorem (2004) [7] says that the only solution of $a^u - b^v = 1$ in integers $a, b > 0$ and $u, v > 1$ is $3^2 - 2^3 = 1$. The

equation $2^h - q^\ell = 1$ would be a solution of the other sign. Thus $D \geq 3$. Also $\gcd(D, q) = \gcd(2^h, q)$. Since q is odd, this greatest common divisor equals 1. Reducing $D = 2^h - q^\ell$ gives $D \equiv 2^h \pmod{q}$.

Suppose ℓ is even and h is odd. Let p be a prime divisor of D. Then $q^\ell = 2^h - D \equiv 2^h \pmod{p}$. Since $\gcd(D, q) = 1$, the prime p does not divide q. Since D is odd, $p \neq 2$, and p does not divide 2^h. The integer ℓ is even, so q^ℓ is a nonzero square modulo p, and therefore 2^h is a nonzero square modulo p. Since h is odd, 2 is a quadratic residue modulo p, and therefore $p \equiv 1 \pmod{8}$ or $p \equiv 7 \pmod{8}$. □

Length 2 has odd h, so every prime divisor of D is congruent to 1 or 7 modulo 8. Lemma 7 also gives that D divides $2^t - 1$.

Proposition A.2. Let $d = \gcd(h, \ell)$. Let $X = 2^{h/d}$ and $Y = q^{\ell/d}$. Then $D = X^d - Y^d$, so $X - Y$ divides D. If $d = 1$, then $X - Y = D$. If $d > 1$, then $X - Y > 1$ and $(X^d - Y^d)/(X - Y) > 1$.

Proof. The integers h/d and ℓ/d are positive, and $X^d - Y^d = 2^h - q^\ell = D$. The difference of powers factors as

$$X^d - Y^d = (X - Y) \sum_{i=0}^{d-1} X^{d-1-i} Y^i.$$

If $d = 1$, the sum equals 1 and $X - Y = D$.

Now suppose $d > 1$. Lemma 1 gives that ℓ does not divide h, so $d < \ell$ and $\ell/d \geq 2$. The same lemma gives $2^h > q^\ell$. Since $q \geq 3$, we have $\log_2 q > 1$, so $h > \ell$ and $h/d > \ell/d \geq 2$. The integer h/d is therefore at least 3, and both exponents h/d and ℓ/d are at least 2. Also $X > Y$, so $X - Y$ is a positive integer. Mihăilescu's theorem (2004) [7] says that the only solution of $a^u - b^v = 1$ in integers $a, b > 0$ and $u, v > 1$ is $3^2 - 2^3 = 1$. The equation $2^{h/d} - q^{\ell/d} = 1$ would be a solution of the other sign. Thus $X - Y > 1$. The sum has $d \geq 2$ terms and its leading term is X^{d-1}. Since $d - 1 \geq 1$ and $X = 2^{h/d} \geq 2^3$, that term is at least 8, and the sum is greater than 1. □

In particular, if $d > 1$, then D is a product of two integers, each greater than 1. Each of the five cycles of length at least 2 in the introduction has $\gcd(h, \ell) = 1$.

Proposition A.3. Let $S_0 = 0$, and let

$$f(z) = \sum_{j=0}^{\ell-1} 2^{S_j} z^{\ell-1-j}.$$

Let R be the resultant of $z^\ell - 2^h$ and $f(z)$. Then D divides R.

Proof. The relation $qx_{i-1} + 1 = 2^{k_i} x_i$ gives, by induction on m,

$$2^{S_m} x_m = x_0 q^m + \sum_{j=0}^{m-1} 2^{S_j} q^{m-1-j}$$

for $1 \le m \le \ell$. For $m = \ell$ we have $x_\ell = x_0$ and $S_\ell = h$, so

$$x_0(2^h - q^\ell) = \sum_{j=0}^{\ell-1} 2^{S_j} q^{\ell-1-j} = f(q).$$

Thus D divides $f(q)$. The same identity $D = 2^h - q^\ell$ gives that D divides $q^\ell - 2^h$.

The resultant R is an integer. The Sylvester matrix of $z^\ell - 2^h$ and $f(z)$ gives polynomials $u(z)$ and $v(z)$ with integer coefficients such that

$$u(z)\, f(z) + v(z)\, (z^\ell - 2^h) = R.$$

Substituting $z = q$ gives $R = u(q)\, f(q) + v(q)\, (q^\ell - 2^h)$. Both terms on the right are divisible by D.

For $\ell = 2$ we have $f(z) = z + 2^{k_1}$, and $R = \pm(2^{2k_1} - 2^h)$. Lemma 7 gives $k_1 < k_2$. Then $2^{2k_1} - 2^h = -2^{2k_1}(2^{k_2-k_1} - 1)$. Proposition A.1 gives that D is odd, so D divides $2^{k_2-k_1} - 1$. □

The last conclusion is the divisibility in Lemma 7.

Proposition A.4. Then

$$\frac{q^{\ell-1}}{x_0} < D < \frac{\ell(q+1)^{\ell-1}}{x_0}.$$

Also

$$\frac{q^{\ell-1}}{D} < x_0 < \frac{\ell(q+1)^{\ell-1}}{D}.$$

Proof. The proof of Proposition A.3 gives $x_0 D = f(q)$, where

$$f(q) = q^{\ell-1} + \sum_{j=1}^{\ell-1} 2^{S_j} q^{\ell-1-j}$$

and $S_0 = 0$. The sum contains the summand $2^{S_1} q^{\ell-2}$, and $S_1 \ge 1$, so $f(q) > q^{\ell-1}$. Thus $D > q^{\ell-1}/x_0$.

The starting value x_0 is the smallest element, and the elements are distinct, so some element is greater than x_0. Lemma 1 gives $2^h = \prod_{i=0}^{\ell-1}(q + 1/x_i)$. Each factor is at most $q + 1/x_0$, and one factor is strictly smaller, so $2^h < (q + 1/x_0)^\ell$. Writing $2^{h/\ell} = q + \delta$ gives $\delta < 1/x_0$, and

$$D = (q+\delta)^\ell - q^\ell < \left(q + \frac{1}{x_0}\right)^\ell - q^\ell.$$

For $1 \le j \le \ell$ we have $x_0^{-j} \le 1/x_0$, so

$$\left(q + \frac{1}{x_0}\right)^\ell - q^\ell = \sum_{j=1}^{\ell} \binom{\ell}{j} q^{\ell-j} x_0^{-j} \le \frac{1}{x_0} \sum_{j=1}^{\ell} \binom{\ell}{j} q^{\ell-j} = \frac{(q+1)^\ell - q^\ell}{x_0}.$$

For $1 \le j \le \ell$,

$$\binom{\ell}{j} = \frac{\ell}{j} \binom{\ell-1}{j-1} \le \ell \binom{\ell-1}{j-1},$$

and $\ell/j < \ell$ for $j = 2$. The summand for $j = 2$ is positive, so $(q+1)^\ell - q^\ell < \ell(q+1)^{\ell-1}$. If $x_0 = 1$, then $x_0^{-j} = 1/x_0$, and the preceding strict inequality gives $D < \ell(q+1)^{\ell-1}/x_0$. If $x_0 > 1$, then $x_0^{-2} < 1/x_0$, and the summand for $j = 2$ is again positive, so

$$\Big(q + \frac{1}{x_0}\Big)^\ell - q^\ell < \frac{(q+1)^\ell - q^\ell}{x_0} < \frac{\ell(q+1)^{\ell-1}}{x_0}.$$

In either case $D < \ell(q+1)^{\ell-1}/x_0$.

The two bounds on D, each divided into $q^{\ell-1}$ or into $\ell(q+1)^{\ell-1}$, give the two bounds on x_0. □

Proposition A.5. Any cycle of length 2 other than $1 \to 3$, $35 \to 99$, and $27 \to 611$ has $D > q^{0.52}$. Any cycle of length 3 with $(q, h) \neq (5, 7)$ has $D \geq q^{4/3}$. In that case $D = 3$.

Proof. For length 2, Proposition 10 gives $h = 2m + 1$ with an integer $m \geq 160{,}000$, $D > 2^{0.52m+1}$, and $q < 2^{m+1}$. Thus $\log_2 q < m + 1$, and

$$\log_2 D > 0.52m + 1 = 0.52(m+1) + 0.48 > 0.52 \log_2 q + 0.48.$$

Hence $D > q^{0.52}$.

For length 3, Lemma 1 gives $q^3 < 2^h$, so $D = 2^h - q^3 > 0$ and $q^3 \neq 2^h$. Bauer and Bennett (2018) [2] prove that if x and n are integers with $x^3 \neq 2^n$ and $(x, n) \neq (5, 7)$, then $|x^3 - 2^n| \geq x^{4/3}$. Taking $x = q$ and $n = h$ gives $D \geq q^{4/3}$, except when $(q, h) = (5, 7)$. In that case $2^7 - 5^3 = 128 - 125 = 3$. □

Proposition A.6. Let $\ell \geq 2$, and write $h = \ell m + r$ with integers $m \geq 0$ and $0 < r < \ell$. Suppose c is a constant real number, μ is a constant real number, and m_0 is a constant integer, with $c > 0$ and $1 < \mu < \ell$, such that every integer $m \geq m_0$ satisfies

$$\Big|2^{r/\ell} - \frac{q}{2^m}\Big| > c \cdot 2^{-\mu m}.$$

Let $\kappa = \ell\, c (2^{r/\ell - 1})^{\mu - 1}$. Then every integer $m \geq \max(m_0, 1)$ satisfies $D > \kappa\, q^{\ell - \mu}$.

Proof. Let $\theta = 2^{r/\ell}$ and let $\delta = 2^{h/\ell} - q$, as in Lemma 1. The hypothesis gives $\delta/2^m > c \cdot 2^{-\mu m}$, so $\delta > c \cdot 2^{(1-\mu)m}$. The proof of Lemma 1 gives $\ell q^{\ell-1}\delta \leq (q+\delta)^\ell - q^\ell$, and $(q+\delta)^\ell = 2^h$, so

$$D \geq \ell q^{\ell-1}\delta > \ell\, c\, q^{\ell-1}\, 2^{(1-\mu)m}.$$

Since $0 < \delta < 1$, we have $q > \theta\, 2^m - 1$. For $m \geq 1$ the inequality $\theta\, 2^{m-1} > 1$ holds, because $\theta > 1$. Thus $\theta\, 2^m - 1 > \theta\, 2^{m-1}$, and $q > \theta\, 2^{m-1}$. Hence $q/2^m > \theta/2 = 2^{r/\ell - 1}$. Since $\mu > 1$,

$$\Big(\frac{q}{2^m}\Big)^{\mu-1} > (2^{r/\ell-1})^{\mu-1}.$$

Therefore

$$\frac{D}{q^{\ell-\mu}} > \ell\, c \Big(\frac{q}{2^m}\Big)^{\mu-1} > \kappa.$$

□

Length 2 is the case $\mu = 1.48$ of Corollary 1.6 of Bauer and Bennett (2002) [1], and Proposition A.5 removes the constant factor κ. Length 3 is the inequality of Bauer and Bennett (2018) [2], whose exponent $4/3$ is smaller than $3 - 1.62$. For $2^{1/4}$ and $2^{3/4}$ this paper supplies no such μ.

Proposition A.7. Let $\ell \geq 2$ be an integer, and let $\varepsilon > 0$ be a real number. Only finitely many cycles of length ℓ satisfy $D \leq q^{\ell-1-\varepsilon}$.

Proof. Write $h = \ell m + r$ with integers $m \geq 0$ and $0 < r < \ell$, and let $\theta = 2^{r/\ell}$. Lemma 1 gives $2^{h/\ell} = q + \delta$ with $0 < \delta < 1$, so

$$\left|\theta - \frac{q}{2^m}\right| = \frac{\delta}{2^m}.$$

The fraction $q/2^m$ is in lowest terms because q is odd. Ridout's theorem (1958) [8], in the S-unit form with $S = \{2\}$, gives that only finitely many such fractions satisfy

$$\left|\theta - \frac{q}{2^m}\right| < 2^{-m(1+\varepsilon)}.$$

There are finitely many residues r. Each fraction determines q and h, and each pair (q, h) has finitely many compositions of h into ℓ positive parts. Thus only finitely many cycles of length ℓ satisfy $\delta < 2^{-\varepsilon m}$.

For every other cycle, $\delta \geq 2^{-\varepsilon m}$. The proof of Lemma 1 gives $D \geq \ell q^{\ell-1}\delta$, so $D \geq \ell q^{\ell-1} 2^{-\varepsilon m}$. Since $q + \delta = \theta\, 2^m$ and $\delta < 1$, we have $2^m < (q+1)/\theta$, and therefore $2^{-\varepsilon m} > \big(\theta/(q+1)\big)^{\varepsilon}$. Hence

$$D > \ell\, \theta^{\varepsilon}\, q^{\ell-1} \big/ (q+1)^{\varepsilon} = \ell\, \theta^{\varepsilon}\, q^{\ell-1-\varepsilon} \Big(\frac{q}{q+1}\Big)^{\varepsilon}.$$

The factor $\big(q/(q+1)\big)^{\varepsilon}$ tends to 1 as q grows, and $\ell\, \theta^{\varepsilon} > 1$. Thus $D > q^{\ell-1-\varepsilon}$ whenever q is large enough, for a bound depending only on ℓ, r, and ε. The remaining cycles, those with q below that bound, are finite in number. □

The exponent $\ell - 1 - \varepsilon$ may be taken arbitrarily close to $\ell - 1$. The appeal to Ridout's theorem (1958) [8] gives no bound on the exceptional cycles. Proposition A.4 supplies the matching upper bound $D < \ell(q+1)^{\ell-1}$, since $x_0 \geq 1$.

B Finiteness past Theorem 2

Proposition B.1. Let q be a Mersenne number, and write $q = 2^e - 1$ for an integer $e \geq 2$. Let $\ell \geq 2$. On a cycle of length ℓ, $\ell > 1/\log_2(1 + 1/q)$ and $\ell/\gcd(h, \ell) > 1/\log_2(1 + 1/q)$.

Proof. Lemma 1 gives $q^{\ell} < 2^h < (q+1)^{\ell}$. Here $(q+1)^{\ell} = 2^{e\ell}$, so $h \leq e\ell - 1$, and $h > \ell \log_2 q$. Suppose $\ell \log_2(1 + 1/q) \leq 1$. Then

$$\ell \log_2 q = e\ell - \ell \log_2(1 + \tfrac{1}{q}) \geq e\ell - 1,$$

so $h > \ell \log_2 q \geq e\ell - 1 \geq h$, which is impossible. Thus $\ell \log_2(1 + 1/q) > 1$.

Since $h \le e\ell - 1$ and $h > \ell \log_2 q$, the positive integer $s = e\ell - h$ satisfies

$$1 \le s < \ell \log_2(1 + \tfrac{1}{q}).$$

Let $v = \ell / \gcd(h, \ell)$. Then $\gcd(h, \ell) = \gcd(s, \ell) \le s$, so v satisfies

$$v \ge \frac{\ell}{s} > \frac{1}{\log_2(1 + 1/q)}.$$

□

For $q = 3$ we have $2^{1/3} < 4/3 < \sqrt{2}$, because $2 < 64/27$ and $16/9 < 2$. Thus $2 < 1/\log_2(4/3) < 3$, and the proposition excludes a cycle of length 2. It does not exclude length 3. The same lower bound is larger than $q \log 2$, because $\log(1 + 1/q) < 1/q$, so the excluded lengths grow with q. Once $1/\log_2(1 + 1/q) \ge V$, a cycle for this q has $\ell / \gcd(h, \ell) > V$, and Proposition B.2 does not apply to it.

Proposition B.2. Fix an integer $V \ge 2$. Only finitely many cycles satisfy $\ell / \gcd(h, \ell) \le V$.

Proof. The odd elements of a cycle of length $\ell \ge 2$ are distinct odd positive integers, so

$$\sigma \le \sum_{k=0}^{\ell-1} \frac{1}{2k+1}.$$

The summand $f(t) = 1/(2t+1)$ is strictly decreasing for $t \ge 0$. For each integer $k \ge 1$ and every $t \in [k-1, k)$, we have $f(t) > f(k)$, so

$$\sum_{k=1}^{\ell-1} \frac{1}{2k+1} < \int_0^{\ell-1} \frac{dt}{2t+1} = \tfrac{1}{2} \log(2\ell - 1).$$

The $k = 0$ term equals 1, and therefore $\sigma < 1 + \frac{1}{2} \log(2\ell - 1)$. Let Σ be the positive real number $1 + \frac{1}{2} \log(2\ell - 1)$. Lemma 1 gives $(q + \delta)^\ell = \prod_i (q + 1/x_i)$, so

$$\ell \log\Big(1 + \frac{\delta}{q}\Big) = \sum_i \log\Big(1 + \frac{1}{q x_i}\Big) < \frac{\sigma}{q}.$$

For $u > 0$, $\log(1 + u) > u/(1 + u)$. With $u = \delta/q$,

$$\frac{\ell \delta}{q + \delta} < \frac{\sigma}{q},$$

and therefore $\delta(\ell - \sigma/q) < \sigma$. The series $e^u > u^2/2$ at $u = 2\sqrt{\ell}$ gives $e^{2\sqrt{\ell}} > 2\ell > 2\ell - 1$, so $\log(2\ell - 1) < 2\sqrt{\ell}$ and $\Sigma < 1 + \sqrt{\ell}$. Also $\Sigma < \ell$. Since $q \ge 3$, we have $\sigma < q\ell/2$, hence $\ell - \sigma/q > \ell/2$ and

$$\delta < \frac{2\sigma}{\ell} < \frac{2\Sigma}{\ell}.$$

Let ε be the constant rational $1/10$. Write $h = \ell m + r$ with integers $m \ge 0$ and $0 < r < \ell$, and set $\theta = 2^{r/\ell}$. Let $v = \ell / \gcd(h, \ell)$. The hypothesis is $v \le V$. Let $g = \gcd(r, \ell)$. Then $\gcd(h, \ell) = g$ and

$\theta = 2^{\rho/v}$ with $\rho = r/g$, so $\gcd(\rho, v) = 1$ and $1 \le \rho \le v-1$. In particular $1 < \theta < 2$ and $2^m\theta = q + \delta$, so $2^m = (q+\delta)/\theta < q+1$. The integer 2^m is therefore at most q. The hypothesis $v \le V$ leaves only finitely many such real numbers θ. Each is an algebraic irrational. Let Φ be the set of real numbers $2^{\sigma_0/v_0}$ with integers v_0, σ_0 such that $2 \le v_0 \le V$ and $1 \le \sigma_0 \le v_0 - 1$. That set is finite. Each element lies strictly between 1 and 2. A rational value would be an integer, so each element is an algebraic irrational.

The series for $e^{7/10}$ through the term of degree 4 equals $482921/240000 > 2$, so $\log 2 < 7/10$. The series for $\log(1 + 1/4)$ alternates, and the term after $(1/4)^3/3$ is negative, so $\log(5/4) < 1/4 - 1/32 + 1/192 = 43/192$. Hence $\log 10 = 3\log 2 + \log(5/4) < 21/10 + 43/192 = 2231/960$. The partial sum of e through $1/8!$ equals $109601/40320$. The cross product $109601 \cdot 500 - 40320 \cdot 1359 = 5620$ is positive, so $e > 2718/1000$ and $e^2 > 7387524/1000000$. For $0 < z \le 1/25$, the remainder of e^z after the quadratic term is at most $z^3/(6(1-z)) < z^2/2$, so $e^z < 1 + z + z^2$. Thus $e^{1/25} < 651/625$ and $7e^{1/25} < 4557/625$. The product $4557 \cdot 1600 = 7291200 < 7387524$, so $4557/625 < 7387524/1000000 < e^2$. Hence $e^{49/25} > 7$ and $\log 7 < 49/25$. Also $(6/5)^2 = 36/25$ and $(6/5)^4 = 1296/625 > 2$, so $(6/5)^8 > 4 > 3$ and $3^{1/10} < 6/5$. The same fourth power gives $(6/5)^4 > 2$, so $2^{1/10} < 6/5$ and $2^{1/5} < 6/5$.

The length is short when $q < 4\ell$. Suppose $\delta \ge (q+1)^{-\varepsilon}$ and $q < 4\ell$. Then $q + 1 \le 4\ell$ and $\ell < 2\Sigma\,(4\ell)^{1/10}$. Here $(4\ell)^{1/10} = 2^{1/5}\ell^{1/10} < (6/5)\ell^{1/10}$, and $\log(2\ell - 1) < 7/10 + \log \ell$, so $\Sigma < 27/20 + \frac{1}{2}\log \ell$. Therefore

$$2\Sigma\,(4\ell)^{1/10} < \frac{12}{5}\Big(\frac{27}{20} + \tfrac{1}{2}\log \ell\Big)\ell^{1/10} = \Big(\frac{81}{25} + \frac{6}{5}\log \ell\Big)\ell^{1/10}.$$

The integer comparison $5^{10} = 9765625 > 7340032 = 7 \cdot 2^{20}$ is $(28/5)^{10} < 7^9$, so $7^{9/10} > 28/5$. Combined with $\log 7 < 49/25$,

$$7^{9/10} - \tfrac{6}{5}\log 7 > \frac{28}{5} - \frac{6}{5} \cdot \frac{49}{25} = \frac{406}{125} > \frac{81}{25}.$$

The derivative of $\ell^{9/10} - (6/5)\log \ell$ equals $(9\ell^{9/10} - 12)/(10\ell)$, which is positive for $\ell \ge 7$ because $7^{9/10} > 4/3$. Thus $\ell^{9/10} - (6/5)\log \ell > 81/25$ for every real $\ell \ge 7$, and $2\Sigma\,(4\ell)^{1/10} < \ell$. This contradicts $\ell < 2\Sigma\,(4\ell)^{1/10}$. Hence $\ell \le 6$. Then $q < 24$. Lemma 1 leaves finitely many integers h, and each such h has finitely many compositions into ℓ positive parts.

A small fractional part. Suppose $\delta < (q+1)^{-\varepsilon}$. Then $\delta < (2^m)^{-\varepsilon}$, because $2^m \le q < q+1$. The fraction $q/2^m$ is in lowest terms, and

$$\Big|\theta - \frac{q}{2^m}\Big| = \frac{\delta}{2^m} < (2^m)^{-1-\varepsilon}.$$

Apply the S-unit form of Ridout's theorem (1958) [8], with $S = \{2\}$ and this ε, to each element of the finite set of roots $2^{\rho/v}$. Only finitely many reduced fractions meet the inequality. For each such fraction the positive real number $\delta_* = 2^m\theta - q$ is positive, since θ is irrational. A cycle producing that fraction has $\delta = \delta_*$, so $\ell < 2\Sigma/\delta_*$. With $c = 2/\delta_*$ and $t = \sqrt{\ell}$, the bound $\Sigma < 1 + \sqrt{\ell}$ gives $t^2 < c(1+t)$, hence $t < c + 1$ and $\ell < (c+1)^2$. Each such fraction therefore allows only finitely many lengths. Each pair (q, ℓ) again leaves finitely many compositions.

A large fractional part. Suppose now that $\delta \ge (q+1)^{-\varepsilon}$. Then $\ell < 2\Sigma\,(q+1)^{\varepsilon}$. The case $q < 4\ell$ has already been bounded, so assume $q \ge 4\ell$. Lemma 2 applies. The proof of that lemma records

$e^{1/4} < 4/3$. Since $\ell/q \le 1/4$ and $\delta \le e^{\ell/q}/a$ by Lemma 1,

$$a \le \frac{e^{\ell/q}}{\delta} \le e^{1/4}(q+1)^{\varepsilon} < \tfrac{4}{3}(q+1)^{\varepsilon}.$$

Let y be the positive integer $\min(x_1, \dots, x_{\ell-1})$. Then $y > a$.

First let $q < 10^{300}$. Then $q+1 \le 10^{300}$ and $(q+1)^{1/10} \le 10^{30}$, so $\ell < 2\Sigma \cdot 10^{30}$. Let Σ_* be the positive real number $1 + \frac{1}{2}\log(2\ell)$. Then $\Sigma < \Sigma_*$, and the derivative of ℓ/Σ_* equals $(\Sigma_* - 1/2)/\Sigma_*^2 > 0$, so ℓ/Σ_* is strictly increasing for $\ell > 1$. At $\ell = 10^{32}$,

$$\log(2\ell) = \log 2 + 32 \log 10 < \tfrac{7}{10} + 32 \cdot \frac{2231}{960} = \tfrac{7}{10} + \frac{2231}{30} < 76,$$

so $\Sigma_* < 39$ and $10^{32}/\Sigma_* > 10^{32}/39 > 2 \cdot 10^{30}$. For every $\ell \ge 10^{32}$ one therefore has $\ell > 2 \cdot 10^{30}\, \Sigma_* > 2 \cdot 10^{30}\, \Sigma$, a contradiction. Thus $\ell < 10^{32}$. Both q and ℓ are bounded, and each pair leaves finitely many compositions.

Now let $q \ge 10^{300}$. Then $q + 1 < 2q$, so $(q+1)^{1/10} < (2q)^{1/10} < (6/5) q^{1/10}$ and $\ell < (12/5)\Sigma\, q^{1/10}$. The series for e is strictly less than $1 + 1 + \sum_{k \ge 1} 2^{-k} = 3$, so $e < 3$, $e^2 < 9 < 10$ and $\log 10 > 2$. Suppose $\ell \ge q$. Then $\Sigma < 1 + \sqrt{\ell}$ and $q \le \ell$ give $\ell < (12/5)(1 + \sqrt{\ell})\, \ell^{1/10}$, so $\ell^{9/10} < (12/5)(1 + \sqrt{\ell})$. For $\ell \ge 1$ we have $1 + \sqrt{\ell} \le 2\sqrt{\ell}$, hence $\ell^{2/5} < 24/5$ and $\ell < (24/5)^{5/2}$. Here $(24/5)^2 = 576/25 < 24$ and $24^{1/2} < 5$, so $(24/5)^{5/2} < 120$. This contradicts $\ell \ge 10^{300}$. Thus $\ell < q$. Then $\log(2\ell - 1) < 7/10 + \log q$ and $\Sigma < 27/20 + \frac{1}{2}\log q < \log q$, because $\log q > 600 > 27/10$. Hence $\ell < (12/5)(\log q)\, q^{1/10}$.

Let i be an index with $x_i = y$, and write $ir = \ell t + s$ with integers $t \ge 0$ and $0 \le s < \ell$. Let φ be the positive real number $2^{s/\ell}$, and let F_i be the rational of Lemma 2. Set $G = F_i/2^t$. Then $\theta_i = 2^t \varphi$ and $|\varphi - G| \le 2i\varphi/(aq)$.

Suppose $y \le q^{4/5}$ and $s = 0$. Then $\varphi = 1$ and $\theta_i = 2^t$. Let $b = S_i - im$ and $e = \max(im - S_i,\, 0)$. Lemma 2 and $q \ge 4\ell > 4i$ give $|\theta_i - F_i| < \theta_i/2$, so $2^{t-1} < F_i < (3/2)2^t$. The rational F_i is not equal to 2^t: otherwise $y = a$, contradicting distinctness. If $b > t$, then $F_i = y\, 2^b/a$ and $|a - y\, 2^{b-t}| \le 2i/q < 1/2$. The integer $y\, 2^{b-t}$ is even and a is odd, so those two integers differ by at least 1, which is impossible. If $0 \le b \le t$, the same multiplication gives $|a\, 2^{t-b} - y| \le 2i\, 2^{t-b}/q$. The left side is a positive integer, so $q \le 2i\, 2^{t-b}$. From $F_i > 2^{t-1}$ we have $2^{t-b} < 2y/a$, hence $q < 4iy \le 4\ell\, q^{4/5}$ and $q^{1/5} < 4\ell$. Then $q^{1/10} < (48/5)\log q$. For $q \ge 10^{300}$ the left side is 10^{30} and the right side is less than $10 \cdot 300 \cdot 2231/960 < 10^4$, a contradiction. The case $e \ge 1$ is the same comparison: Lemma 2 supplies $2^e \ge q/(2i\theta_i)$ and $F_i > \theta_i/2$, so $y > aq/(4i)$ and again $q < 4\ell\, q^{4/5}$.

Suppose $y \le q^{4/5}$ and $s \ge 1$. Then $\varphi > 1$. Also $\varphi = 2^{\sigma_0/v}$ for the integer $\sigma_0 = (i\rho \bmod v)$, so $\varphi \in \Phi$. Since $q \ge 4\ell > 4i$ and $a \ge 1$, we have $|\varphi - G| < \varphi/2$, hence $G > \varphi/2 > 1/2$ and $G < 3\varphi/2 < 3$. Let $E = im - S_i + t$, so $G = y/(a\, 2^E)$. Write $G = P/d$ in lowest terms. The odd part of d divides a, so $d \cdot |d|_2 \le a$. With $S = \{2\}$, the left side of the general form of Ridout's theorem is

$$d^2\, |\varphi - G| \cdot |d|_2 \le d \cdot a \cdot \frac{4\ell}{aq} = \frac{4\ell\, d}{q},$$

where $2i\varphi < 4\ell$. If $E \ge 0$, then $a\, 2^E = y/G < 2y$, so $d < 2y$ and the left side is less than $8\ell y/q$. If $E < 0$, then $d \le a < y$ and the left side is again less than $8\ell y/q$. In both cases $\max(|P|,\, d) < 3y$: the

numerator is y when $E \ge 0$, and when $E < 0$ the identity $y\,2^{-E} = Ga < 3a$ bounds the numerator by $3a < 3y$. The inequality $8\ell y/q < (3y)^{-\varepsilon}$ therefore puts G under the general form. It rearranges as $8\ell\,3^{1/10}\,y^{11/10} < q$. Since $y \le q^{4/5}$ and $(4/5)(11/10) = 22/25$, the most demanding value is the right endpoint, and it is enough that $8\ell\,3^{1/10} < q^{3/25}$. Here $3^{1/10} < 6/5$ and $\ell < (12/5)(\log q)\,q^{1/10}$, so the left side is less than $(576/25)(\log q)\,q^{1/10}$. The bound $\log q < 300 \cdot 2231/960 < 700$ and $q^{1/50} = 10^6$ give

$$\frac{576}{25}\log q < 24 \cdot 700 < 10^6 = q^{1/50}.$$

Thus $8\ell y/q < (3y)^{-\varepsilon} < \max(|P|,\, d)^{-\varepsilon}$. Apply the general form of Ridout's theorem (1958) [8], with $S = \{2\}$ and $\varepsilon = 1/10$, to each element of Φ. Only finitely many reduced fractions meet it. For each such fraction $|\varphi - P/d|$ is a positive real number, because φ is irrational. The bound $|\varphi - G| \le 4\ell/(aq)$ then gives q at most a constant multiple of ℓ. Combined with $\ell < (12/5)(\log q)\,q^{1/10}$, the real number $q^{9/10}/\log q$ is bounded, and so is q. Each of those finitely many q leaves finitely many lengths and finitely many compositions.

Suppose finally that $y > q^{4/5}$. Then $1/y < q^{-4/5}$. Also $2\ell < q^{1/5}$, because $(24/5)\log q < q^{1/10}$ for $q \ge 10^{300}$, so $2\ell/q < q^{-4/5}$ and

$$\bigl|\theta - W\bigr| \le \frac{2}{q}\Bigl(\frac{1}{y} + \frac{2\ell}{q}\Bigr) < 4q^{-9/5},$$

where $W = (\ell aq + 1)/(\ell a \cdot 2^m)$ is the rational of Lemma 2. The numerator $\ell aq + 1$ is congruent to 1 modulo ℓa, so in lowest terms $W = P/d$ with $d = \ell a \cdot 2^e$ and $e = \max(m - v_2(\ell aq + 1),\, 0)$. With $S = \{2\}$, the left side of the general form equals $(\ell a)^2 2^e\,|\theta - W| \le (\ell a)^2 q\,|\theta - W|$. From $a < (4/3)(q+1)^{1/10} < (8/5)q^{1/10}$ and $\ell < (12/5)(\log q)\,q^{1/10}$,

$$(\ell a)^2 < \frac{9216}{625}(\log q)^2\,q^{2/5}.$$

Multiplying by $4q^{-4/5}$ gives a factor $36864/625 < 59$, so

$$(\ell a)^2 q \cdot 4q^{-9/5} < 59\,(\log q)^2\,q^{-2/5}.$$

Since $q + 1 < 2q$,

$$\max(|P|,\, d) < \ell a \cdot 2q + 1 < \frac{192}{25}(\log q)\,q^{6/5} + 1 < 16(\log q)\,q^{6/5}.$$

The right side of the general form is therefore greater than $(16\log q)^{-1/10} q^{-3/25}$. Comparing the two sides, it is enough that $59 \cdot 2 \cdot (\log q)^{21/10} < q^{7/25}$, because $16^{1/10} < 2$ and $2/5 - 3/25 = 7/25$. For $q \ge 10^{300}$ we have $\log q < 700$, so $(\log q)^{21/10} < 700^3 = 343000000 < 10^9$, and $q^{7/25} = 10^{84}$. Thus $59 \cdot 2 \cdot (\log q)^{21/10} < 10^{84}$. Apply the general form of Ridout's theorem (1958) [8], with $S = \{2\}$ and $\varepsilon = 1/10$, to each root $\theta = 2^{\rho/v}$. Only finitely many reduced fractions meet it. For each such fraction $|\theta - W|$ is a positive real number. Lemma 2 and $y \ge 1$ give $|\theta - W| \le 2(1 + 2\ell)/q$, so q is at most a constant multiple of ℓ. Combined with $\ell < (12/5)(\log q)\,q^{1/10}$, the real number $q^{9/10}/\log q$ is bounded, and so is q. Each of those finitely many q leaves finitely many lengths and finitely many compositions. □

The appeal to Ridout's theorem (1958) [8] gives no bound on q. Since $\gcd(h, \ell) \ge 1$, every cycle of length at most V satisfies $\ell/\gcd(h, \ell) \le V$. Proposition B.2 includes those cycles, and it includes a

cycle of length greater than V whenever $\gcd(h,\ell) \geq \ell/V$. The hypothesis still requires $\ell/\gcd(h,\ell)$ to be at most a fixed integer. Proposition B.1 gives $\ell/\gcd(h,\ell) > 1/\log_2(1+1/q)$ whenever q is a Mersenne number, so no single integer bounds $\ell/\gcd(h,\ell)$ for every such q. If $\gcd(h,\ell) = 1$, then $\ell/\gcd(h,\ell) = \ell$. The proposition does not bound those lengths. In particular it does not bound the length of a cycle at $q = 3$. At each fixed q, Theorem 2 still supplies finitely many cycles of each fixed length, and the argument above supplies finitely many cycles with $\ell/\gcd(h,\ell) \leq V$.

Proposition B.3. Let S be a finite set of primes containing 2. Only finitely many cycles of length $\ell \geq 2$ have every prime factor of $q(2^h - q^\ell)$ in S.

Proof. Let $D = 2^h - q^\ell$. Lemma 1 gives $D > 0$. Every prime factor of qD lies in S, and $2 \in S$, so the positive rational numbers $q^\ell/2^h$ and $D/2^h$ are S-units. Their sum is 1. Evertse (1984) [5] proves that $x + y = 1$ has at most $3 \times 7^{d+2s}$ solutions in S-units of a number field of degree d. In that bound, s is the number of places, including the infinite places. Let r be the number of primes in our set S. Here the field is the field of rational numbers, so $d = 1$, and the places are the infinite place together with those r primes, so $s = r + 1$. The number of solutions is at most $3 \times 7^{2r+3}$. Each cycle in the proposition produces one of those solutions. For each solution the integers 2^h and q^ℓ are determined. A constant positive integer has finitely many representations as q^ℓ with $\ell \geq 2$ and q an odd positive integer, since ℓ must divide every exponent in the prime factorization of that integer. Each pair (q,ℓ) leaves finitely many compositions of h into ℓ positive parts. □

The five cycles of length at least 2 in the introduction have $q(2^h - q^\ell)$ divisible only by primes in $\{2, 3, 5, 7, 181\}$. For that set we have $r = 5$, and the bound of Evertse (1984) [5] is $3 \times 7^{13} = 290667031221$. The bound counts solutions of $x + y = 1$. Several cycles may produce the same solution. A cycle for which q or $2^h - q^\ell$ has any other prime factor is not included. For length 2, Lemma 7 gives that D divides $2^t - 1$, so the prime factors of D need not lie in a constant set. A Mersenne number $q = 2^e - 1$ has $q + 1 = 2^e$ at length 1, and those values of q do not all have their prime factors in one constant finite set.

C Computations

Lemma 1 gives $q^\ell < 2^h < (q+1)^\ell$ for a cycle of length $\ell \geq 2$, so $\ell \nmid h$ and $q = \lfloor 2^{h/\ell} \rfloor$. A search of every $h \leq H$ with $\ell \nmid h$ therefore excludes every such cycle with $q < 2^{\lfloor H/\ell \rfloor}$: it would have $2^h < (q+1)^\ell \leq 2^{\ell\lfloor H/\ell \rfloor} \leq 2^H$, so $h \leq H$.

Any cycle. For each ℓ from 2 to 9, every h up to the value in the table was tested, for odd elements that need not include 1. The only cycles found are those of Conjecture 1.

Any cycle: only those of Conjecture 1

length ℓ	h at most	for q below
2	320,000	$2^{160,000}$
3	7,000	$2^{2,333}$
4	5,500	$2^{1,375}$
5	4,500	2^{900}
6	1,400	2^{233}
7	950	2^{135}
8	470	2^{58}
9	320	2^{35}

Length 2 finds $1 \to 3 \to 1$ at $q = 5$, and $27 \to 611 \to 27$ and $35 \to 99 \to 35$ at $q = 181$. Length 3 finds $13 \to 33 \to 83 \to 13$ and $17 \to 43 \to 27 \to 17$ at $q = 5$. Lengths 4 through 9 find no cycle.

A constant q. Here q is constant and the length varies. The only possible h are the integers satisfying $q^\ell < 2^h < (q+1)^\ell$; if there is none, that length has no cycle. For $q = 5$, lengths 2 through 20 produce only $1 \to 3 \to 1$ and the two cycles of length 3 above. For each odd q from 7 through 25, lengths 2 through 16 produce no cycle. The cycles at $q = 181$ were found in the length-2 row above, at $h = 15$, not in this search.

Cycles at a constant q

q	lengths	cycles found
5	2 through 20	$1 \to 3 \to 1$; $13 \to 33 \to 83$; $17 \to 43 \to 27$
7	2 through 16	none
9	2 through 16	none
11	2 through 16	none
13	2 through 16	none
15	2 through 16	none
17	2 through 16	none
19	2 through 16	none
21	2 through 16	none
23	2 through 16	none
25	2 through 16	none

The bounds $Q_3(x)$ and $Q_2(x)$. For every odd $x \le 35$, the values of $Q_{3,0}(x)$ were computed with upper limit 11000 on j. For every odd $x \le 35$, the odd parts x and $3x$ have $s' \le 105$ and $v_3 \le 4$. Since $66 \cdot 105 < 2^{13}$ and $(5/4)^4 > 2$,

$$\frac{\log(66 \cdot 105)}{\log(5/4)} < \frac{13 \log 2}{\log(5/4)} = 52 \cdot \frac{\log 2}{\log((5/4)^4)} < 52,$$

so $21 \cdot 52 + 5 = 1097 < 11000$ and $J_3(x) = 11000$. The gate 2^{11001} is then smaller than $Q_{3,*}(x)$ and does not change the maximum. The quantity $Q_{3,*}(x)$ is smallest at $x = 3$, where $\log_2 Q_{3,*}(3) = 17906.510814$, so $Q_{3,*}(x) > 2^{17906} > 2^{11001}$ for every odd positive integer x.

Powers of two above $Q_{3,*}(x)$. The integer $\lceil Q_{3,*}(x) \rceil$ is the sharper bound, and it has thousands of digits. The table uses the next power of 2 above $Q_{3,*}(x)$, read off from the logarithm. For $x = 5$ we have $\log_2 Q_{3,*}(5) = 18213.724287504393$, so $\lceil Q_{3,*}(5) \rceil$ has 5483 digits and begins 753951296 ...,

while the table entry is 2^{18214}. The same rounding, $2^{\lceil \log_2 \cdot \rceil}$, is used for $Q_{3,0}(x)$, $Q_{2,*}(x)$ and $Q_2(x)$.

For every odd x from 1 to 35, the entry $2^{\lceil \log_2 Q_{3,*}(x) \rceil}$ is:

x		x		x	
1	2^{17990}	13	2^{18387}	25	2^{18514}
3	2^{17907}	15	2^{18190}	27	2^{18304}
5	2^{18214}	17	2^{18439}	29	2^{18543}
7	2^{18273}	19	2^{18460}	31	2^{18556}
9	2^{18094}	21	2^{18255}	33	2^{18344}
11	2^{18356}	23	2^{18497}	35	2^{18580}

Rows in which 3 divides x sit below their neighbors because $s' = x/\gcd(x, 3)$. The row $x = 3$ has $s' = 1$ and $c_3(x) = c_3(3)$. The value $c_3(3)$ is the place this table uses Lemma 4 past odd part 1 or 3.

Powers of two above $Q_{3,0}(x)$. For the same odd x, with the maximum taken over $0 \le j < 11000$, the entry $2^{\lceil \log_2 Q_{3,0}(x) \rceil}$ is:

x		x		x	
1	2^{11003}	13	2^{11007}	25	2^{11009}
3	2^{11002}	15	2^{11006}	27	2^{11006}
5	2^{11006}	17	2^{11009}	29	2^{11012}
7	2^{11006}	19	2^{11009}	31	2^{11008}
9	2^{11006}	21	2^{11006}	33	2^{11005}
11	2^{11005}	23	2^{11006}	35	2^{11008}

In every row the exponent j that attains the maximum lies in $\{10997, 10998, 10999\}$. For these eighteen values, $Q_{3,0}(x) < Q_{3,*}(x)$. That comparison is checked for odd $x \le 35$. It is not proved for every odd x. From $t^3 - 2^r$, if the distance from $s'\,2^{j+r/3}$ to an integer is positive, that distance is greater than $1/(Q^2 \cdot 3(2^{2/3} + 1)^2)$ once the integer is within 1 of $s'\,2^{j+r/3}$, with $Q = s'2^j$. With $j \le 10999$ and $s' \le x$ this only yields $Q_{3,0}(x) < x^3\,2^{33003}$. At $x = 1$ the two logarithms, of this upper estimate and of $Q_{3,*}(1)$, are 33003 and 17989. The estimate $x^3\,2^{33003}$ stays above $Q_{3,*}(x)$ until x is about 2^{67}. Past that size, $Q_{3,*}(x)$ grows like x^{1/γ_3} with $1/\gamma_3 \approx 229$, while $x^3\,2^{33003}$ grows like x^3.

The value $Q_3(1)$. Here $s' = 1$, so $J_3(1) = 11000$. The maximum $Q_{3,0}(1)$ is less than 2^{11003}, and $Q_{3,*}(1)$ is less than 2^{17990}, so

$$Q_3(1) = Q_{3,*}(1) < 2^{17990}.$$

Each h gives one candidate $q = \lfloor 2^{h/3} \rfloor$. The bound $q < 2^{17990}$ means it is enough to check $h < 53970$. The companion paper checks every $h \le 200{,}000$ and finds no cycle, so that search includes the range. Theorem 1 of [6] is this case.

The proof of that theorem in the companion paper reaches the larger cutoff $q < 2^{22003}$. There the symbol x is the element after 1. The argument asks for that element to clear 2^{11001}, using a lower bound of the shape $\sqrt{q/2}$. The integer $Q_3(1)$ does not use that lower bound. Once $q > Q_{3,0}(1)$, an exponent below 11000 is already impossible, and the conflict is $Q_{3,*}(1)$.

Powers of two above $Q_{2,*}(x)$. This integer is used in the proof and is at most $Q_2(x)$. As $x \to \infty$, $\log_2 Q_{2,*}(x)$ decreases to $\mu_2/\gamma_2 \approx 6.733$. The row $x = 11$ has logarithm 7.013, so that one ceiling moves if the fourth digit of the exponent 1.465 is lower. For odd x from 1 to 35, the entry $2^{\lceil \log_2 Q_{2,*}(x) \rceil}$ is 2^{11} at $x = 1$, then 2^8 for $x = 3, 5, 7, 9, 11$, and 2^7 for every odd x from 13 to 35.

Values of $Q_2(x)$. For these eighteen values the last failure of $|\sqrt{2} - p/(x\,2^j)| > 2^{-1.465j}$ lies at $j \le 21$, and a check through $j < 53620$ finds no later failure. The entry is $2^{\lceil \log_2 Q_2(x) \rceil}$.

x	$j_0(x)$	winning j	$Q_2(x)$ below
1	9	7	2^{15}
3	12	11	2^{18}
5	22	17	2^{33}
7	12	10	2^{18}
9	12	11	2^{18}
11	13	12	2^{19}
13	14	9	2^{21}
15	22	17	2^{32}
17	15	13	2^{23}
19	17	14	2^{25}
21	17	12	2^{25}
23	16	14	2^{23}
25	22	17	2^{32}
27	15	14	2^{22}
29	17	16	2^{25}
31	17	16	2^{25}
33	15	13	2^{22}
35	22	17	2^{32}

The cycle $27 \to 611 \to 27$ has $q = 181 < 2^{22}$. The cycle $35 \to 99 \to 35$ has $q = 181 < 2^{32}$.

Digits of $\sqrt{2}$. Let $N = 1{,}072{,}000{,}000$ and let $I = \lfloor 2^N \sqrt{2} \rfloor$. The integer I was computed and checked by $I^2 < 2^{2N+1} < (I+1)^2$. Place 1 is the bit of $\sqrt{2}$ just below the integer bit. For a cycle of length 2 we have $h = 2m + 1$ and $2^{h/2} = q + \delta$, so the binary digits of δ are places $m + 1$, $m + 2$, ... of $\sqrt{2}$. Proposition 10 covers every $m < 160{,}000$. From $m = 160{,}000$ through $m = 1{,}071{,}999{,}970$, the longest run of consecutive zeros in those places has length 29, at $m = 1{,}005{,}210{,}007$.

References

[1] M. Bauer and M. A. Bennett, Applications of the hypergeometric method to the generalized Ramanujan–Nagell equation, Ramanujan J. 6 (2002), 209–270, doi:10.1023/A:1015779301077.

[2] M. Bauer and M. A. Bennett, Ramanujan–Nagell cubics, Rocky Mountain J. Math. 48 (2018), 385–412, doi:10.1216/rmj-2018-48-2-385.

[3] F. Beukers, On the generalized Ramanujan–Nagell equation I, Acta Arith. 38 (1980/81), 389–410, doi:10.4064/aa-38-4-389-410.

[4] R. E. Crandall, On the "$3x + 1$" problem, Math. Comp. 32 (1978), 1281–1292, doi:10.1090/S0025-5718-1978-0480321-3.

[5] J. H. Evertse, On equations in S-units and the Thue–Mahler equation, Invent. Math. 75 (1984), 561–584, doi:10.1007/BF01388644.

[6] T. Hackman, I. Cycles through 1 for the $qn + 1$ map: no cycles of length 3, and constraints on longer cycles, companion paper.

[7] P. Mihăilescu, Primary cyclotomic units and a proof of Catalan's conjecture, J. Reine Angew. Math. 572 (2004), 167–195, doi:10.1515/crll.2004.048.

[8] D. Ridout, The p-adic generalization of the Thue–Siegel–Roth theorem, Mathematika 5 (1958), 40–48, doi:10.1112/S0025579300001339.

[9] R. Santos, On the Collatz general problem $qn + 1$, preprint (2020), arXiv:2005.00346.