

Bairdii ND4

POPULATION SIZE, MIGRATION, DIVERGENCE, ASSIGNMENT, HISTORY


Bayesian inference using the structured coalescent

Migrate-n version 4.4.4(git:v4-series-26-ge85c6ff) [June-1-2019]

Compiled for a SYMMETRIC multiprocessors (Grandcentral)

Program started at Thu May 20 09:28:39 2021

Program finished at Thu May 20 18:12:25 2021 [Runtime:0000:08:43:46]



Options

Inheritance scalers in use for Thetas:

All loci use an inheritance scaler of 1.0

[The locus with a scaler of 1.0 used as reference]

Random number seed: (with internal timer) 3305160015

Start parameters:

Theta values were generated RANDOM start value from the prior

M values were generated RANDOM start value from the prior

Connection matrix:

m = average (average over a group of Thetas or M,

s = symmetric migration M, S = symmetric 4Nm,

0 = zero, and not estimated,

* = migration free to vary, Thetas are on diagonal

d = row population split off column population, D = split and then migration

Population	1	2
1 GSL	*	d
2 SEV	0	*

Order of parameters:

1	Θ_1	<displayed>
2	Θ_2	<displayed>
3	$\Delta_{2 \rightarrow 1}$	<displayed>

4 $\sigma_{2 \rightarrow 1}$ <displayed>

Mutation rate among loci: Mutation rate is constant

Analysis strategy: Bayesian inference

-Population size estimation: Exponential Distribution

-Geneflow estimation: Exponential Distribution

-Divergence time estimation: Normal Distribution Shortcut (mean and standard dev.)

Proposal distributions for parameter

Parameter	Proposal
Theta	Metropolis sampling
M	Metropolis sampling
Divergence	Metropolis sampling
Divergence Spread	Metropolis sampling
Genealogy	Metropolis-Hastings

Prior distribution for parameter

Parameter	Prior	Minimum	Mean	Maximum	Delta	Bins	UpdateFreq
1 Theta **	Uniform	0.000000	0.050	0.100	0.010	1500	0.12500
2 Theta **	Uniform	0.000000	0.050	0.100	0.010	1500	0.12500
3 Splittime mean 10	Uniform	0.000000	0.010	100.0	10.00	1500	0.12500
4 Splittime std 10	Uniform	0.000000	0.010	100.0	10.00	1500	0.12500

[* * means priors were set globally]

Markov chain settings: Long chain

Number of chains	1
Recorded steps [a]	10000
Increment (record every x step [b])	1000
Number of concurrent chains (replicates) [c]	2
Visited (sampled) parameter values [a*b*c]	20000000
Number of discard trees per chain (burn-in)	1000

Multiple Markov chains:

Static heating scheme	4 chains with temperatures
1000000.00	3.00 1.50 1.00
	Swapping interval is 1

Print options:

Data file:	infile
Haplotyping is turned on:	NO
Output file:	outfile
Log file:	logfile

Posterior distribution raw histogram file:	bayesfile
Raw data from the MCMC run:	bayesallfile.gz
Print data:	No
Print genealogies [only some for some data type]:	None

Data summary

Data file: infile
 Datatype: Haplotype data
 Number of loci: 1

Mutationmodel:

Locus	Sublocus	Mutationmodel	Mutationmodel parameters
1	1	HKY	[Bf:0.26 0.33 0.13 0.29, kappa=4.630]

Sites per locus

Locus	Sites
1	363

Site rate variation and probabilities:

Locus	Sublocus	Region type	Rate of change	Probability	Patch size
-------	----------	-------------	----------------	-------------	------------

1	1	1	1.000	1.000	1.000	
Population					Locus	Gene copies data (missing)
1 GSL					1	203
2 SEV					1	141
Total of all populations					1	344 (0)





Bayesian Analysis: Posterior distribution table

Locus	Parameter	2.5%	25.0%	Mode	75.0%	97.5%	Median	Mean
1	Θ_1	0.02633	0.03760	0.04150	0.04527	0.05667	0.04157	0.04206
1	Θ_2	0.01540	0.02387	0.02917	0.03420	0.04280	0.02923	0.02922
1	$D_{2 \rightarrow 1}$	0.00000	0.00000	0.03333	12.46667	45.20000	12.50000	31.22200
1	$S_{2 \rightarrow 1}$	1.80000	18.93333	37.16667	41.26667	46.20000	26.56667	53.03280

Citation suggestions:

Beerli P., 2006. Comparison of Bayesian and maximum-likelihood inference of population genetic parameters. *Bioinformatics* 22:341-345

Beerli P., 2009. How to use MIGRATE or why are Markov chain Monte Carlo programs difficult to use?
In *Population Genetics for Animal Conservation*, G. Bertorelle, M. W. Bruford, H. C. Hauffe, A. Rizzoli, and C. Vernesi, eds., vol. 17 of *Conservation Biology*, Cambridge University Press, Cambridge UK, pp. 42-79.

Bayesian Analysis: Posterior distribution for locus 1

Log-Probability of the data given the model (marginal likelihood)

Use this value for Bayes factor calculations:

$BF = \text{Exp}[\ln(\text{Prob}(D \mid \text{thisModel}) - \ln(\text{Prob}(D \mid \text{otherModel}))]$

or as $LBF = 2 (\ln(\text{Prob}(D \mid \text{thisModel}) - \ln(\text{Prob}(D \mid \text{otherModel})))$

shows the support for thisModel]

Method	$\ln(\text{Prob}(D \mid \text{Model}))$	Notes
Thermodynamic integration	-1733.634962	(1a)
	-1254.295073	(1b)
Harmonic mean	-935.469983	(2)

(1a, 1b and 2) are approximations to the marginal likelihood, make sure that the program run long enough!

(1a, 1b) and (2) should give similar results, in principle.

But (2) is overestimating the likelihood, it is presented for historical reasons and should not be used

(1a, 1b) needs heating with chains that span a temperature range of 1.0 to at least 100,000.

(1b) is using a Bezier-curve to get better approximations for runs with low number of heated chains

Citation suggestions:

Beerli P. and M. Palczewski, 2010. Unified framework to evaluate panmixia and migration direction among multiple sampling locations, *Genetics*, 185: 313-326.

Acceptance ratios for all parameters and the genealogies

Parameter	Accepted changes	Ratio
Θ_1	1597549/2500727	0.63883
Θ_2	912497/2500594	0.36491
$\Delta_{2 \rightarrow 1}$	1821620/2498656	0.72904
$\sigma_{2 \rightarrow 1}$	1961660/2502541	0.78387
Genealogies	2601467/9997482	0.26021

MCMC-Autocorrelation and Effective MCMC Sample Size

Parameter	Autocorrelation	Effective Sample Size
Θ_1	0.80371	2433.85
Θ_2	0.59694	5048.59
$\Delta_{2 \rightarrow 1}$	-0.00266	20104.80
$\sigma_{2 \rightarrow 1}$	0.00931	19634.45
Genealogies	0.00931	19634.45

Potential Problems

This section reports potential problems with your run, but such reporting is often not very accurate. With many parameters in a multilocus analysis, it is very common that some parameters for some loci will not be very informative, triggering suggestions (for example to increase the prior range) that are not sensible. This suggestion tool will improve with time, therefore do not blindly follow its suggestions. If some parameters are flagged, inspect the tables carefully and judge whether an action is required. For example, if you run a Bayesian inference with sequence data, for macroscopic species there is rarely the need to increase the prior for Theta beyond 0.1; but if you use microsatellites it is rather common that your prior distribution for Theta should have a range from 0.0 to 100 or more. With many populations (>3) it is also very common that some migration routes are estimated poorly because the data contains little or no information for that route. Increasing the range will not help in such situations, reducing number of parameters may help in such situations.

No warning was recorded during the run