


Combinatorics in Kazhdan–Lusztig polynomials of paving matroids

by Trevor K. Karn (U. Minnesota)
(joint with George Nasr, Nick Proudfoot, and Lorenzo Vecchi)
on Saturday, March 18, 2023

A classical story

Our story




Euclid stated five postulates for rigorous geometry.

We can drop the fifth and still do geometry.

A partition $\lambda \vdash n$ is a weakly decreasing sequence of nonnegative integers $\lambda_1 \geq \lambda_2 \geq \cdots$ summing to n .

Example

$\lambda = (5, 4, 4, 2, 1) \vdash 16$ has Ferrers diagram



A Young tableau T is a filling of a Ferrers diagram by positive integers. T is standard if it is filled by $\{1, 2, \dots, n\}$ and increasing in rows and columns. Define f^λ as the number of standard tableaux of shape λ .

Example

One of $f^{(5,4,4,2,1)} = 549120$ standard Young tableaux:

| | | | | |
|---|---|----|----|----|
| 1 | 6 | 10 | 13 | 16 |
| 2 | 7 | 11 | 14 | |
| 3 | 8 | 12 | 15 | |
| 4 | 9 | | | |
| 5 | | | | |

Fact

Fix n . Then

$$\sum_{\lambda \vdash n} (f^\lambda)^2 = n!$$

Fact

Fix n . Then

$$\sum_{\lambda \vdash n} (f^\lambda)^2 = n!$$

Proof 1:

The Robinson-Schensted bijection:

pairs of standard tableaux of same shape \longleftrightarrow symmetric group \mathfrak{S}_n

Fact

Fix n . Then

$$\sum_{\lambda \vdash n} (f^\lambda)^2 = n!$$


Proof 1:

The Robinson-Schensted bijection:

pairs of standard tableaux of same shape \longleftrightarrow symmetric group \mathfrak{S}_n

Proof 2:

Uses that irreducible representations of \mathfrak{S}_n are indexed by $\lambda \vdash n$
and have dimension f^λ




Let v_1, \dots, v_n be vectors in a vector space V . Then any two bases A, B for the span of v_1, \dots, v_n satisfy the following requirements:

- 1) There must be at least one basis
- 2) If $a \in A - B$ then there is a $b \in B$ with $(A - a) \cup b$ a basis.




Let v_1, \dots, v_n be vectors in a vector space V . Then any two bases A, B for the span of v_1, \dots, v_n satisfy the following requirements:

- 1) There must be at least one basis
- 2) If $a \in A - B$ then there is a $b \in B$ with $(A - a) \cup b$ a basis.



Let v_1, \dots, v_n be vectors in a vector space V . Then any two bases A, B for the span of v_1, \dots, v_n satisfy the following requirements:

- 1) There must be at least one basis
- 2) If $a \in A - B$ then there is a $b \in B$ with $(A - a) \cup b$ a basis.



Definition

A **matroid** $M = (E, \mathcal{B})$ is a set E with $\emptyset \neq \mathcal{B} \subseteq 2^E$ such that if $A, B \in \mathcal{B}$ and $a \in A$, there exists $b \in B$ such that

$$(A - a) \cup b \in \mathcal{B}$$

Call \mathcal{B} the bases of the matroid.

Definition

A **matroid** $M = (E, \mathcal{B})$ is a set E with $\emptyset \neq \mathcal{B} \subseteq 2^E$ such that if $A, B \in \mathcal{B}$ and $a \in A$, there exists $b \in B$ such that

$$(A - a) \cup b \in \mathcal{B}$$

Call \mathcal{B} the bases of the matroid.

Matroids have the combinatorics of vectors without the “zeroth postulate”.

» Vocab

| Finite set of vectors | Matroids |
|----------------------------|---------------------------|
| Maximally independent sets | Bases \mathcal{B} |
| Minimally dependent sets | Circuits \mathcal{C} |
| Dimension of span | Rank |
| Codimension 1 subspaces | Hyperplanes \mathcal{H} |

» Vocab

| Finite set of vectors | Matroids |
|----------------------------|---------------------------|
| Maximally independent sets | Bases \mathcal{B} |
| Minimally dependent sets | Circuits \mathcal{C} |
| Dimension of span | Rank |
| Codimension 1 subspaces | Hyperplanes \mathcal{H} |

Definition

$\mathcal{CH} = \mathcal{C} \cap \mathcal{H}$ is the set of circuit hyperplanes.

A hyperplane H is stressed if every subset of H of size $\text{rk}(E)$ is in \mathcal{C} . Denote the set of (nontrivial) stressed hyperplanes by \mathcal{SH} .

Fact

There is a very large class of matroids called (sparse) paving matroids.

Fact

There is a very large class of matroids called (sparse) paving matroids.


Fact

The matroid Kazhdan–Lusztig polynomial $P_M(t)$ is an interesting polynomial invariant of a matroid M , introduced by Elias, Proudfoot, and Wakefield in 2016.

A skew partition λ/μ is a pair of partitions where the diagram of μ is contained in the diagram of λ

Example



If $\lambda = (4, 3, 2, 1)$ and $\mu = (2, 1)$ then λ/μ has diagram



A skew tableau T is a filling of a skew diagram by positive integers. T is standard if it is filled by $\{1, 2, \dots, |\lambda| - |\mu|\}$ and increasing in rows and columns. Define $f^{\lambda/\mu}$ as the number of standard skew tableaux of shape λ/μ .

Example

Two of $f^{(4,3,2,1)/(2,1)} = 272$ standard skew tableaux:



Theorem (Lee, Nasr, Radcliffe '21)



Let M be a rank- k , sparse paving matroid with $E = [n]$ and circuit hyperplanes \mathcal{CH} . The t^i coefficient in $P_M(t)$ is

$$f^{\lambda/\mu} - |\mathcal{CH}| f^{\lambda'/\mu'}$$

where

$$\lambda = [n - 2i, (k - 2i + 1)^i], \mu = [(k - 2i - 1)^i]$$

$$\lambda' = [(k - 2i + 1)^{i+1}], \mu' = [k - 2i, (k - 2i - 1)^{i-1}]$$



Theorem (Lee, Nasr, Radcliffe '21)

For a sparse paving matroid M , the t^i coefficient in $P_M(t)$ is

$$f^{\lambda/\mu} - |\mathcal{CH}| f^{\lambda'/\mu'}$$

Proof 1 (LNR '21): Combinatorial argument with recursion.

Fact

There is a (reducible) \mathfrak{S}_n representation $S^{\lambda/\mu}$ of dimension $f^{\lambda/\mu}$.

Theorem (Lee, Nasr, Radcliffe '21)


For a sparse paving matroid M , the t^i coefficient in $P_M(t)$ is

$$f^{\lambda/\mu} - |\mathcal{CH}| f^{\lambda'/\mu'}$$


Proof 1 (LNR '21): Combinatorial argument with recursion.

Proof 2 (KNPV '22): $\dim(\text{some } S^{\lambda/\mu} \text{ coming from } M)$.

» Example: Vámos matroid



$$\lambda = [6, 3], \mu = [1] \longrightarrow$$


$$\lambda' = [3, 3], \mu' = [2] \longrightarrow$$


$$|\mathcal{CH}| = 5$$

$$f^{\lambda/\mu} - 5f^{\lambda'/\mu'} = 48 - 15 = 33$$

$$P_V(t) = 1 + 33t$$

» Example: Projective plane over \mathbb{F}_3

$$\lambda/\mu = \begin{array}{|c|c|c|c|c|c|c|c|c|c|c|} \hline & & & & & & & & & & \\ \hline & & & & & & & & & & \\ \hline \end{array}$$

$$\mathcal{CH} = \emptyset$$

$$f^{\lambda/\mu} = 65 \neq 0$$

$$P_M(t) = 1$$

» Example: Projective plane over \mathbb{F}_3

$$\lambda/\mu =$$


$$|\mathcal{SH}| = 13$$

$$\lambda'/\mu' =$$


$$f^{\lambda/\mu} - 13f^{\lambda'/\mu'} = 65 - 13 * 5 = 0$$

$$P_M(t) = 1$$

Theorem

For a (arbitrary!) paving matroid M , the t^i coefficient in $P_M(t)$ is



$$f^{\lambda/\mu} = \sum_{H \in \mathcal{SH}} f^{\lambda'(|H|)/\mu'}$$

where


$$\lambda = [n - 2i, (k - 2i + 1)^i], \mu = [(k - 2i - 1)^i]$$

$$\lambda'(h) = [h - 2i + 1, (k - 2i + 1)^i], \mu' = [h - 2i, (k - 2i - 1)^{i-1}]$$

Proof: Our proof of LNR's theorem implies this more general result



Certain $P_M(t)$ in terms of standard skew Young tableaux



Combinatorial proof [LNR21]

Algebraic proof [KNPV22]

THANK YOU!

» References



Ben Elias, Nicholas Proudfoot, and Max Wakefield, The Kazhdan–Lusztig polynomial of a matroid, Advances in Mathematics **299** (2016), 36–70.



Luisa Ferroni, George D. Nasr, and Lorenzo Vecchi, Stressed hyperplanes and Kazhdan–Lusztig gamma-positivity for matroids, 2021.



Katie R. Gedeon, Nicholas Proudfoot, and Benjamin Young, The equivariant Kazhdan–Lusztig polynomial of a matroid, J. Comb. Theory, Ser. A **150** (2017), 267–294.



Alice L. L. Gao and Matthew H. Y. Xie, The inverse Kazhdan–Lusztig polynomial of a matroid, J. Combin. Theory Ser. B **151** (2021), 375–392. MR 4294228

» References (cont.)



Alice L. L. Gao, Matthew H.Y. Xie, and Arthur L. B. Yang, The equivariant inverse Kazhdan–Lusztig polynomials of uniform matroids, 2021.



Trevor Karn, George Nasr, Nicholas Proudfoot, and Lorenzo Vecchi, Equivariant Kazhdan–Lusztig theory of paving matroids, 2022.



Kyungyong Lee, George D. Nasr, and Jamie Radcliffe, A combinatorial formula for Kazhdan–Lusztig polynomials of sparse paving matroids, *Electron. J. Comb.* **28** (2021).



Dillon Mayhew, Mike Newman, Dominic Welsh, and Geoff Whittle, On the asymptotic proportion of connected matroids, *European J. Combin.* **32** (2011), no. 6, 882–890. MR 2821559

» References (cont.)



Peter Orlik and Louis Solomon, Combinatorics and topology of complements of hyperplanes, *Inventiones mathematicae* **56** (1980), no. 2, 167–189 (eng).



Rudi Pendavingh and Jorn van der Pol, On the number of matroids compared to the number of sparse paving matroids, *Electron. J. Combin.* **22** (2015), no. 2, Paper 2.51, 17. MR 3367294



Nicholas Proudfoot, Yuan Xu, and Benjamin Young, The Z-polynomial of a matroid, *Electron. J. Comb.* **25** (2018), P1.26.