

Combinatorics in matroid Kazhdan–Lusztig polynomials

by Trevor K. Karn (U. Minnesota - Twin Cities)
(joint with George Nasr, Nick Proudfoot, and Lorenzo Vecchi)
on Thursday, March 23, 2023

A classical story

Matroids


Combinatorial formulas



A partition $\lambda \vdash n$ is a weakly decreasing sequence of nonnegative integers $\lambda_1 \geq \lambda_2 \geq \cdots$ summing to n .

Example

$\lambda = (5, 4, 4, 2, 1) \vdash 16$ has Ferrers diagram



A Young tableau T is a filling of a Ferrers diagram by positive integers. T is standard if it is filled by $\{1, 2, \dots, n\}$ and increasing in rows and columns. Define f^λ as the number of standard tableaux of shape λ .

Example

One of $f^{(5,4,4,2,1)} = 549120$ standard Young tableaux:

1	6	10	13	16
2	7	11	14	
3	8	12	15	
4	9			
5				

Fact

Fix n . Then

$$\sum_{\lambda \vdash n} (f^\lambda)^2 = n!$$

Fact

Fix n . Then

$$\sum_{\lambda \vdash n} (f^\lambda)^2 = n!$$

Proof 1:

The Robinson-Schensted bijection:

pairs of standard tableaux of same shape \longleftrightarrow permutation in \mathfrak{S}_n

Definition

A representation of a group G is a homomorphism

$$\rho : G \rightarrow \mathrm{GL}_n(\mathbb{C}).$$

A representation is irreducible if there is no G -stable subspace $W \subseteq \mathbb{C}^n$.

Example

There is a representation of \mathfrak{S}_3 defined by sending

$$(12) \mapsto \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$(23) \mapsto \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}.$$

The action: $\pi \in \mathfrak{S}_3$ permutes coordinates. So

$$(12) \cdot \langle 1, 2, 3 \rangle = \langle 2, 1, 3 \rangle,$$

and

$$\pi \cdot \langle 1, 1, 1 \rangle = \langle 1, 1, 1 \rangle.$$

Fact

Irreducible \mathfrak{S}_n representations are indexed by $\lambda \vdash n$.
Denote them by S^λ . Then

$$\dim S^\lambda = f^\lambda.$$

Fact

Irreducible \mathfrak{S}_n representations are indexed by $\lambda \vdash n$.
Denote them by S^λ . Then

$$\dim S^\lambda = f^\lambda.$$

Example

The representation on the last slide contains $S^{(3)}$ and $S^{(2,1)}$.

Fact

Irreducible \mathfrak{S}_n representations are indexed by $\lambda \vdash n$.
Denote them by S^λ . Then

$$\dim S^\lambda = f^\lambda.$$

Example

The representation on the last slide contains $S^{(3)}$ and $S^{(2,1)}$.

Fact

Let d_1, d_2, \dots, d_r be the dimensions of all irreducible representations of a finite group. Then

$$\sum_i d_i^2 = |G|.$$


Fact

Fix n . Then

$$\sum_{\lambda \vdash n} (f^\lambda)^2 = n!$$

Proof 2:

$$\sum_{\lambda} (f^\lambda)^2 = \sum_i d_i^2 = |G| = |\mathfrak{S}_n| = n!$$




Let v_1, \dots, v_n be vectors in a vector space V (not all 0). Then any two bases A, B for the span of v_1, \dots, v_n satisfy the following requirements:

- 1) There must be at least one basis
- 2) If $a \in A - B$ then there is a $b \in B$ with $(A - a) \cup b$ a basis.




Let v_1, \dots, v_n be vectors in a vector space V (not all 0). Then any two bases A, B for the span of v_1, \dots, v_n satisfy the following requirements:

- 1) There must be at least one basis
- 2) If $a \in A - B$ then there is a $b \in B$ with $(A - a) \cup b$ a basis.



Let v_1, \dots, v_n be vectors in a vector space V (not all 0). Then any two bases A, B for the span of v_1, \dots, v_n satisfy the following requirements:

- 1) There must be at least one basis
- 2) If $a \in A - B$ then there is a $b \in B$ with $(A - a) \cup b$ a basis.



Definition 1

A **matroid** $M = (E, \mathcal{B})$ is a finite set E with $\emptyset \neq \mathcal{B} \subseteq 2^E$ such that if $A, B \in \mathcal{B}$ and $a \in A$, there exists $b \in B$ such that

$$(A - a) \cup b \in \mathcal{B}$$

Call \mathcal{B} the bases of the matroid.

Definition 1

A **matroid** $M = (E, \mathcal{B})$ is a finite set E with $\emptyset \neq \mathcal{B} \subseteq 2^E$ such that if $A, B \in \mathcal{B}$ and $a \in A$, there exists $b \in B$ such that

$$(A - a) \cup b \in \mathcal{B}$$

Call \mathcal{B} the bases of the matroid.

Matroids have the combinatorics of vectors without the “zeroth postulate”.

Definition 2

A **matroid** $M = (E, \mathcal{C})$ is a set E with $\mathcal{C} \subseteq 2^E$ such that

$$\emptyset \notin \mathcal{C}$$

If $C_1, C_2 \in \mathcal{C}$ with $C_1 \subseteq C_2$, then $C_1 = C_2$.

If $C_1, C_2 \in \mathcal{C}$ are distinct, and $e \in C_1 \cap C_2$, then there is a $C_3 \in \mathcal{C}$ such that $C_3 \subseteq (C_1 \cup C_2) - e$

Call \mathcal{C} the circuits.

Example

The **uniform matroid** $U_{k,n}$ models n -many
 k -dimensional vectors in general position

Bases \longleftrightarrow any set of k -many vectors

Circuits \longleftrightarrow any set of $k + 1$ -many vectors

Example

The **uniform matroid** $U_{k,n}$ models n -many k -dimensional vectors in general position

Bases \longleftrightarrow any set of k -many vectors

Circuits \longleftrightarrow any set of $k + 1$ -many vectors

Example of the example

$U_{3,12}$ corresponds to 12 generic vectors in \mathbb{R}^3 . One choice of basis is $\{e_1, e_2, e_3\}$. On the other hand $\{e_1, e_2, e_3, v\}$ is dependent for any $v \in \mathbb{R}^3$.

Example

A graph Γ with edges E forms a matroid:

Bases \longleftrightarrow spanning trees

Circuits \longleftrightarrow cycles



Example

The columns of a matrix form a matroid:

$$\begin{bmatrix} 1 & 2 & 0 & 0 \\ 2 & 4 & 2 & 4 \end{bmatrix}$$



Example

The projective plane is the set of lines through the origin in \mathbb{F}^3 .

Example

The projective plane is the set of lines through the origin in \mathbb{F}^3 .


If $\mathbb{F} = \mathbb{F}_2$,

$$\begin{bmatrix} 1 & 0 & 0 & 1 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 1 & 1 \end{bmatrix}$$

Theorem (Nelson, 2018)

Almost all matroids cannot be written as a matrix.

» Example: Vámos matroid



» Vocab

Finite set of vectors	Matroids
Maximally independent sets	Bases \mathcal{B}
Minimally dependent sets	Circuits \mathcal{C}
Dimension of span	Rank
Subspaces	Flats \mathcal{F}
Codimension 1 subspaces	Hyperplanes \mathcal{H}

» Vocab

Finite set of vectors	Matroids
Maximally independent sets	Bases \mathcal{B}
Minimally dependent sets	Circuits \mathcal{C}
Dimension of span	Rank
Subspaces	Flats \mathcal{F}
Codimension 1 subspaces	Hyperplanes \mathcal{H}

Definition

$\mathcal{CH} = \mathcal{C} \cap \mathcal{H}$ is the set of circuit hyperplanes.

A hyperplane H is stressed if every subset of H of size $\text{rk}(E)$ is in \mathcal{C} . Denote the set of (nontrivial) stressed hyperplanes by \mathcal{SH} .

M is a paving matroid if all circuits have size at least $k = \text{rk}(E)$

A paving matroid is sparse if $\binom{E}{k} = \mathcal{CH} \sqcup \mathcal{B}$

M is a paving matroid if all circuits have size at least $k = \text{rk}(E)$

A paving matroid is sparse if $\binom{E}{k} = \mathcal{CH} \sqcup \mathcal{B}$

Conjecture (Mayhew, Newman, Welsh, Whittle '11)

Asymptotically almost all matroids are sparse paving

Theorem (Pendavingh, van der Pol '15)

Asymptotically logarithmically almost all matroids are sparse paving

M is a paving matroid if all circuits have size at least $k = \text{rk}(E)$

A paving matroid is sparse if $\binom{E}{k} = \mathcal{CH} \sqcup \mathcal{B}$

Example

$U_{k,n} \leftrightarrow$ sparse paving

M is a paving matroid if all circuits have size at least $k = \text{rk}(E)$

A paving matroid is sparse if $\binom{E}{k} = \mathcal{CH} \sqcup \mathcal{B}$

Example

$U_{k,n} \leftrightarrow$ sparse paving

$\begin{bmatrix} 1 & 2 & 0 & 0 \\ 2 & 4 & 2 & 4 \end{bmatrix} \leftrightarrow$ paving

M is a paving matroid if all circuits have size at least $k = \text{rk}(E)$

A paving matroid is sparse if $\binom{E}{k} = \mathcal{CH} \sqcup \mathcal{B}$

Example

$U_{k,n} \leftrightarrow$ sparse paving

$\begin{bmatrix} 1 & 2 & 0 & 0 \\ 2 & 4 & 2 & 4 \end{bmatrix} \leftrightarrow$ paving

$\mathbb{P}\mathbb{F}_3 \leftrightarrow$ paving

M is a paving matroid if all circuits have size at least $k = \text{rk}(E)$

A paving matroid is sparse if $\binom{E}{k} = \mathcal{CH} \sqcup \mathcal{B}$

Example

$U_{k,n} \leftrightarrow$ sparse paving

$\begin{bmatrix} 1 & 2 & 0 & 0 \\ 2 & 4 & 2 & 4 \end{bmatrix} \leftrightarrow$ paving

$\mathbb{PF}_3 \leftrightarrow$ paving

$V \leftrightarrow$ sparse paving

Fact


The matroid Kazhdan–Lusztig polynomial $P_M(t)$ is an interesting polynomial invariant of a matroid M , introduced by Elias, Proudfoot, and Wakefield in 2016.

It is defined in terms of \mathcal{F} .

A skew partition λ/μ is a pair of partitions where the diagram of μ is contained in the diagram of λ

Example



If $\lambda = (4, 3, 2, 1)$ and $\mu = (2, 1)$ then λ/μ has diagram



A skew tableau T is a filling of a skew diagram by positive integers. T is standard if it is filled by $\{1, 2, \dots, |\lambda| - |\mu|\}$ and increasing in rows and columns. Define $f^{\lambda/\mu}$ as the number of standard skew tableaux of shape λ/μ .

Example

Two of $f^{(4,3,2,1)/(2,1)} = 272$ standard skew tableaux:



Theorem (Lee, Nasr, Radcliffe '21)



Let M be a rank- k , sparse paving matroid with $E = [n]$ and circuit hyperplanes \mathcal{CH} . The t^i coefficient in $P_M(t)$ is

$$f^{\lambda/\mu} - |\mathcal{CH}| f^{\lambda'/\mu'}$$

where

$$\lambda = [n - 2i, (k - 2i + 1)^i], \mu = [(k - 2i - 1)^i]$$

$$\lambda' = [(k - 2i + 1)^{i+1}], \mu' = [k - 2i, (k - 2i - 1)^{i-1}]$$



Theorem (Lee, Nasr, Radcliffe '21)

For a sparse paving matroid M , the t^i coefficient in $P_M(t)$ is

$$f^{\lambda/\mu} - |\mathcal{CH}| f^{\lambda'/\mu'}$$

Proof 1 (LNR '21): Combinatorial argument with recursion.

Fact

There is a (reducible) \mathfrak{S}_n representation $S^{\lambda/\mu}$ of dimension $f^{\lambda/\mu}$.

Theorem (Lee, Nasr, Radcliffe '21)


For a sparse paving matroid M , the t^i coefficient in $P_M(t)$ is

$$f^{\lambda/\mu} - |\mathcal{CH}| f^{\lambda'/\mu'}$$

Proof 1 (LNR '21): Combinatorial argument with recursion.

Proof 2 (KNPV '22): $\dim(\text{some } S^{\lambda/\mu} \text{ coming from } M)$.

» Example: Vámos matroid



Some general facts:

Know $P_M(t)$ always has constant term 1.

Know $\deg P_M(t) < \frac{\text{rk } E}{2}$.

Some general facts:

Know $P_M(t)$ always has constant term 1.

Know $\deg P_M(t) < \frac{\text{rk } E}{2}$.

$\text{rk } V = 4$ so $P_V(t) = 1 + ?t$.

Some general facts:


Know $P_M(t)$ always has constant term 1.

Know $\deg P_M(t) < \frac{\text{rk } E}{2}$.

$\text{rk } V = 4$ so $P_V(t) = 1 + ?t$.

Only need to compute linear coefficient!

$$\lambda = [6, 3], \mu = [1] \longrightarrow$$


$$\lambda' = [3, 3], \mu' = [2] \longrightarrow$$


$$|\mathcal{CH}| = 5$$

$$f^{\lambda/\mu} - 5f^{\lambda'/\mu'} = 48 - 15 = 33$$

$$P_V(t) = 1 + 33t$$

» Example: Projective plane over \mathbb{F}_3



$$\mathcal{CH} = \emptyset$$

$$f^{\lambda/\mu} = 65 \neq 0$$

From Elias, Proudfoot, and Wakefield, we know

$$P_M(t) = 1$$

» Example: Projective plane over \mathbb{F}_3

$$\lambda/\mu =$$


$$|\mathcal{SH}| = 13$$

$$\lambda'/\mu' =$$


$$f^{\lambda/\mu} - 13f^{\lambda'/\mu'} = 65 - 13 * 5 = 0$$

From Elias, Proudfoot, and Wakefield, we know

$$P_M(t) = 1$$

Theorem

For a (arbitrary!) paving matroid M , the t^i coefficient in $P_M(t)$ is



$$f^{\lambda/\mu} = \sum_{H \in \mathcal{SH}} f^{\lambda'(|H|)/\mu'}$$

where

$$\lambda = [n - 2i, (k - 2i + 1)^i], \mu = [(k - 2i - 1)^i]$$

$$\lambda'(h) = [h - 2i + 1, (k - 2i + 1)^i], \mu' = [h - 2i, (k - 2i - 1)^{i-1}]$$

Proof: Our proof of LNR's theorem implies this more general result



» Proof idea

In a stressed hyperplane, all size- k subsets are circuits.
Create a new matroid by turning all circuits in H into bases

» Proof idea

In a stressed hyperplane, all size- k subsets are circuits.
Create a new matroid by turning all circuits in H into bases



turns into




» Proof idea

Do this until you obtain $U_{k,n}$


» Proof idea

Do this until you obtain $U_{k,n}$



» Proof idea

Do this until you obtain $U_{k,n}$



Each step accounts for $S^{\lambda'(h)/\mu'}$.

Certain $P_M(t)$ in terms of standard skew Young tableaux

Combinatorial proof [LNR21]

Algebraic proof [KNPV22]
+ extension

THANK YOU!

» References



Ben Elias, Nicholas Proudfoot, and Max Wakefield, The Kazhdan–Lusztig polynomial of a matroid, Advances in Mathematics **299** (2016), 36–70.



Luisa Ferroni, George D. Nasr, and Lorenzo Vecchi, Stressed hyperplanes and Kazhdan–Lusztig gamma-positivity for matroids, 2021.



Katie R. Gedeon, Nicholas Proudfoot, and Benjamin Young, The equivariant Kazhdan–Lusztig polynomial of a matroid, J. Comb. Theory, Ser. A **150** (2017), 267–294.



Alice L. L. Gao and Matthew H. Y. Xie, The inverse Kazhdan–Lusztig polynomial of a matroid, J. Combin. Theory Ser. B **151** (2021), 375–392. MR 4294228

» References (cont.)



Alice L. L. Gao, Matthew H.Y. Xie, and Arthur L. B. Yang, The equivariant inverse Kazhdan–Lusztig polynomials of uniform matroids, 2021.



Trevor Karn, George Nasr, Nicholas Proudfoot, and Lorenzo Vecchi, Equivariant Kazhdan–Lusztig theory of paving matroids, 2022.



Kyungyong Lee, George D. Nasr, and Jamie Radcliffe, A combinatorial formula for Kazhdan–Lusztig polynomials of sparse paving matroids, Electron. J. Comb. **28** (2021).



Dillon Mayhew, Mike Newman, Dominic Welsh, and Geoff Whittle, On the asymptotic proportion of connected matroids, European J. Combin. **32** (2011), no. 6, 882–890. MR 2821559

» References (cont.)



Peter Orlik and Louis Solomon, Combinatorics and topology of complements of hyperplanes, *Inventiones mathematicae* **56** (1980), no. 2, 167–189 (eng).



Rudi Pendavingh and Jorn van der Pol, On the number of matroids compared to the number of sparse paving matroids, *Electron. J. Combin.* **22** (2015), no. 2, Paper 2.51, 17. MR 3367294



Nicholas Proudfoot, Yuan Xu, and Benjamin Young, The Z-polynomial of a matroid, *Electron. J. Comb.* **25** (2018), P1.26.