

GALERKIN APPROXIMATIONS OF DELAY DIFFERENTIAL EQUATIONS

HONGHU LIU AND TREVOR NORTON

Abstract.

1. Introduction.

2. Preliminaries.

2.1. The type of DDE. We are interested in approximating the solution to the following DDE:

$$(2.1) \quad \begin{aligned} \frac{dx}{dt} &= ax(t) + bx(t - \tau) + F(x(t - \tau)), \quad t > 0 \\ x(t) &= \varphi(t), \quad t \in [-\tau, 0] \end{aligned}$$

for $\varphi \in L^2([-\tau, 0]; \mathbb{R})$ and where $F : \mathbb{R} \rightarrow \mathbb{R}$ is Lipschitz with constant L . It is appropriate to formulate this problem into the space $\mathcal{H} := L^2([-\tau, 0]; \mathbb{R}) \times \mathbb{R}$.

2.2. Properties and Basic Results of Koornwinder Polynomials. From [2, Eq. (2.1)], the sequence of Koornwinder polynomials $\{K_n\}$ can be built from the Legendre polynomials L_n by

$$(2.2) \quad K_n(s) := -(1+s) \frac{d}{ds} L_n(s) + (n^2 + n + 1) L_n(s), \quad s \in [-1, 1], \quad n \in \mathbb{N}_0.$$

Furthermore, we reproduce from [1, Prop. 3.1] some simple properties that $\{K_n\}$ satisfy.

PROPOSITION 2.1. *The polynomial K_n defined in (2.2) is of degree n and admits the following expansion in terms of the Legendre polynomials:*

$$(2.3) \quad K_n(s) = - \sum_{j=0}^{n-1} (2j+1) L_j(s) + (n^2 + 1) L_n(s), \quad n \in \mathbb{N}_0;$$

and the following normalization property holds:

$$(2.4) \quad K_n(1) = 1, \quad n \in \mathbb{N}_0.$$

Moreover, the sequence given by

$$(2.5) \quad \{\mathcal{K}_n := (K_n, K_n(1)) : n \in \mathbb{N}_0\}$$

forms an orthogonal basis of the product space

$$(2.6) \quad \mathcal{E} := L^2([-1, 1]; \mathbb{R}) \times \mathbb{R},$$

where \mathcal{E} is endowed with the following inner product:

$$(2.7) \quad \langle (f, a), (g, b) \rangle_{\mathcal{E}} = \frac{1}{2} \int_{-1}^1 f(s)g(s) ds + ab, \quad (f, a), (g, b) \in \mathcal{E}.$$

Moreover $\left\{ \frac{\mathcal{K}_n}{\|\mathcal{K}_n\|_{\mathcal{E}}} \right\}$ forms a Hilbert basis of \mathcal{E} where the norm $\|\mathcal{K}_n\|_{\mathcal{E}}$ of \mathcal{K}_n induced by $\langle \cdot, \cdot \rangle_{\mathcal{E}}$ possesses the following analytic expression:

$$(2.8) \quad \|\mathcal{K}_n\|_{\mathcal{E}} = \sqrt{\frac{(n^2 + 1)((n + 1)^2 + 1)}{2n + 1}}, \quad n \in \mathbb{N}_0.$$

Suppose that Π_N is the N -dimensional standard projection into $\text{span}\{\mathcal{K}_n : n \leq N\} \subset \mathcal{E}$. It will be relevant to discuss when we have convergence of $[\Pi_N u]^D$ for $u \in \mathcal{E}$. In particular, we will focus on uniform convergence. We define for $f \in L^2([-1, 1], \mathbb{R})$ the following:

$$(2.9) \quad a_n(f) := \frac{2n + 1}{2} \int_{-1}^1 f(x) L_n(x) dx.$$

PROPOSITION 2.2. Let $f \in C^2([-1, 1]; \mathbb{R})$ and denote $\psi = (f, f(0)) \in \mathcal{E}$. Then the series

$$(2.10) \quad [\Pi_N \psi]^D = \sum_{n=0}^N \frac{\langle \psi, \mathcal{K}_n \rangle_{\mathcal{E}}}{\|\mathcal{K}_n\|_{\mathcal{E}}^2} \mathcal{K}_n$$

converges uniformly to f .

Proof. It is easy to show based on (2.3) we have for $\theta \in [-1, 1]$

$$(2.11) \quad \begin{aligned} |K_n(\theta)| &\leq (n^2 + 1)|L_n(\theta)| + \sum_{j=0}^{n-1} (2j + 1)|L_j(\theta)| \\ &\leq (n^2 + 1) + \sum_{j=0}^{n-1} (2j + 1) \\ &= 2n^2 + 1, \end{aligned}$$

i.e., $\|K_n\|_{\infty} \leq 2n^2 + 1$.

By the definition of $\langle \cdot, \cdot \rangle_{\mathcal{E}}$ and the Koornwinder polynomials, we have that for $n \in \mathbb{N}_0$

$$(2.12) \quad \begin{aligned} \langle \psi, \mathcal{K}_n \rangle_{\mathcal{E}} &= \frac{1}{2} \int_{-1}^1 f(x) K_n(x) dx + f(1) \\ &= \frac{1}{2} \left[- \int_{-1}^1 f(x)(1 + x) L'_n(x) dx + (n^2 + n + 1) \int_{-1}^1 f(x) L_n(x) dx \right] + f(1). \end{aligned}$$

If we use integration by parts, we find that

$$(2.13) \quad - \int_{-1}^1 f(x)(1 + x) L'_n(x) dx = -2f(1) + \int_{-1}^1 f'(x)(1 + x) L_n(x) dx + \int_{-1}^1 f(x) L_n(x) dx.$$

Applying (2.13) to (2.12) gives that

$$(2.14) \quad \begin{aligned} \langle \psi, \mathcal{K}_n \rangle_{\mathcal{E}} &= \frac{1}{2} \int_{-1}^1 f'(x)(1 + x) L_n(x) dx + \frac{n^2 + n + 2}{2} \int_{-1}^1 f(x) L_n(x) dx \\ &= \frac{1}{2} \int_{-1}^1 f'(x)(1 + x) L_n(x) dx + \frac{n^2 + n + 2}{2n + 1} a_n(f). \end{aligned}$$

We can also note that by applying the Hölder inequality we get

$$(2.15) \quad \left| \int_{-1}^1 f'(x)(1+x)L_n(x) dx \right| \leq \|f'\|_\infty \left(\int_{-1}^1 (1+x) dx \right)^{1/2} \|L_n\|_{L^2} \\ = \frac{4\|f'\|_\infty}{\sqrt{6n+3}}.$$

Furthermore, from [4, Thm. 2.1] we have

$$(2.16) \quad |a_n(f)| \leq \frac{V_1}{n - \frac{1}{2}} \sqrt{\frac{\pi}{2n}},$$

where $V_1 := \int_{-1}^1 \frac{f''(x)}{\sqrt{1-x^2}} dx < \infty$. Thus,

$$(2.17) \quad |\langle \psi, \mathcal{K}_n \rangle_{\mathcal{H}}| \leq \frac{2\|f'\|_\infty}{\sqrt{6n+3}} + V_1 \sqrt{2\pi} \frac{n^2 + n + 2}{\sqrt{n}(4n^2 + 1)},$$

and so

$$(2.18) \quad \frac{|\langle \psi, \mathcal{K}_n \rangle_{\mathcal{H}}|}{\|\mathcal{K}_n\|_{\mathcal{H}}^2} \|K_n\|_\infty \leq \left[\frac{2\|f'\|_\infty}{\sqrt{6n+3}} + V_1 \sqrt{2\pi} \frac{n^2 + n + 2}{\sqrt{n}(4n^2 + 1)} \right] \times \left[\frac{(2n+1)(2n^2+1)}{(n^2+1)((n+1)^2+1)} \right] \\ = O\left(\frac{1}{n^{3/2}}\right).$$

By the Weierstrass M-test, the series (2.10) converges uniformly.

Note also that (2.10) is simply the functional part of the Koornwinder expansion of ψ in \mathcal{H} . So the series converges in $L^2([-1, 1]; \mathbb{R})$ to $\psi^D = f$. Therefore, since the series converges uniformly, it must converge uniformly to f . \square

It will also be necessary to prove certain properties of the series of Koornwinder polynomials

$$(2.19) \quad S_N(x) := \sum_{n=0}^N \frac{K_n}{\|\mathcal{K}_n\|_{\mathcal{E}}^2}, \quad N \in \mathbb{N}_0, \quad x \in [-1, 1].$$

If we were to denote $\psi = (0, 1) \in L^2([-1, 1]) \times \mathbb{R}$, then S_N would simply be the functional part of $\Pi_N \psi$. The following lemma allows us to express S_N in terms of Legendre Polynomials.

LEMMA 2.1. *The functions S_N defined in (2.19) can be expressed as*

$$(2.20) \quad S_N(x) = \frac{1}{(N+1)^2 + 1} \sum_{n=0}^N (2n+1)L_n, \quad x \in [-1, 1].$$

Proof. Using (2.3), we can show that for $m \leq N \in \mathbb{N}_0$

$$(2.21) \quad \int_{-1}^1 S_N(x)L_m(x) dx = \sum_{n=0}^N \frac{1}{\|\mathcal{K}_n\|_{\mathcal{E}}^2} \int_{-1}^1 K_n(x)L_m(x) dx \\ = \|L_m\|_{L^2([-1, 1])}^2 \left[(m^2 + 1) \frac{1}{\|\mathcal{K}_m\|_{\mathcal{E}}^2} - (2m+1) \sum_{k=m+1}^N \frac{1}{\|\mathcal{K}_k\|_{\mathcal{E}}^2} \right],$$

and so

$$(2.22) \quad S_N(x) = \sum_{n=0}^N \left[\frac{n^2 + 1}{\|\mathcal{K}_n\|_{\mathcal{E}}^2} - (2n + 1) \sum_{m=n+1}^N \frac{1}{\|\mathcal{K}_m\|_{\mathcal{E}}^2} \right] L_n(x).$$

It is easy to show that

$$(2.23) \quad \begin{aligned} \sum_{n=0}^N \frac{1}{\|\mathcal{K}_n\|_{\mathcal{E}}^2} &= \sum_{n=0}^N \frac{2n + 1}{(n^2 + 1)((n + 1)^2 + 1)} \\ &= \sum_{n=0}^N \left[\frac{1}{n^2 + 1} - \frac{1}{(n + 1)^2 + 1} \right] \\ &= 1 - \frac{1}{(N + 1)^2 + 1} \end{aligned}$$

and

$$(2.24) \quad \begin{aligned} \sum_{m=n+1}^N \frac{1}{\|\mathcal{K}_m\|_{\mathcal{E}}^2} &= \sum_{m=0}^N \frac{1}{\|\mathcal{K}_m\|_{\mathcal{E}}^2} - \sum_{m=0}^n \frac{1}{\|\mathcal{K}_m\|_{\mathcal{E}}^2} \\ &= \frac{1}{(n + 1)^2 + 1} - \frac{1}{(N + 1)^2 + 1}. \end{aligned}$$

Applying (2.24) to (2.22) gives

$$(2.25) \quad \begin{aligned} S_N(x) &= \sum_{n=0}^N \left[\frac{n^2 + 1}{\|\mathcal{K}_n\|_{\mathcal{E}}^2} - (2n + 1) \left(\frac{1}{(n + 1)^2 + 1} - \frac{1}{(N + 1)^2 + 1} \right) \right] L_n(x) \\ &= \sum_{n=0}^N \left[\frac{2n + 1}{(n + 1)^2 + 1} - \frac{2n + 1}{(n + 1)^2 + 1} + \frac{2n + 1}{(N + 1)^2 + 1} \right] L_n(x) \quad \square \\ &= \sum_{n=0}^N \frac{2n + 1}{(N + 1)^2 + 1} L_n(x). \end{aligned}$$

Now that we have this expression, we can prove the properties of S_N that will be useful when showing the main result.

PROPOSITION 2.3. *For the functions S_N defined in (2.19), we have that*

$$(2.26) \quad |S_N(x)| < 1, \quad \forall N \in \mathbb{N}_0, \quad \forall x \in [-1, 1].$$

Furthermore,

$$(2.27) \quad \lim_{N \rightarrow \infty} S_N(x) = 0, \quad \forall x \in (-1, 1).$$

Proof. It is known that

$$(2.28) \quad |L_n(x)| \leq 1, \quad \forall x \in [-1, 1], \quad \forall n \in \mathbb{N}_0.$$

Thus for $x \in [-1, 1]$ and $N \in \mathbb{N}_0$

$$\begin{aligned}
(2.29) \quad |S_N(x)| &\leq \frac{1}{(N+1)^2+1} \sum_{n=0}^N (2n+1) |L_n(x)| \\
&\leq \frac{1}{(N+1)^2+1} \sum_{n=0}^N (2n+1) \\
&= \frac{N^2+1}{(N+1)^2+1} \\
&< 1.
\end{aligned}$$

From [3, Thm. 61], we also have that for $n \geq 1$ and $x \in (-1, 1)$

$$(2.30) \quad |L_n(x)| < \sqrt{\frac{\pi}{2n(1-x^2)}}.$$

Then for $x \in (-1, 1)$ and $N \in \mathbb{N}_0$

$$\begin{aligned}
(2.31) \quad |S_N(x)| &\leq \frac{1}{(N+1)^2+1} \left[1 + \sum_{n=1}^N (2n+1) |L_n(x)| \right] \\
&\leq \frac{1}{(N+1)^2+1} \left[1 + 3 \sum_{n=1}^N n \cdot \sqrt{\frac{\pi}{2n(1-x^2)}} \right] \\
&= \frac{1}{(N+1)^2+1} \left[1 + 3 \cdot \sqrt{\frac{\pi}{2(1-x^2)}} \cdot \sum_{n=1}^N \sqrt{n} \right].
\end{aligned}$$

We can note that

$$\begin{aligned}
(2.32) \quad \sum_{n=1}^N \sqrt{n} &\leq \int_1^{N+1} \sqrt{x} \, dx \\
&= \frac{2}{3} (N+1)^{3/2} - \frac{2}{3}.
\end{aligned}$$

So

$$(2.33) \quad |S_N(x)| \leq \frac{1}{(N+1)^2+1} \left[1 + \sqrt{\frac{2\pi}{1-x^2}} \left((N+1)^{3/2} - 1 \right) \right],$$

where the right-hand side converges to 0 as $N \rightarrow \infty$ for fixed $x \in (-1, 1)$. Thus $S_N(x) \rightarrow 0$ as $N \rightarrow \infty$ for $x \in (-1, 1)$. \square

2.3. The Space X . We define the following inner product space with elements in

$$(2.34) \quad X := \mathcal{C}([-\tau, 0]; \mathbb{R}) \times \mathbb{R}$$

and the inner product defined by

$$(2.35) \quad (\Phi, \Psi)_X := \Phi^S \Psi^S + \frac{1}{\tau} (\Phi^D, \Psi^D)_{L^2([-\tau, 0])} + \Phi^D(-\tau) \Psi^D(-\tau), \quad \Phi, \Psi \in X.$$

It is relatively straight-forward to verify that $(\cdot, \cdot)_X$ is symmetric, bilinear, and positive definite and thus is an inner product. We will also make use of the norm $\|\cdot\|_X$ induced from this inner product. Note that X is **not** a Banach space since Cauchy sequences might not converge in X .

3. Uniform Convergence of Galerkin Solutions.

3.1. Pointwise Convergence in X .

It will be helpful to prove a lemma.

LEMMA 3.1. *There is $C > 0$ such that for any $N \in \mathbb{N}_0$.*

$$(3.1) \quad \|T_N(t-s)\Pi_N(\mathcal{F}(u(s)) - \mathcal{F}(u_N(s)))\|_X \leq C\|u(s) - u_N(s)\|_X,$$

where $t \in [0, T]$ and $s \in [0, t]$.

Proof. We have that

$$(3.2) \quad \begin{aligned} \|T_N(t-s)\Pi_N(\mathcal{F}(u(s)) - \mathcal{F}(u_N(s)))\|_X^2 &= \|T_N(t-s)\Pi_N(\mathcal{F}(u(s)) - \mathcal{F}(u_N(s)))\|_{\mathcal{H}}^2 \\ &\quad + \left| [T_N(t-s)\Pi_N(\mathcal{F}(u(s)) - \mathcal{F}(u_N(s)))]^D(-\tau) \right|^2. \end{aligned}$$

Note that for the first term on the right-hand side of (3.2), we have that

$$(3.3) \quad \begin{aligned} \|T_N(t-s)\Pi_N(\mathcal{F}(u(s)) - \mathcal{F}(u_N(s)))\|_{\mathcal{H}} &\leq Me^{\omega(t-s)}\|\mathcal{F}(u(s)) - \mathcal{F}(u_N(s))\|_{\mathcal{H}} \\ &\leq Me^{\omega T}\|\mathcal{F}(u(s)) - \mathcal{F}(u_N(s))\|_{\mathcal{H}} \\ &= Me^{\omega T} |f([u(s)]^D(-\tau)) - f([u_N(s)]^D(-\tau))| \\ &\leq LMe^{\omega T} |[u(s)]^D(-\tau) - [u_N(s)]^D(-\tau)| \\ &\leq LMe^{\omega T}\|u(s) - u_N(s)\|_X. \end{aligned}$$

For the second term on the right-hand side of (3.2), we consider first the case when $t-s \geq \tau$. Then

$$(3.4) \quad \begin{aligned} \left| [T_N(t-s)\Pi_N(\mathcal{F}(u(s)) - \mathcal{F}(u_N(s)))]^D(-\tau) \right| &\leq \|T_N(t-s-\tau)\Pi(\mathcal{F}(u(s)) - \mathcal{F}(u_N(s)))\|_{\mathcal{H}} \\ &\leq Me^{\omega T}\|\mathcal{F}(u(s)) - \mathcal{F}(u_N(s))\|_{\mathcal{H}} \\ &\leq LMe^{\omega T}\|u(s) - u_N(s)\|_X. \end{aligned}$$

Now consider the case when $t-s < \tau$. So we have that

$$(3.5) \quad \begin{aligned} \left| [T_N(t-s)\Pi_N(\mathcal{F}(u(s)) - \mathcal{F}(u_N(s)))]^D(-\tau) \right| &= \left| [\Pi_N(\mathcal{F}(u(s)) - \mathcal{F}(u_N(s)))]^D(t-s-\tau) \right| \\ &= |f([u(s)]^D(-\tau)) - f([u_N(s)]^D(-\tau))| \cdot |S_N^T(t-s-\tau)| \\ &\leq L |[u(s)]^D(-\tau) - [u_N(s)]^D(-\tau)| \\ &\leq L\|u(s) - u_N(s)\|_X. \end{aligned}$$

If we define

$$(3.6) \quad C := \sqrt{2} \cdot \max\{L, LMe^{\omega T}\}$$

and apply (3.3), (3.4), and (3.5) to (3.2), then we get that

$$(3.7) \quad \|T_N(t-s)\Pi_N(\mathcal{F}(u(s)) - \mathcal{F}(u_N(s)))\|_X \leq C\|u(s) - u_N(s)\|_X. \quad \square$$

We introduce the following definitions:

$$\begin{aligned}
(3.8) \quad r_N(t) &:= \|u(t) - u_N(t)\|_X, \\
\epsilon_N(t) &:= \|T(t)u_0 - T_N(t)\Pi_N u_0\|_X, \\
d_N(t, s) &:= \|(T(t-s) - T_N(t-s)\Pi_N)\mathcal{F}(u(s))\|_X.
\end{aligned}$$

One can apply the variation-of-constants formula and the above definitions to get that

$$\begin{aligned}
(3.9) \quad r_N(t) &\leq \epsilon_N(t) + \int_0^t d_N(t, s) \, ds + \int_0^t \|T_N(t-s)\Pi_N(\mathcal{F}(u(s)) - \mathcal{F}(u_N(s)))\|_X \, ds \\
&\leq \epsilon_N(t) + \int_0^t d_N(t, s) \, ds + C \int_0^t r_N(s) \, ds.
\end{aligned}$$

Applying Grönwall's inequality to (3.9) gives

$$\begin{aligned}
(3.10) \quad r_N(t) &\leq \left[\epsilon_N(t) + \int_0^t d_N(t, s) \, ds \right] + \int_0^t C e^{C(t-s)} \left[\epsilon_N(s) + \int_0^s d_N(s, r) \, dr \right] \, ds \\
&\leq \left[\epsilon_N(t) + \int_0^t d_N(t, s) \, ds \right] + C e^{CT} \int_0^t \left[\epsilon_N(s) + \int_0^s d_N(s, r) \, dr \right] \, ds.
\end{aligned}$$

We wish to show that $r_N(t) \rightarrow 0$ as $N \rightarrow \infty$ for each fixed $t \in [0, T]$. To this end, we show that each term on the right-hand side of (3.10) converges to 0 as $N \rightarrow \infty$ and $t \in [0, T]$ fixed. The following propositions will show this.

PROPOSITION 3.1. *For fixed $t \in [0, T]$,*

$$(3.11) \quad \epsilon_N(t) \rightarrow 0 \text{ and } \int_0^t \epsilon_N(s) \, ds \rightarrow 0$$

as $N \rightarrow \infty$.

Proof. From the definition of the X norm, we have that

$$(3.12) \quad \epsilon_N(t)^2 = \|T(t)u_0 - T_N(t)\Pi_N u_0\|_{\mathcal{H}}^2 + \|[T(t)u_0]^D(-\tau) - [T_N(t)\Pi_N u_0]^D(-\tau)\|^2.$$

The first term on the right-hand side converges uniformly to 0 by the Trotter-Kato theorem. For the second case, we again consider the case when $t \geq \tau$. Here we can apply the Trotter-Kato theorem again to $\|T(t-\tau)u_0 - T_N(t-\tau)\Pi_N u_0\|_{\mathcal{H}}^2$ to get the term converges to zero. When $t < \tau$, the second term becomes

$$(3.13) \quad |u_0^D(t-\tau) - [\Pi_N u_0]^D(t-\tau)|^2$$

which converges to 0 uniformly by Proposition 2.2. This gives that $\epsilon_N(t) \rightarrow 0$.

To show the other convergence, note that $\epsilon_N(s)$ converges pointwisely to 0 on $[0, t]$. Furthermore, we may uniformly bound $\epsilon_N(s)$ by again observing the equality (3.12) and applying the uniform bounds on $\|T_N(\cdot)\|_{\mathcal{H}}$ and on $[\Pi_N u_0]^D$. Then by the Bounded Convergence Theorem, we have $\int_0^t \epsilon_N(s) \, ds \rightarrow 0$. \square

PROPOSITION 3.2. For fixed $t \in [0, T]$,

$$(3.14) \quad \int_0^t d_N(t, s) ds \rightarrow 0 \text{ and } \int_0^t \int_0^s d_N(s, r) dr ds \rightarrow 0,$$

as $N \rightarrow \infty$.

Proof. We can again apply the definition of $\|\cdot\|_X$ to get that

$$(3.15) \quad d_N^2(t, s) = \|(T(t-s) - T_N(t-s)\Pi_N)\mathcal{F}(u(s))\|_{\mathcal{H}}^2 \\ + |[T(t-s)\mathcal{F}(u(s))]^D(-\tau) - [T_N(t-s)\Pi_N\mathcal{F}(u(s))]^D(-\tau)|^2.$$

For fixed t and s , the first term of the right-hand side converges to zero. For $t-s \geq \tau$ the second term will similarly converge to 0. For $t-s < \tau$, the second term will become

$$(3.16) \quad |0 - [\Pi_N\mathcal{F}(u(s))]^D(t-s-\tau)| = |f([u(s)]^D(-\tau))| \cdot |S_N(t-s-\tau)|,$$

which converges a.e. to 0 by (2.27). So for fixed t , $d_N(t, s)$ converges a.e. to 0 for $s \in [0, t]$. Furthermore, we can uniformly bound $d_N(t, s)$ by (2.26). Thus by the Bounded Convergence Theorem, we have $\int_0^t d_N(t, s) ds \rightarrow 0$ as $N \rightarrow \infty$.

The second convergence follows by the observations that $\int_0^t d_N(\cdot, r) dr$ converges pointwise to 0 by our earlier work and can uniformly bounded on $[0, t]$. This allows us to apply the Bounded Convergence Theorem to get that $\int_0^t \int_0^s d_N(s, r) dr ds \rightarrow 0$ as $N \rightarrow \infty$. \square

We may now state our final result.

THEOREM 3.3. For $t \in [0, T]$,

$$(3.17) \quad \lim_{N \rightarrow \infty} \|u(t) - u_N(t)\|_X = 0.$$

Proof. Apply propositions (3.1) and (3.2) to the inequality in (??). \square

3.2. Uniform Convergence.

LEMMA 3.2. The following convergences hold:

$$(3.18) \quad \lim_{N \rightarrow \infty} [u_N(\cdot)]^D(-\tau) = [u(\cdot)]^D(-\tau) \text{ with respect to } L^2([0, T]; \mathbb{R}),$$

and

$$(3.19) \quad \lim_{N \rightarrow \infty} \mathcal{F}(u_N(\cdot)) = \mathcal{F}(u(\cdot)) \text{ with respect to } L^1([0, T]; \mathcal{H}).$$

Proof. Note that

$$(3.20) \quad \int_0^T |[u_N(s)]^D(-\tau) - [u(s)]^D(-\tau)|^2 ds \leq \sum_{k=0}^m \int_{-\tau}^0 |[u_N(k\tau)]^D(\theta) - [u(k\tau)]^D(\theta)|^2 d\theta,$$

for m such that $T - \tau \leq m\tau < T$. In other words,

$$(3.21) \quad \|[u_N(\cdot)]^D(-\tau) - [u(\cdot)]^D(-\tau)\|_{L^2([0, T]; \mathbb{R})}^2 \leq \sum_{k=0}^m \|[u_N(k\tau)]^D - [u(k\tau)]^D\|_{L^2([-\tau, 0]; \mathbb{R})}^2.$$

It has been shown that $\|[u_N(t)]^D - [u(t)]^D\|_{L^2([0,T];\mathbb{R})} \rightarrow 0$ as $N \rightarrow \infty$ for any $t \in [0, T]$. This gives that the right side of (3.21) converges to 0 as $N \rightarrow \infty$, and thus the left side of (3.21) also converges to 0 as $N \rightarrow \infty$. This proves (3.18).

To prove the other convergence, note that

$$\begin{aligned}
(3.22) \quad \int_0^T \|\mathcal{F}(u_N(s)) - \mathcal{F}(u(s))\|_{\mathcal{H}} ds &= \int_0^T |f([u_N(s)]^D(-\tau)) - f([u(s)]^D(-\tau))| ds \\
&\leq L \int_0^T |[u_N(s)]^D(-\tau) - [u(s)]^D(-\tau)| ds \\
&= L \|[u_N(\cdot)]^D(-\tau) - [u(\cdot)]^D(-\tau)\|_{L^1([0,T];\mathbb{R})}.
\end{aligned}$$

Noting that $L^2([0, T]; \mathbb{R})$ is continuously embedded in $L^1([0, T]; \mathbb{R})$ and applying (3.18) proves that (3.19) holds. \square

THEOREM 3.4. *The sequence of functions $\{u_N\}_{N=0}^\infty$, where*

$$(3.23) \quad u_N : [0, T] \mapsto \mathcal{H}, \quad N \in \mathbb{N}_0,$$

is uniformly equicontinuous.

Proof. Suppose $t_0, t_1 \in [0, T]$ and $t_0 \leq t_1$. Denote $\delta := t_1 - t_0$. Applying the variation-of-constants formula, we have that for $N \in \mathbb{N}_0$

$$\begin{aligned}
(3.24) \quad \|u_N(t_0) - u_N(t_1)\|_{\mathcal{H}} &\leq \underbrace{\|(T_N(t_0) - T_N(t_0 + \delta))\Pi_N u_0\|_{\mathcal{H}}}_{\text{I}(\delta, N)} \\
&\quad + \underbrace{\left\| \int_0^{t_0} [T_N(t_0 - s) - T_N(t_0 + \delta - s)]\Pi_N \mathcal{F}(u_N(s)) ds \right\|_{\mathcal{H}}}_{\text{II}(\delta, N)} \\
&\quad + \underbrace{\left\| \int_{t_0}^{t_0 + \delta} T_N(t_0 + \delta - s)\Pi_N \mathcal{F}(u_N(s)) ds \right\|_{\mathcal{H}}}_{\text{III}(\delta, N)}.
\end{aligned}$$

We show that for each of these terms, the dependence on δ and N can be separated.

I. We have that

$$\begin{aligned}
(3.25) \quad \text{I}(\delta, N) &= \|T_N(t_0)(I - T_N(\delta))\Pi_N u_0\|_{\mathcal{H}} \\
&\leq M e^{\omega t_0} \|(I - T_N(\delta))\Pi_N u_0\|_{\mathcal{H}} \\
&= M e^{\omega t_0} \|(\Pi_N - T_N(\delta)\Pi_N)u_0\|_{\mathcal{H}} \\
&\leq M e^{\omega t_0} \|(I - T_N(\delta)\Pi_N)u_0\|_{\mathcal{H}} \\
&\leq M e^{\omega T} \|(I - T_N(\delta)\Pi_N)u_0\|_{\mathcal{H}} \\
&\leq M e^{\omega T} [\|(I - T(\delta))u_0\|_{\mathcal{H}} + \|(T(\delta) - T_N(\delta)\Pi_N)u_0\|_{\mathcal{H}}] \\
&\leq M e^{\omega T} \left[\|(I - T(\delta))u_0\|_{\mathcal{H}} + \sup_{t \in [0, T]} \|(T(t) - T_N(t)\Pi_N)u_0\|_{\mathcal{H}} \right].
\end{aligned}$$

Now define the following functions:

$$(3.26) \quad \mathbf{I}^*(\delta) := Me^{\omega T} \times \|(I - T(\delta))u_0\|_{\mathcal{H}}$$

and

$$(3.27) \quad \mathbf{I}^{**}(N) := Me^{\omega T} \times \sup_{t \in [0, T]} \|(T(t) - T_N(t)\Pi_N)u_0\|_{\mathcal{H}}$$

Note that $\lim_{\delta \rightarrow 0^+} \mathbf{I}^*(\delta) = 0$ by the continuity of $T(t)$ and $\lim_{N \rightarrow \infty} \mathbf{I}^{**}(N) = 0$ by the Trotter-Kato theorem.

II. We have that

$$(3.28) \quad \begin{aligned} \mathbf{II}(\delta, N) &\leq \int_0^{t_0} \|(T_N(t_0 - s) - T_N(t_0 + \delta - s))\Pi_N \mathcal{F}(u_N(s))\|_{\mathcal{H}} \, ds \\ &\leq Me^{\omega T} \int_0^{t_0} \|(I - T_N(\delta)\Pi_N) \mathcal{F}(u_N(s))\|_{\mathcal{H}} \, ds \\ &\leq Me^{\omega T} \left[\underbrace{\int_0^{t_0} \|(I - T_N(\delta)\Pi_N) \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds}_A \right. \\ &\quad \left. + \underbrace{\int_0^{t_0} \|(I - T_N(\delta)\Pi_N)(\mathcal{F}(u_N(s)) - \mathcal{F}(u(s)))\|_{\mathcal{H}} \, ds}_B \right]. \end{aligned}$$

From here, we can note that

$$(3.29) \quad \begin{aligned} A &\leq \int_0^T \|(I - T(\delta)) \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds + \int_0^T \|(T(\delta) - T_N(\delta)) \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds \\ &\leq \int_0^T \|(I - T(\delta)) \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds + \int_0^T \sup_{t \in [0, T]} \|(T(t) - T_N(t)) \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds, \end{aligned}$$

where both of these terms can easily be shown to converge to zero as $\delta \rightarrow 0$ and $N \rightarrow \infty$, respectively. Namely, we can apply the Lebesgue Dominated Convergence Theorem. Also note that

$$(3.30) \quad B \leq (1 + Me^{\omega T}) \int_0^T \|\mathcal{F}(u_N(s)) - \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds,$$

where the right-hand side converges to zero as $N \rightarrow \infty$ by (3.19). Now we set

$$(3.31) \quad \mathbf{II}^*(\delta) := Me^{\omega T} \int_0^T \|(I - T(\delta)) \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds$$

and

$$(3.32) \quad \begin{aligned} \Pi^{**}(N) := & M e^{\omega T} \left[\int_0^T \sup_{t \in [0, T]} \|(T(t) - T_N(t))\mathcal{F}(u(s))\|_{\mathcal{H}} \, ds \right. \\ & \left. + (1 + M e^{\omega T}) \int_0^T \|\mathcal{F}(u_N(s)) - \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds \right]. \end{aligned}$$

III. We have that

$$(3.33) \quad \begin{aligned} \text{III}(\delta, N) &\leq \int_{t_0}^{t_0+\delta} \|T_N(t_0 + \delta - s) \Pi_N \mathcal{F}(u_N(s))\|_{\mathcal{H}} \, ds \\ &\leq M e^{\omega T} \int_{t_0}^{t_0+\delta} \|\mathcal{F}(u_N(s))\|_{\mathcal{H}} \, ds \\ &\leq M e^{\omega T} \left[\int_{t_0}^{t_0+\delta} \|\mathcal{F}(u(s))\|_{\mathcal{H}} \, ds + \int_{t_0}^{t_0+\delta} \|\mathcal{F}(u_N(s)) - \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds \right] \\ &\leq M e^{\omega T} \left[\delta \times \sup_{t \in [0, T]} \|\mathcal{F}(u(t))\|_{\mathcal{H}} + \int_0^T \|\mathcal{F}(u_N(s)) - \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds \right]. \end{aligned}$$

Note that $\sup_{t \in [0, T]} \|\mathcal{F}(u(t))\|_{\mathcal{H}}$ is finite since $\|\mathcal{F}(u(t))\|_{\mathcal{H}}$ is a continuous function. Now let

$$(3.34) \quad \text{III}^*(\delta) := M e^{\omega T} \delta \times \sup_{t \in [0, T]} \|\mathcal{F}(u(t))\|_{\mathcal{H}}$$

and

$$(3.35) \quad \text{III}^{**}(N) := M e^{\omega T} \int_0^T \|\mathcal{F}(u_N(s)) - \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds.$$

Clearly $\lim_{\delta \rightarrow 0^+} \text{III}^*(\delta) = 0$. Also from (3.19) we have that $\lim_{N \rightarrow \infty} \text{III}^{**}(N) = 0$. Thus,

$$(3.36) \quad \begin{aligned} \|u_N(t_0) - u_N(t_1)\|_{\mathcal{H}} &\leq \text{I}(\delta, N) + \text{II}(\delta, N) + \text{III}(\delta, N) \\ &\leq [\text{I}^*(\delta) + \text{II}^*(\delta) + \text{III}^*(\delta)] + [\text{I}^{**}(N) + \text{II}^{**}(N) + \text{III}^{**}(N)]. \end{aligned}$$

Let $\epsilon > 0$. We wish to choose $\delta > 0$ such that $\|u_n(t) - u_n(t')\|_{\mathcal{H}} < \epsilon$ for any $n \in \mathbb{N}_0$ and $t, t' \in [0, T]$ with $|t - t'| < \delta$. Choosing δ^* small enough so that $\text{I}^*(\delta^*) + \text{II}^*(\delta^*) + \text{III}^*(\delta^*) < \epsilon/2$ and N large enough such that $\text{I}^{**}(N) + \text{II}^{**}(N) + \text{III}^{**}(N) < \epsilon/2$, we get that

$$(3.37) \quad \|u_n(t) - u_n(t')\|_{\mathcal{H}} < \epsilon,$$

where $|t - t'| < \delta^*$ and $n \geq N$. For each $n \in \mathbb{N}_0$ that are less than N , we pick $\delta_n > 0$ such that $\|u_n(t) - u_n(t')\|_{\mathcal{H}} < \epsilon$ for $|t - t'| < \delta_n$. This is possible since u_n is uniformly continuous on $[0, T]$. Let $\delta = \min\{\delta^*, \delta_0, \dots, \delta_{N-1}\}$. Then δ satisfies the challenge from ϵ . This proves uniform equicontinuity. \square

THEOREM 3.5. *For $T > 0$, we have that*

$$(3.38) \quad \lim_{N \rightarrow \infty} \sup_{t \in [0, T]} \|u_N(t) - u(t)\|_{\mathcal{H}} = 0.$$

Proof. The above result follows directly from [Theorem 3.3](#) and [Theorem 3.4](#). \square

4. Uniform Equicontinuity of Galerkin Solutions.

4.1. Initial Lemmas.

LEMMA 4.1. *The following convergences hold:*

$$(4.1) \quad \lim_{N \rightarrow \infty} [u_N(\cdot)]^D(-\tau) = [u(\cdot)]^D(-\tau) \text{ with respect to } L^2([0, T]; \mathbb{R}),$$

and

$$(4.2) \quad \lim_{N \rightarrow \infty} \mathcal{F}(u_N(\cdot)) = \mathcal{F}(u(\cdot)) \text{ with respect to } L^1([0, T]; \mathcal{H}).$$

Proof. Note that

$$(4.3) \quad \int_0^T |[u_N(s)]^D(-\tau) - [u(s)]^D(-\tau)|^2 \, ds \leq \sum_{k=0}^m \int_{-\tau}^0 |[u_N(k\tau)]^D(\theta) - [u(k\tau)]^D(\theta)|^2 \, d\theta.$$

In other words,

$$(4.4) \quad \|[u_N(\cdot)]^D(-\tau) - [u(\cdot)]^D(-\tau)\|_{L^2([0, T]; \mathbb{R})}^2 \leq \sum_{k=0}^m \|[u_N(k\tau)]^D - [u(k\tau)]^D\|_{L^2([0, T]; \mathbb{R})}^2.$$

It has been shown that $\|[u_N(t)]^D - [u(t)]^D\|_{L^2([0, T]; \mathbb{R})} \rightarrow 0$ as $N \rightarrow \infty$ for any $t \in [0, T]$. This gives that the right side of (4.4) converges to 0 as $N \rightarrow \infty$, and thus the left side of (4.4) also converges to 0 as $N \rightarrow \infty$. This proves (4.1).

To prove the other convergence, note that

$$(4.5) \quad \begin{aligned} \int_0^T \|\mathcal{F}(u_N(s)) - \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds &= \int_0^T |f([u_N(s)]^D(-\tau)) - f([u(s)]^D(-\tau))| \, ds \\ &\leq L \int_0^T |[u_N(s)]^D(-\tau) - [u(s)]^D(-\tau)| \, ds \\ &= L \|[u_N(\cdot)]^D(-\tau) - [u(\cdot)]^D(-\tau)\|_{L^1([0, T]; \mathbb{R})}. \end{aligned}$$

Noting that $L^2([0, T]; \mathbb{R})$ is continuously embedded in $L^1([0, T]; \mathbb{R})$ and applying (4.1) proves that (4.2) holds. \square

4.2. Uniform Equicontinuity.

THEOREM 4.1. *The sequence of functions $\{u_N\}_{N=0}^\infty$, where*

$$(4.6) \quad u_N : [0, T] \mapsto \mathcal{H}, \quad N \in \mathbb{N}_0,$$

is uniformly equicontinuous.

Proof. Suppose $t_0, t_1 \in [0, T]$ and $t_0 \leq t_1$. Denote $\delta := t_1 - t_0$. Applying the variation-of-constants formula, we have that for $N \in \mathbb{N}_0$

$$\begin{aligned}
(4.7) \quad \|u_N(t_0) - u_N(t_1)\|_{\mathcal{H}} &\leq \underbrace{\|(T_N(t_0) - T_N(t_0 + \delta))\Pi_N u_0\|_{\mathcal{H}}}_{\text{I}(\delta, N)} \\
&\quad + \underbrace{\left\| \int_0^{t_0} [T_N(t_0 - s) - T_N(t_0 + \delta - s)]\Pi_N \mathcal{F}(u_N(s)) \, ds \right\|_{\mathcal{H}}}_{\text{II}(\delta, N)} \\
&\quad + \underbrace{\left\| \int_{t_0}^{t_0 + \delta} T_N(t_0 + \delta - s)\Pi_N \mathcal{F}(u_N(s)) \, ds \right\|_{\mathcal{H}}}_{\text{III}(\delta, N)}.
\end{aligned}$$

We show that for each of these terms, the dependence on δ and N can be separated.

I. We have that

$$\begin{aligned}
(4.8) \quad \text{I}(\delta, N) &= \|T_N(t_0)(I - T_N(\delta))\Pi_N u_0\|_{\mathcal{H}} \\
&\leq M e^{\omega t_0} \|(I - T_N(\delta))\Pi_N u_0\|_{\mathcal{H}} \\
&= M e^{\omega t_0} \|(\Pi_N - T_N(\delta)\Pi_N)u_0\|_{\mathcal{H}} \\
&\leq M e^{\omega t_0} \|(I - T_N(\delta)\Pi_N)u_0\|_{\mathcal{H}} \\
&\leq M e^{\omega T} \|(I - T_N(\delta)\Pi_N)u_0\|_{\mathcal{H}} \\
&\leq M e^{\omega T} [\|(I - T(\delta))u_0\|_{\mathcal{H}} + \|(T(\delta) - T_N(\delta)\Pi_N)u_0\|_{\mathcal{H}}] \\
&\leq M e^{\omega T} \left[\|(I - T(\delta))u_0\|_{\mathcal{H}} + \sup_{t \in [0, T]} \|(T(t) - T_N(t)\Pi_N)u_0\|_{\mathcal{H}} \right].
\end{aligned}$$

Now define the following functions:

$$(4.9) \quad \text{I}^*(\delta) := M e^{\omega T} \times \|(I - T(\delta))u_0\|_{\mathcal{H}}$$

and

$$(4.10) \quad \text{I}^{**}(N) := M e^{\omega T} \times \sup_{t \in [0, T]} \|(T(t) - T_N(t)\Pi_N)u_0\|_{\mathcal{H}}$$

Note that $\lim_{\delta \rightarrow 0^+} \text{I}^*(\delta) = 0$ by the continuity of $T(t)$ and $\lim_{N \rightarrow \infty} \text{I}^{**}(N) = 0$ by the Trotter-Kato theorem.

II. We have that

$$\begin{aligned}
(4.11) \quad \Pi(\delta, N) &\leq \int_0^{t_0} \|(T_N(t_0 - s) - T_N(t_0 + \delta - s))\Pi_N \mathcal{F}(u_N(s))\|_{\mathcal{H}} \, ds \\
&\leq M e^{\omega T} \int_0^{t_0} \|(I - T_N(\delta)\Pi_N) \mathcal{F}(u_N(s))\|_{\mathcal{H}} \, ds \\
&\leq M e^{\omega T} \left[\underbrace{\int_0^{t_0} \|(I - T_N(\delta)\Pi_N) \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds}_A \right. \\
&\quad \left. + \underbrace{\int_0^{t_0} \|(I - T_N(\delta)\Pi_N)(\mathcal{F}(u_N(s)) - \mathcal{F}(u(s)))\|_{\mathcal{H}} \, ds}_B \right].
\end{aligned}$$

From here, we can note that

$$\begin{aligned}
(4.12) \quad A &\leq \int_0^T \|(I - T(\delta)) \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds + \int_0^T \|(T(\delta) + T_N(\delta)) \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds \\
&\leq \int_0^T \|(I - T(\delta)) \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds + \int_0^T \sup_{t \in [0, T]} \|(T(t) + T_N(t)) \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds,
\end{aligned}$$

where both of these terms can easily be shown to converge to zero as $\delta \rightarrow 0$ and $N \rightarrow \infty$, respectively. Namely, we can apply the Lebesgue Dominated Convergence Theorem. Also note that

$$(4.13) \quad B \leq (1 + M e^{\omega T}) \int_0^T \|\mathcal{F}(u_N(s)) - \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds,$$

where the right-hand side converges to zero as $N \rightarrow \infty$ by (4.2). Now we set

$$(4.14) \quad \Pi^*(\delta) := M e^{\omega T} \int_0^T \|(I - T(\delta)) \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds$$

and

$$\begin{aligned}
(4.15) \quad \Pi^{**}(N) &:= M e^{\omega T} \left[\int_0^T \sup_{t \in [0, T]} \|(T(t) + T_N(t)) \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds \right. \\
&\quad \left. + (1 + M e^{\omega T}) \int_0^T \|\mathcal{F}(u_N(s)) - \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds \right].
\end{aligned}$$

III. We have that

$$\begin{aligned}
\text{III}(\delta, N) &\leq \int_{t_0}^{t_0+\delta} \|T_N(t_0 + \delta - s) \Pi_N \mathcal{F}(u_N(s))\|_{\mathcal{H}} \, ds \\
&\leq M e^{\omega T} \int_{t_0}^{t_0+\delta} \|\mathcal{F}(u_N(s))\|_{\mathcal{H}} \, ds \\
(4.16) \quad &\leq M e^{\omega T} \left[\int_{t_0}^{t_0+\delta} \|\mathcal{F}(u(s))\|_{\mathcal{H}} \, ds + \int_{t_0}^{t_0+\delta} \|\mathcal{F}(u_N(s)) - \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds \right] \\
&\leq M e^{\omega T} \left[\delta \times \sup_{t \in [0, T]} \|\mathcal{F}(u(t))\|_{\mathcal{H}} + \int_0^T \|\mathcal{F}(u_N(s)) - \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds \right].
\end{aligned}$$

Note that $\sup_{t \in [0, T]} \|\mathcal{F}(u(t))\|_{\mathcal{H}}$ is finite since $\|\mathcal{F}(u(t))\|_{\mathcal{H}}$ is a continuous function. Now let

$$(4.17) \quad \text{III}^*(\delta) := M e^{\omega T} \delta \times \sup_{t \in [0, T]} \|\mathcal{F}(u(t))\|_{\mathcal{H}}$$

and

$$(4.18) \quad \text{III}^{**}(N) := M e^{\omega T} \int_0^T \|\mathcal{F}(u_N(s)) - \mathcal{F}(u(s))\|_{\mathcal{H}} \, ds.$$

Clearly $\lim_{\delta \rightarrow 0^+} \text{III}^*(\delta) = 0$. Also from (4.2) we have that $\lim_{N \rightarrow \infty} \text{III}^{**}(N) = 0$. Thus,

$$\begin{aligned}
(4.19) \quad \|u_N(t_0) - u_N(t_1)\|_{\mathcal{H}} &\leq \text{I}(\delta, N) + \text{II}(\delta, N) + \text{III}(\delta, N) \\
&\leq [\text{I}^*(\delta) + \text{II}^*(\delta) + \text{III}^*(\delta)] + [\text{I}^{**}(N) + \text{II}^{**}(N) + \text{III}^{**}(N)].
\end{aligned}$$

Let $\epsilon > 0$. We wish to choose $\delta > 0$ such that $\|u_n(t) - u_n(t')\|_{\mathcal{H}} < \epsilon$ for any $n \in \mathbb{N}_0$ and $t, t' \in [0, T]$ with $|t - t'| < \delta$. Choosing δ^* small enough so that $\text{I}^*(\delta^*) + \text{II}^*(\delta^*) + \text{III}^*(\delta^*) < \epsilon/2$ and N large enough such that $\text{I}^{**}(N) + \text{II}^{**}(N) + \text{III}^{**}(N) < \epsilon/2$, we get that

$$(4.20) \quad \|u_n(t) - u_n(t')\|_{\mathcal{H}} < \epsilon,$$

where $|t - t'| < \delta^*$ and $n \geq N$. For each $n \in \mathbb{N}_0$ that are less than N , we pick $\delta_n > 0$ such that $\|u_n(t) - u_n(t')\|_{\mathcal{H}} < \epsilon$ for $|t - t'| < \delta_n$. This is possible since u_n is uniformly continuous on $[0, T]$. Let $\delta = \min\{\delta^*, \delta_0, \dots, \delta_{N-1}\}$. Then δ satisfies the challenge from ϵ . This proves uniform equicontinuity. \square

REFERENCES

- [1] M. D. CHEKROUN, M. GHIL, H. LIU, AND S. WANG, *Low-dimensional galerkin approximations of nonlinear delay differential equations*, Discrete and Continuous Dynamical Systems, 36 (2016), pp. 4133–4177.
- [2] T. H. KOORNWINDER, *Orthogonal polynomials with weight function $(1-x)^\alpha(1+x)^\beta + M\delta(x+1) + N\delta(x-1)$* , Canadian mathematical bulletin, 27 (1984), pp. 205–214.
- [3] E. D. RAINVILLE, *Special functions*, Macmillan, New York, 1971.
- [4] H. WANG AND S. XIANG, *On the convergence rates of legendre approximation*, Mathematics of Computation, 81 (2012), pp. 861–877.