



## The Structure of our Milky Way Disk:



Diameter  $\sim 100,000$  light years  
 Mass  $\sim 60$  billion times the mass of the Sun

1

## A Model of the Milky Way: A Spiral Galaxy



2

## Thick/Thin Disk





Fig 2.8 (Reid, Knude) 'Galaxies in the Universe' Sparke/Gallagher CUP 2007

3

## Thick / Thin Disks

- Stars of different ages also have **different kinematics**:



- We can define 1 D velocity dispersion of a group of stars in a given direction:  

$$\sigma_z^2 \equiv \langle v_z^2 \rangle - \langle v_z \rangle^2$$
- Young stars (age  $< 3\text{Gyr}$ ) have  $\sigma_z \sim 15\text{km/s}$  while old, low metallicity, thick disk stars have  $\sigma_z \sim 25\text{-}40\text{km/s}$ ,  $\sigma_R$  is typically significantly larger!

4

## An aside: what is meant by [ Fe/ H]

$$[A/B] = \log_{10} \left[ \frac{(\text{number of A atoms}/\text{number of B atoms})_\star}{(\text{number of A atoms}/\text{number of B atoms})_\odot} \right]$$

$$[A/B] \equiv \log_{10} \left( \frac{A}{B} \right)_\star - \log_{10} \left( \frac{A}{B} \right)_\odot$$

$\alpha$  refers to "alpha process elements" which are the result of a class of fusion processes involving helium. The  $\alpha$  elements multiples of helium nuclei (" $\alpha$  particles"): integer multiple of 4 nucleons. The  $\alpha$  elements are O, Ne, Mg, Si, S, Ar, Ca, Ti

5


## An Evolving Picture: Thick Disk

*How much of the total stellar surface mass density at R0 comes from stars with a scale-height  $h_z$ ?*

If there were a distinct 'thin' and 'thick' disk, one would expect a bimodal distribution in  $\Sigma_{R0}(h_z)$ , with  $\approx 85\%$  of it in a thin disk peak (with scale heights covering the range  $h_z \approx 100$  to  $250$  pc) and  $\approx 15\%$  in a thick disk (with  $h_z \approx 700$  pc) for a canonical disk decomposition (e.g., Juric et al 2008). However, the Figure shows a different picture. There is a continuous distribution of  $\Sigma_{R0}(h_z)$ , with no sign of bimodality, a distribution that is quite well-approximated by  $\Sigma_{R0}(h_z) \propto \exp(-h_z/280$  pc). Rix & Bovy 2013


6

Some recent work questions whether there is a distinction between the "thin" and "thick" disks.




No bimodality in scale height with age or metallicity!

## Milky Way at different wavelengths



## The MW Bar

Short  $\sim 3.0$  kpc (Freudenreich 1997)  
Long  $\sim 4.5$  kpc (Cabrera-Lambers et al 08, Benjamin 2009)




7

8

## An Evolving Picture: X-Shaped Bulge/Bar

Ness & Lang 2016




The WISE W1 and W2 image fit by a simple exponential disk model, making the X structure more apparent. Top-left: Data. Top-middle: Data, masking out the top and bottom 5% of pixels based on W1 – W2 color, as well as pixels with negative flux. The diagonal structure at the top of the image is due to scattered light from the Moon in the unWISE coadds. Top-right: Exponential disk model fit. Bottom-left: Residuals (data minus model). Bottom-middle: Masked residuals. Bottom-right: 50-pixel ( $\sim 1.7^\circ$ ) median filter of masked residuals (median of unmasked pixels).

9

## Formation of an X-Shaped/Boxy Bulge


Abbott, Valluri + 2017



10

## Formation of an X-Shaped/Boxy Bulge

Abbott, Valluri + 2017




BP/X shape in the N-body model is produced largely by the box orbit family that constitutes the majority of orbits in the bar.

Bar Buckling: when  $\sigma_z / \sigma_x < 0.4$

Figure 8. Projected excess-mass density on the x-z plane for each of the five orbit families from Model A as labelled. The colour gradient as shown in the bar on the right of each plot is in units of  $M_\odot \text{ kpc}^{-3}$ . The box orbits (top row left) and very resonant 2:-2:1 x1+banana orbits (top row right) show a clear BP/X-shape at large radii. The x-tube orbits (top row middle) show a hint of an X-shape but at very low-density levels. Resonant boxlet 3:-2:0 fish/prezel orbits (bottom row left) and resonant 3:0:-5 brezel orbits (bottom row right) show a BP/X-shape at small radii ( $\le 2$  kpc).

## The Magellanic System & the CGM




$$M_{\text{Gas}} \text{ outside } \sim 2 \times 10^9 M_\odot (d/55 \text{ kpc})^2 > 2 \times M_{\text{Gas LMC+SMC}}$$

11


12

## Circumgalactic Medium (CGM) in Cold Gas (HI)




13

## Circumgalactic Medium (CGM): Baryon Cycle




14

## Stellar Halo



Bullock & Johnson 2005



**Figure 4.** Initial number density profiles of the stellar halo ( $\#/ \text{kpc}^3$ ), built by applying the stellar tracer method, outlined in Laporte et al. (2013a,b), and using the observed density profiles for K-giants (MW-X; dashed line Xue et al. 2015) and RR Lyrae (Hernitschek et al. 2018, MW-H; dotted line). The DM density profile for the MW halo (solid line) is shown for comparison.

Garavito-Camargo + 2019

15

## Stellar Halo: Einasto Profiles


$$\begin{aligned} \text{Number of stars per } \text{kpc}^3 \\ \nu(r) = \nu_e e^{-dn(r/r_{\text{eff}})^{(1/n)-1}} \\ dn = 3n - 1/3 + 0.0079/n. \end{aligned}$$

	MW-X	MW-H
Density profile	Einasto	Einasto
$(n; r_{\text{eff}})$	(3.1; 15 kpc)	(9.53; 1.07 kpc)
Distances (kpc)	10–80	20–131
Tracers	K-giants	RRLyR
Reference	Xue et al. (2015)	Hernitschek et al. (2018)

**Table 4.** MW stellar halo model parameters from observations of K-giants and RR Lyrae (RRLyR).

16


## Dark Matter vs. Stars



Wetzel+2016

17


## Our Milky Way has many Neighbors



26 Satellites as of Feb 2015

18

Possible New Dwarfs found by DES Survey  
+PANSTARRS +SMASH survey = ~ 50 ?




Bechtol + 2015 (DES); Koposov+2015; Laevens +2015;  
Martin, Nidever, Besla+ 2015 (SMASH), Drlica-Wagner+2015  
(DES), Homma+2016 ... more to come with LSST!

19

## GAIA

Gaia is a space observatory operated by the European Space Agency. It is designed for astrometry.

**ASTROMETRY:**  
measuring the positions, distances and velocities of stars with unprecedented precision.



20





Fig 2.1 'Galaxies in the Universe' Sparke/Gallagher CUP 2007

## Gaia is orbiting L2


Positions to 20 micro arc second! Each star is observed 70 times: parallaxes and proper motions (motion on the sky).  
Hipparcos: measured positions of bright stars to only 0.001"



21

22

## Gaia: First Geometric Indication of the Galactic Bar



## $V_{\tan}$ --> Proper Motion

The **proper motion** of an object is the measurement of its angular change in position on the sky over time.

### Barnard's Star:

Highest proper motion of any star visible from the Earth.

True motion: 140 km/s




Credit: Steve Quirk


23

24

## GAIA



25



26

## HSTPROMO

The HST Proper Motion Collaboration

<http://www.stsci.edu/~marel/hstpromo.html>

- Characteristic velocity accuracy necessary  
~10 km/s at 70 kpc (Milky Way halo/satellite dynamics)
- Corresponding PM accuracy  
~ 30  $\mu$ as / yr

(~ speed of human hair growth  
at distance of the Moon)

With HST we can measure a  
change of 0.006 ACS/WFC pixels  
over a 10 yr baseline



## The Need for HST:

Both Gaia & HST in concert are needed to develop a model for the dynamics of our Local Group.

Galaxy	$\Delta\text{PM}$ (km/s)	
	HST	Gaia (2017) <sup>2018</sup>
Horologium I	23	45
Pictoris II	24	119
Phoenix II	25	125
Hydra II	27	138
Grus I	26	138
Eridanus III	23	151
Tucana II	24	59

< 30 kpc: Gaia  
30-80 kpc: Gaia & HST  
> 80 kpc: HST




HST Cycle 23 Large  
PI Kallivayalil

28


29

## Our Local Group of Galaxies



30

What is the 3D velocity of M31?




31

HSTPROMO: The First Direct Proper Motion Measurement of M31



Sohn + 2012 (12 μas accuracy) - M31 is coming straight at us!

Andromeda is heading DIRECTLY towards us!



32


33

With the new M31 proper motion measurement we can predict the timing of the collision between the MW & M31:  $3.87^{+0.42}_{-0.32}$  Gyr van der Marel,Besla+2012



34

How might the night sky change?




Credit: Besla, Frank Summers

36

## Simulating a Galaxy Collision

The N-body problem:  
solve for the dynamics of  
a group of N objects in  
3D space under the  
influence of mutual  
gravity.



Doing this by hand would suck!  
Turn to computers!

35

Q: What are the Future Prospects  
for Astronomy in 6 Billion Years?



37