

# Primer on Topology for AlgTop

This is a collection of notes intended to provide the minimal necessary topology background to ramp someone up to algebraic topology.

Trey Plante

## Contents

<b>1</b>	<b>Topological Spaces</b>	<b>1</b>
1.1	Bases . . . . .	2
1.2	Product Topology . . . . .	3

## 1 Topological Spaces

**Definition 1.1** (Lee). A *topology* on a set  $X$  is a collection  $\mathcal{T}$  of subsets of  $X$ , called *open sets*, satisfying the following:

1.  $X \in \mathcal{T}$  and  $\emptyset \in \mathcal{T}$
2. If  $U_1, \dots, U_n \in \mathcal{T}$ , then  $U_1 \cap \dots \cap U_n \in \mathcal{T}$
3. If  $\{U_\alpha\}_{\alpha \in A}$  is a collection of elements of  $\mathcal{T}$ , then  $\cup_{\alpha \in A} U_\alpha$  is in  $\mathcal{T}$ .

**Definition 1.2** (Lee). A *topological space* is a pair  $(X, \mathcal{T})$  consisting of a set  $X$  and a topology  $\mathcal{T}$  on  $X$ .

*Remark 1.3* (Lee). A neighborhood of  $q \in X$  is an open set containing  $q$ .

**Lemma 1.4** (Exercise 2.1 (Munkres)). *Let  $(X, \mathcal{T})$  be a topological space; let  $A$  be a subset of  $X$ . Suppose that for each  $x \in A$  there is an open set  $U$  containing  $x$  such that  $U \subset A$ . Then,  $A$  is open in  $X$ .*

*Proof.* Suppose that  $A \subset X$  and that for all  $x \in A$  there is an open set  $U$  containing  $x$  such that  $U \subset A$ . By definition,  $\cup_{x \in A} U_x \in \mathcal{T}$  because each  $U_x \in \mathcal{T}$ . Then,  $\cup_{x \in A} U_x = A$  so  $A \in \mathcal{T}$ .  $\square$

**Lemma 1.5** (Exercise 2.3 (Munkres)). *Is the collection*

$$\mathcal{T}_{\text{inf}} = \{U | X - U \text{ is infinite or empty or all of } X\}$$

*a topology on  $X$ ?*

*Proof.*

1. If the condition is all of  $X$ , then  $U = \emptyset$ , so  $\emptyset \in \mathcal{T}$ . If the condition is empty, then  $U = X$ , so  $X \in \mathcal{T}$ .
2. Let  $U_1, \dots, U_n \in \mathcal{T}$ . For  $U = \cap_i U_i$ , we have

$$X - U = X - \cap_i U_i = \cup_i (X - U_i)$$

.

If any of the  $X - U_i$  is infinite, all of  $X$ , or empty, the union meets the corresponding condition.

3. Let  $U_1, U_2, \dots \in \mathcal{T}$ . For  $U = \cup_i U_i$ , we have

$$X - U = X - \cup_i U_i = \cap_i (X - U_i)$$

.

If all of the  $X - U_i$  are infinite, the intersection does not necessarily meet any of the conditions. Thus,  $\mathcal{T}_{\text{inf}}$  is not a topology on  $X$ .

□

**Definition 1.6** (Munkres). A topology  $\mathcal{T}$  is finer than  $\mathcal{T}'$  if  $\mathcal{T}' \subseteq \mathcal{T}$  and, analogously,  $\mathcal{T}'$  is coarser.

## 1.1 Bases

**Definition 1.7** (Munkres). If  $X$  is a set, a *basis* for a topology on  $X$  is a collection  $\mathcal{B}$  of subsets of  $X$  such that

1. For each  $x \in X$ , there is at least one basis element  $B$  containing  $x$ .
2. If  $x$  belongs to the intersection of two basis elements  $B_1$  and  $B_2$ , then there is a basis element  $B_3$  containing  $x$  such that  $B_3 \subset B_1 \cap B_2$ .

*Remark 1.8.* If  $\mathcal{B}$  satisfies both of the above conditions, then we can define the topology  $\mathcal{T}$  generated by  $\mathcal{B}$  as follows: A subset  $U$  of  $X$  is said to be open in  $X$  if for each  $x \in U$ , there is a basis element  $B \in \mathcal{B}$  such that  $x \in B$  and  $B \subset U$ . Each basis element itself is an element of  $\mathcal{T}$ .

**Lemma 1.9.** *Let  $\mathcal{T}$  be the collection generated by the basis  $\mathcal{B}$  on  $X$ .  $\mathcal{T}$  is a topology.*

*Proof.* •  $X \in \mathcal{T}$  because for each  $x \in X$  there is a basis element  $B$  with  $x \in B$  and  $B \subset X$ .  $\emptyset \in \mathcal{T}$  vacuously because for each  $x \in \emptyset$  is a vacuous statement.

- Suppose  $U_1, \dots, U_n \in \mathcal{T}$ . We want to show  $U_1 \cap \dots \cap U_n \in \mathcal{T}$ . For each  $x \in U_1 \cap \dots \cap U_n$ , we have that there are basis elements  $x \in B_1, \dots, x \in B_n$  because each  $U_i \in \mathcal{T}$ . We also know  $U_i \subset B_i$ . By the definition of basis, there is some basis element  $B \subset B_1 \cap \dots \cap B_n$  such that  $x \in B$ . Then,  $x \in B$  and  $B \subset U_1 \cap \dots \cap U_n$  (by  $B \subset U_i$ ). Thus,  $U_1 \cap \dots \cap U_n \in \mathcal{T}$ .

- Suppose  $U_i \in \mathcal{T}$ . We want to show that  $\cup_i U_i \in \mathcal{T}$ . For each  $x \in \cup_i U_i$ , we have that  $x \in U_i$  and there is a basis element  $x \in B$  with  $B \subset U_i$ . Then,  $B \subset \cup_i U_i$ . Thus,  $\cup_i U_i \in \mathcal{T}$ .

□

exercises + lemmas

**Lemma 1.10** (Munkres). *Let  $X$  be a set and  $\mathcal{B}$  be a basis for a topology  $\mathcal{T}$  on  $X$ . Then  $\mathcal{T}$  is the collection of all unions of elements of  $\mathcal{B}$ .*

*Remark 1.11.* This lemma means that we can express every open set  $U$  of  $X$  as a union of basis elements.

**Lemma 1.12** (Munkres). *Let  $X$  be a topological space. Suppose that  $\mathcal{C}$  is a collection of open sets of  $X$  such that for each open set  $U$  and  $x \in U$ , there is  $C \in \mathcal{C}$  such that  $x \in C \subset U$ . Then  $\mathcal{C}$  is a basis for the topology of  $X$ .*

*Proof.* It suffices to show (1) that  $\mathcal{C}$  is a basis and (2) the topology generated by  $\mathcal{C}$  is the same as the topology of  $X$ .

*Part 1:  $\mathcal{C}$  is a basis*

For the first condition, we must check that for each  $x \in X$ , there is a  $C \in \mathcal{C}$  containing  $x$ . Let  $x \in X$ . Because  $X$  is an open set of  $X$ , we have, by supposition, that there is  $C \in \mathcal{C}$  such that  $x \in C \subset X$ .

For the second condition, we must check that if  $x \in C_1 \cap C_2$ , then there is  $C_3 \subset C_1 \cap C_2$ . Let  $x \in C_1 \cap C_2$ . Because  $C_1$  and  $C_2$  are open, so is  $C_1 \cap C_2$ . By supposition, there is  $C_3 \in \mathcal{C}$  such that  $x \in C_3 \subset C_1 \cap C_2$ .

*Part 2:  $\mathcal{C}$  generates the topology of  $X$*

Let  $\mathcal{T}$  be the collection of open sets of  $X$  and let  $\mathcal{T}'$  be the topology generated by  $\mathcal{C}$ .

Suppose that  $U$  belongs to  $\mathcal{T}$ . Then for each  $x \in U$ , there is  $C \in \mathcal{C}$  such that  $x \in C \subset U$ . Then, by the definition of basis,  $U$  is in  $\mathcal{T}'$ .

Suppose that  $U$  belongs to  $\mathcal{T}'$ . Then,  $U$  is a union of elements of  $\mathcal{C}$  by the preceding lemma. Each element of  $\mathcal{C}$  belongs to  $\mathcal{T}$  and a topology is closed under arbitrary unions so  $U$  belongs to  $\mathcal{T}$ . □

## 1.2 Product Topology

**Definition 1.13** (Lee). Suppose  $X_1, \dots, X_n$  are topological spaces. Then let  $\mathcal{B} = \{U_1 \times \dots \times U_n : U_i \text{ is open in } X_i, i = 1, \dots, n\}$ . The topology generated by  $\mathcal{B}$  is the product topology.

*Remark 1.14.* We can say that the product topology  $X \times Y$  is the topology that has as its basis  $\mathcal{B}$  the collection of all sets of the form  $U \times V$  where  $U$  is open in  $X$  and  $V$  is open in  $Y$ .

Then,  $\mathcal{B}$  meets the first condition of a basis because  $X \times Y$  is itself a basis element. For the second condition, consider  $(U_1 \times V_1) \cap (U_2 \times V_2) = (U_1 \cap U_2) \times (V_1 \cap V_2)$ . Because the binary intersections are in  $X$  and  $Y$  (open), we have that the intersection is a basis element. We can generalize this to  $n$ -ary products.

**Theorem 1.15.** *If  $\mathcal{B}$  is a basis for the topology of  $X$  and  $\mathcal{C}$  is a basis for the topology of  $Y$ , then*

$$\mathcal{D} = \{B \times C \mid B \in \mathcal{B} \text{ and } C \in \mathcal{C}\}$$

*is a basis for the topology of  $X \times Y$ .*

*Proof.* Suppose that  $\mathcal{B}$  is a basis for the topology of  $X$  and  $\mathcal{C}$  is a basis for the topology of  $Y$ .

By Lemma 1.12, it suffices to show that  $\mathcal{D}$  is a collection of open sets of  $X \times Y$  such that for each open set  $Z$  and  $z \in Z$ , there is a  $D \in \mathcal{D}$  such that  $x \in D \subset Z$ .

Let  $Z$  be an arbitrary open set of  $X \times Y$  and  $z \in Z$ . □

**Definition 1.16.** Subspace topology ...  
ch3

**Definition 1.17.** Closed set and limit point ...

**Definition 1.18.** Continuous function ...

**Definition 1.19.** Metric space ....

**Definition 1.20.** Quotient topology ...

**Definition 1.21.** Connected space

**Definition 1.22.** Component and path component ...

**Definition 1.23.** Compact space ...

**Definition 1.24.** hausdorff space ...

**Definition 1.25.** The separation axioms, Urysohn's lemma and the Tietze extension theorem

**Definition 1.26.** homotopy and the fundamental group ... ch7

**Definition 1.27.** group theory ...  
ch9