Archived learning-rate diagnostic judgments

	Seed 947	Seed 732	Seed 619
S4 flat	adequate	adequate	adequate
S4ND 2-D	adequate	too low	adequate
S4ND 3-D	adequate	adequate	adequate
S4ND 4-D	adequate	adequate	too high
Mamba-2 flat	adequate	adequate	adequate
Mamba-ND 2-D	adequate	adequate	adequate
Mamba-ND 3-D	adequate	adequate	adequate
Mamba-ND 4-D	adequate	adequate	adequate
GRU flat	adequate	adequate	adequate
GRU ASA 2-D	adequate	adequate	adequate
GRU ASA 3-D	adequate	too high	adequate
GRU ASA 4-D	adequate	adequate	adequate
GRU CoFA 2-D	adequate	adequate	adequate
GRU CoFA 3-D	adequate	adequate	adequate
GRU CoFA 4-D	adequate	adequate	adequate
GRU FA 2-D	too high	adequate	adequate
GRU FA 3-D	too high	adequate	adequate
GRU FA 4-D	adequate	too high	adequate
LSTM flat	adequate	adequate	adequate
LSTM ASA 2-D	adequate	too low	adequate
LSTM ASA 3-D	adequate	adequate	adequate
LSTM ASA 4-D	adequate	too low	adequate
LSTM CoFA 2-D	adequate	too low	adequate
LSTM CoFA 3-D	adequate	adequate	adequate
LSTM CoFA 4-D	adequate	adequate	adequate
LSTM FA 2-D	adequate	adequate	adequate
LSTM FA 3-D	adequate	adequate	adequate
LSTM FA 4-D	adequate	adequate	adequate
Tr flat	adequate	too high	too high
Tr ASA 2-D	too high	adequate	adequate
Tr ASA 3-D	adequate	adequate	too high
Tr ASA 4-D	too high	too high	adequate
Tr CoFA 2-D	adequate	adequate	adequate
Tr CoFA 3-D	adequate	adequate	adequate
Tr CoFA 4-D	adequate	adequate	adequate
Tr FA 2-D	adequate	adequate	too high
Tr FA 3-D	adequate	adequate	adequate
Tr FA 4-D	adequate	adequate	adequate

38 displayed configurations. These categories are diagnostic hypotheses, not demonstrated optimal learning rates.