

Quantifying Predicates

Having predicates by themselves is no real improvement on propositional logic. We need a way to be able to talk about multiple propositions (even an infinite number of them) at the same time. For this we have quantifiers

There are two common quantifiers: \forall and \exists .

The Universal Quantifier \forall

We use the **universal quantifier** when we want to assert that a predicate is true for every element in a given set. So

$$(\forall x \in S)[P(x)]$$

says that $P(x)$ is true for every x in the set S . Notice that it is now a proposition, either true or false.

The Universal Quantifier \forall

We use the **universal quantifier** when we want to assert that a predicate is true for every element in a given set. So

$$(\forall x \in S)[P(x)]$$

says that $P(x)$ is true for every x in the set S . Notice that it is now a proposition, either true or false.

For example $(\forall x \in \mathbb{R})[x^2 \geq 0]$ is a true sentence, but $(\forall x \in \mathbb{R})[x^2 \geq 42]$ is false.

The Universal Quantifier \forall

We use the **universal quantifier** when we want to assert that a predicate is true for every element in a given set. So

$$(\forall x \in S)[P(x)]$$

says that $P(x)$ is true for every x in the set S . Notice that it is now a proposition, either true or false.

For example $(\forall x \in \mathbb{R})[x^2 \geq 0]$ is a true sentence, but $(\forall x \in \mathbb{R})[x^2 \geq 42]$ is false.

If the set is obvious (or the set of everything) then the set can be omitted and at times we can include extra information in the quantifier.

The Universal Quantifier \forall

We use the **universal quantifier** when we want to assert that a predicate is true for every element in a given set. So

$$(\forall x \in S)[P(x)]$$

says that $P(x)$ is true for every x in the set S . Notice that it is now a proposition, either true or false.

For example $(\forall x \in \mathbb{R})[x^2 \geq 0]$ is a true sentence, but $(\forall x \in \mathbb{R})[x^2 \geq 42]$ is false.

If the set is obvious (or the set of everything) then the set can be omitted and at times we can include extra information in the quantifier. E.g., in $(\forall x > 0)[e^{\ln x} = x]$ the fact that $x \in \mathbb{R}$ is implied and

The Universal Quantifier \forall

We use the **universal quantifier** when we want to assert that a predicate is true for every element in a given set. So

$$(\forall x \in S)[P(x)]$$

says that $P(x)$ is true for every x in the set S . Notice that it is now a proposition, either true or false.

For example $(\forall x \in \mathbb{R})[x^2 \geq 0]$ is a true sentence, but $(\forall x \in \mathbb{R})[x^2 \geq 42]$ is false.

If the set is obvious (or the set of everything) then the set can be omitted and at times we can include extra information in the quantifier. E.g., in $(\forall x > 0)[e^{\ln x} = x]$ the fact that $x \in \mathbb{R}$ is implied and the inequality in the quantifier is a short way of saying

$$(\forall x \in \mathbb{R})[(x > 0) \Rightarrow (e^{\ln x} = x)].$$

Lincoln's Predicate

As an example, consider President Abraham Lincoln's reference to "... the proposition that all men are created equal".

Lincoln's Predicate

As an example, consider President Abraham Lincoln's reference to "... the proposition that all men are created equal". Taking a modern interpretation and creating the predicates $\text{Person}(x)$ to mean that "x is a person" and $\text{CreatedEqual}(x, y)$ to mean that "x and y were created equal" we can make the sentence/proposition

$$(\forall x)(\forall y)[(\text{Person}(x) \wedge \text{Person}(y)) \Rightarrow \text{CreatedEqual}(x, y)].$$

Lincoln's Predicate

As an example, consider President Abraham Lincoln's reference to "... the proposition that all men are created equal". Taking a modern interpretation and creating the predicates $\text{Person}(x)$ to mean that "x is a person" and $\text{CreatedEqual}(x, y)$ to mean that "x and y were created equal" we can make the sentence/proposition

$$(\forall x)(\forall y)[(\text{Person}(x) \wedge \text{Person}(y)) \Rightarrow \text{CreatedEqual}(x, y)].$$

Or we could look at Aristotle's dictum "All men are mortal".

Lincoln's Predicate

As an example, consider President Abraham Lincoln's reference to "... the proposition that all men are created equal". Taking a modern interpretation and creating the predicates $\text{Person}(x)$ to mean that "x is a person" and $\text{CreatedEqual}(x, y)$ to mean that "x and y were created equal" we can make the sentence/proposition

$$(\forall x)(\forall y)[(\text{Person}(x) \wedge \text{Person}(y)) \Rightarrow \text{CreatedEqual}(x, y)].$$

Or we could look at Aristotle's dictum "All men are mortal". Let's define our predicates as $P(x)$, meaning "x is a person", and $M(x)$, meaning "x is mortal". Then the statement would be written as

Lincoln's Predicate

As an example, consider President Abraham Lincoln's reference to "... the proposition that all men are created equal". Taking a modern interpretation and creating the predicates $\text{Person}(x)$ to mean that "x is a person" and $\text{CreatedEqual}(x, y)$ to mean that "x and y were created equal" we can make the sentence/proposition

$$(\forall x)(\forall y)[(\text{Person}(x) \wedge \text{Person}(y)) \Rightarrow \text{CreatedEqual}(x, y)].$$

Or we could look at Aristotle's dictum "All men are mortal". Let's define our predicates as $P(x)$, meaning "x is a person", and $M(x)$, meaning "x is mortal". Then the statement would be written as

$$(\forall x)[P(x) \Rightarrow M(x)].$$

Proof by Induction

Proof by induction or **mathematical induction** is a method of proving a predicate is true for all of an infinite sequence of numbers, usually the natural numbers. It uses the “argument” below

$$\frac{\begin{array}{l} P(0) \\ (\forall k \in \mathbb{N})[P(k) \Rightarrow P(k+1)] \end{array}}{\therefore (\forall n \in \mathbb{N})[P(n)]}$$

Base Case (pointing to $P(0)$)
Inductive step (pointing to $P(k) \Rightarrow P(k+1)$)

$$\begin{array}{l} P(0) \\ (k=0) \quad P(0) \Rightarrow P(1) \\ \hline \therefore P(1) \\ (k=1) \quad P(1) \Rightarrow P(2) \\ \hline \therefore P(2) \end{array}$$

to prove that something is true for all natural numbers.

(Equivalently and more commonly, we could prove that that $P(k-1) \Rightarrow P(k)$ for almost all natural numbers (i.e. not zero). Also zero doesn't have to be the first number that P is true for.)

A First Example of Proof by Induction

Lemma

For all natural numbers n , $\sum_{k=1}^n k = \frac{n(n+1)}{2}$.

Proof.

BASE CASE: ($n=1$) ✓
 $\sum_{k=1}^1 k = 1 \stackrel{?}{=} \frac{1(1+1)}{2}$

INDUCTIVE STEP: $\left(\sum_{k=1}^L k = \frac{L(L+1)}{2} \Rightarrow \sum_{k=1}^{L+1} k = \frac{(L+1)(L+2)}{2} \right)$

Assume $\sum_{k=1}^L k = \frac{L(L+1)}{2}$

$$\sum_{k=1}^{L+1} k = \sum_{k=1}^L k + (L+1) = \frac{L(L+1)}{2} + (L+1) = \frac{L^2+L}{2} + \frac{2L+2}{2} = \frac{L^2+3L+2}{2}$$

$$1+2+3+\dots+(L-1)+L+(L+1)$$

$$\frac{(L+1)(L+2)}{2}$$

□

The size of the power set, second visit

Lemma

If S is a finite set then $|\mathcal{P}(S)| = 2^{|S|}$.

The size of the power set, second visit

Lemma

If S is a finite set then $|\mathcal{P}(S)| = 2^{|S|}$.

The idea behind this proof is that the formula works for the empty set (because $\mathcal{P}(\emptyset) = \{\emptyset\}$ so $|\mathcal{P}(\emptyset)| = 1 = 2^0 = 2^{|\emptyset|}$)

The size of the power set, second visit

Lemma

If S is a finite set then $|\mathcal{P}(S)| = 2^{|S|}$.

The idea behind this proof is that the formula works for the empty set (because $\mathcal{P}(\emptyset) = \{\emptyset\}$ so $|\mathcal{P}(\emptyset)| = 1 = 2^0 = 2^{|\emptyset|}$) and we could construct any finite set starting with the empty set and repeatedly adding elements of the desired set.

The size of the power set, second visit

Lemma

If S is a finite set then $|\mathcal{P}(S)| = 2^{|S|}$.

The idea behind this proof is that the formula works for the empty set (because $\mathcal{P}(\emptyset) = \{\emptyset\}$ so $|\mathcal{P}(\emptyset)| = 1 = 2^0 = 2^{|\emptyset|}$) and we could construct any finite set starting with the empty set and repeatedly adding elements of the desired set.

Now if T is a set such that $|\mathcal{P}(T)| = 2^{|T|}$ and S is set made by adding a new element to T (i.e. $S = T \cup \{a\}$, where $a \notin T$)

The size of the power set, second visit

Lemma

If S is a finite set then $|\mathcal{P}(S)| = 2^{|S|}$.

The idea behind this proof is that the formula works for the empty set (because $\mathcal{P}(\emptyset) = \{\emptyset\}$ so $|\mathcal{P}(\emptyset)| = 1 = 2^0 = 2^{|\emptyset|}$) and we could construct any finite set starting with the empty set and repeatedly adding elements of the desired set.

Now if T is a set such that $|\mathcal{P}(T)| = 2^{|T|}$ and S is set made by adding a new element to T (i.e. $S = T \cup \{a\}$, where $a \notin T$) then S 's subsets will either be subsets of T or subsets of T plus the new element.

The size of the power set, second visit

Lemma

If S is a finite set then $|\mathcal{P}(S)| = 2^{|S|}$.

The idea behind this proof is that the formula works for the empty set (because $\mathcal{P}(\emptyset) = \{\emptyset\}$ so $|\mathcal{P}(\emptyset)| = 1 = 2^0 = 2^{|\emptyset|}$) and we could construct any finite set starting with the empty set and repeatedly adding elements of the desired set.

Now if T is a set such that $|\mathcal{P}(T)| = 2^{|T|}$ and S is set made by adding a new element to T (i.e. $S = T \cup \{a\}$, where $a \notin T$) then S 's subsets will either be subsets of T or subsets of T plus the new element. There are $2^{|T|}$ of the first type (because of the T we chose) and since each of the subsets of the second type are associated with exactly one of T 's subsets there are $2^{|T|}$ of those as well.

The size of the power set, second visit

Lemma

If S is a finite set then $|\mathcal{P}(S)| = 2^{|S|}$.

The idea behind this proof is that the formula works for the empty set (because $\mathcal{P}(\emptyset) = \{\emptyset\}$ so $|\mathcal{P}(\emptyset)| = 1 = 2^0 = 2^{|\emptyset|}$) and we could construct any finite set starting with the empty set and repeatedly adding elements of the desired set.

Now if T is a set such that $|\mathcal{P}(T)| = 2^{|T|}$ and S is set made by adding a new element to T (i.e. $S = T \cup \{a\}$, where $a \notin T$) then S 's subsets will either be subsets of T or subsets of T plus the new element. There are $2^{|T|}$ of the first type (because of the T we chose) and since each of the subsets of the second type are associated with exactly one of T 's subsets there are $2^{|T|}$ of those as well. So S has $2 \cdot 2^{|T|} = 2^{|T|+1} = 2^{|S|}$ subsets.

Strong Induction

This type of induction is sometimes called **weak induction** because there's another type called **strong induction** shown by the following argument:

$$\frac{\begin{array}{l} P(0) \\ (\forall n \in \mathbb{N} - \{0\}) [(\forall m < n) [P(m)] \Rightarrow P(n)] \end{array}}{\therefore (\forall n \in \mathbb{N}) [P(n)]}$$

In other words proving $P(n)$'s truth may depend on all previous propositions being true.

Strong Induction

This type of induction is sometimes called **weak induction** because there's another type called **strong induction** shown by the following argument:

$$\frac{P(0) \quad (\forall n \in \mathbb{N} - \{0\}) [(\forall m < n) [P(m)] \Rightarrow P(n)]}{\therefore (\forall n \in \mathbb{N}) [P(n)]}$$

In other words proving $P(n)$'s truth may depend on all previous propositions being true.

We'll return to strong induction when we get to analyzing recursive algorithms.

The Existential Quantifier, \exists

We use the **existential quantifier** when we want to assert that a predicate is true for at least one element of given (or implied) set.

The Existential Quantifier, \exists

We use the **existential quantifier** when we want to assert that a predicate is true for at least one element of given (or implied) set. So

$$(\exists x \in S)P(x)$$

says that there's at least one element of S for which the predicate is true.

The Existential Quantifier, \exists

We use the **existential quantifier** when we want to assert that a predicate is true for at least one element of given (or implied) set. So

$$(\exists x \in S)P(x)$$

says that there's at least one element of S for which the predicate is true.

For example, $(\exists x \in \mathbb{R})[x^2 = 0]$ and $(\exists x \in \mathbb{R})[x^2 = 1]$ are both true, but $(\exists x \in \mathbb{R})[x^2 = -1]$ is false.

A Sillier Phrase to Translate

Let's consider the claim "Some dogs are poodles".

A Sillier Phrase to Translate

Let's consider the claim "Some dogs are poodles". If we let D be the set of all dogs and set $P(x)$ to mean "x is a poodle" then we could write this as


$$(\exists x \in D)[P(x)].$$

A Sillier Phrase to Translate

Let's consider the claim "Some dogs are poodles". If we let D be the set of all dogs and set $P(x)$ to mean "x is a poodle" then we could write this as

$$(\exists x \in D)[P(x)].$$

Or, if we make $D(x)$ mean "x is a dog" we could translate the claim into



The image shows two handwritten mathematical expressions. The first expression is $(\exists x)[P(x) \Rightarrow D(x)]$, and the second is $(\exists x)[P(x) \wedge D(x)]$. A large red 'X' is drawn over the first expression, indicating it is incorrect.

$$\begin{aligned} &(\exists x)[P(x) \Rightarrow D(x)] \\ &(\exists x)[P(x) \wedge D(x)] \end{aligned}$$

A Sillier Phrase to Translate

Let's consider the claim "Some dogs are poodles". If we let D be the set of all dogs and set $P(x)$ to mean "x is a poodle" then we could write this as

$$(\exists x \in D)[P(x)].$$

Or, if we make $D(x)$ mean "x is a dog" we could translate the claim into

$$(\exists x)[D(x) \wedge P(x)].$$

And More Seriously

Remember that every positive real number has a real square root?

And More Seriously

Remember that every positive real number has a real square root? Let's translate that into math without using the square root symbol.

$$(\forall x > 0)(\exists y \in \mathbb{R})[y \cdot y = x]$$

And More Seriously

Remember that every positive real number has a real square root? Let's translate that into math without using the square root symbol.

$$(\forall x \geq 0)(\exists y \in \mathbb{R})[y \cdot y = x]$$

works if we're doing the perfectly reasonable thing of including some constraints in the quantifier.

And More Seriously

Remember that every positive real number has a real square root? Let's translate that into math without using the square root symbol.

$$(\forall x \geq 0)(\exists y \in \mathbb{R})[y \cdot y = x]$$

works if we're doing the perfectly reasonable thing of including some constraints in the quantifier.

And if you need it in full form:

$$(\forall x \in \mathbb{R})[(x \geq 0) \Rightarrow (\exists y \in \mathbb{R})[y \cdot y = x]].$$

And More Seriously

Remember that every positive real number has a real square root? Let's translate that into math without using the square root symbol.

$$(\forall x \geq 0)(\exists y \in \mathbb{R})[y \cdot y = x]$$

works if we're doing the perfectly reasonable thing of including some constraints in the quantifier.

And if you need it in full form:

$$(\forall x \in \mathbb{R})[(x \geq 0) \Rightarrow (\exists y \in \mathbb{R})[y \cdot y = x]].$$

Or even

$$(\forall x \in \mathbb{R})(\exists y \in \mathbb{R})[(x \geq 0) \Rightarrow [y \cdot y = x]],$$

since when $x < 0$ any y will do.

Important Concepts

- A variable in a predicate logic statement is considered **free** if it is not quantified, that is to say, it doesn't appear in a quantifier earlier in the statement.

Important Concepts

- A variable in a predicate logic statement is considered **free** if it is not quantified, that is to say, it doesn't appear in a quantifier earlier in the statement.
- Otherwise it is said to be **bound**.

Important Concepts

- A variable in a predicate logic statement is considered **free** if it is not quantified, that is to say, it doesn't appear in a quantifier earlier in the statement.
- Otherwise it is said to be **bound**.
- A statement in which all variables are bound is considered a **sentence** and is, in essence, a proposition.

Translation Issues (Again)

As usual, natural human language has enough ambiguity that translating back and forth from logic can be difficult.

Translation Issues (Again)

As usual, natural human language has enough ambiguity that translating back and forth from logic can be difficult. Consider, for example, the following English sentence: “If you can solve any problem we come up with for the exam, then you get an A”.

Translation Issues (Again)

As usual, natural human language has enough ambiguity that translating back and forth from logic can be difficult. Consider, for example, the following English sentence: “If you can solve any problem we come up with for the exam, then you get an A”. The phrase “... you can solve any problem ... on the exam ...” is reasonably interpretable as either a universally or an existentially quantified statement.

Translation Issues (Again)

As usual, natural human language has enough ambiguity that translating back and forth from logic can be difficult. Consider, for example, the following English sentence: “If you can solve any problem we come up with for the exam, then you get an A”. The phrase “... you can solve any problem ... on the exam ...” is reasonably interpretable as either a universally or an existentially quantified statement.

Main take-away: the word “any” can be particularly problematic, especially in the phrase “for any”.

Examples from Math ... and Song

Goldbach's conjecture states that every even integer greater than two is the sum of two primes. Let's rewrite that using quantifiers, letting \mathbb{Z} be the integers, and P the set of primes.

$$(\forall n \in \mathbb{Z}) [(2n > 2) \Rightarrow (\exists p, q \in P) [p + q = 2n]]$$

Examples from Math ... and Song

Goldbach's conjecture states that every even integer greater than two is the sum of two primes. Let's rewrite that using quantifiers, letting \mathbb{Z} be the integers, and P the set of primes.

$$(\forall n \in \mathbb{Z})(\exists p, q \in P)[(2n > 2) \Rightarrow (2n = p + q)].$$

Examples from Math ... and Song

Goldbach's conjecture states that every even integer greater than two is the sum of two primes. Let's rewrite that using quantifiers, letting \mathbb{Z} be the integers, and P the set of primes.

$$(\forall n \in \mathbb{Z})(\exists p, q \in P)[(2n > 2) \Rightarrow (2n = p + q)].$$

How about Dean Martin's song "Everybody Loves Somebody Sometime"? First, what predicates do we need?

Examples from Math ... and Song

Goldbach's conjecture states that every even integer greater than two is the sum of two primes. Let's rewrite that using quantifiers, letting \mathbb{Z} be the integers, and P the set of primes.

$$(\forall n \in \mathbb{Z})(\exists p, q \in P)[(2n > 2) \Rightarrow (2n = p + q)].$$

How about Dean Martin's song "Everybody Loves Somebody Sometime"? First, what predicates do we need?

- $P(x)$, "x is a person"
- $T(t)$, "t is a time"
- $L(x, y, t)$, "x loves y at time t"

$$(\forall x) [P(x) \Rightarrow (\exists y)(\exists t) [P(y) \wedge T(t) \wedge L(x, y, t)]]$$

Examples from Math ... and Song

Goldbach's conjecture states that every even integer greater than two is the sum of two primes. Let's rewrite that using quantifiers, letting \mathbb{Z} be the integers, and P the set of primes.

$$(\forall n \in \mathbb{Z})(\exists p, q \in P)[(2n > 2) \Rightarrow (2n = p + q)].$$

How about Dean Martin's song "Everybody Loves Somebody Sometime"? First, what predicates do we need?

- $P(x)$, "x is a person"
- $T(t)$, "t is a time"
- $L(x, y, t)$, "x loves y at time t"

$$(\forall x)(\exists y)(\exists t)[P(x) \Rightarrow (P(y) \wedge T(t) \wedge L(x, y, t))]$$

$$\text{or } (\forall x)[P(x) \Rightarrow (\exists y)(\exists t)[P(y) \wedge T(t) \wedge L(x, y, t)]]$$