

Welcome!

- CSE 2321: Foundations I: Discrete Structures
- Dr. Charles Estill (He/Him)

Class Conventions

- We will be talking about Boolean functions in class

Class Conventions

- We will be talking about Boolean functions in class (that is to say, true/false functions).

Class Conventions

- We will be talking about Boolean functions in class (that is to say, true/false functions). We will use 1 (one) for true and 0 (zero) for false.

Class Conventions

- We will be talking about Boolean functions in class (that is to say, true/false functions). We will use 1 (one) for true and 0 (zero) for false.
- In the pseudocode that I use, I will mostly be using mathematical notation rather than any specific computer language (e.g., \neq instead of $!=$).

Class Conventions

- We will be talking about Boolean functions in class (that is to say, true/false functions). We will use 1 (one) for true and 0 (zero) for false.
- In the pseudocode that I use, I will mostly be using mathematical notation rather than any specific computer language (e.g., \neq instead of $!=$).
- In the pseudocode, indentation will be used to indicate nested code.

Class Conventions

- We will be talking about Boolean functions in class (that is to say, true/false functions). We will use 1 (one) for true and 0 (zero) for false.
- In the pseudocode that I use, I will mostly be using mathematical notation rather than any specific computer language (e.g., \neq instead of $!=$).
- In the pseudocode, indentation will be used to indicate nested code. For example:

```
FUNCTION Total(n)
```

```
     $x \leftarrow 0$ 
```

```
    FOR  $i \leftarrow 1$  TO  $n$  DO
```

```
         $x \leftarrow x + i$ 
```

```
    RETURN( $x$ )
```

Sigma Summation, Σ

Since we will be using a large amount of summation in this class, let's review the notation:

Sigma Summation, \sum

Since we will be using a large amount of summation in this class, let's review the notation:

$$\sum_{k=a}^b f(k) \text{ or sometimes } \sum_{k=a}^b f_k$$

means to add up all the terms of the form $f(k)$ (or f_k) where k starts at an integer a , goes up by one each time, and finishes at an (other) integer b .

Sigma Summation, \sum

Since we will be using a large amount of summation in this class, let's review the notation:

$$\sum_{k=a}^b f(k) \text{ or sometimes } \sum_{k=a}^b f_k$$

means to add up all the terms of the form $f(k)$ (or f_k) where k starts at an integer a , goes up by one each time, and finishes at an (other) integer b . Or, more explicitly,

$$f(a) + f(a+1) + f(a+2) + \cdots + f(b-2) + f(b-1) + f(b).$$

$$\sum_{k=a}^a f(k) = f(a) \text{ and } \sum_{k=3}^1 2 = 0$$

Propositions

A **proposition** or statement is a sentence that is either true or false. For example:

- $8 + 3 = 10$

Propositions

A **proposition** or statement is a sentence that is either true or false. For example:

- $8 + 3 = 10$
- Any field of characteristic zero contains a copy of the rational numbers.

Propositions

A **proposition** or statement is a sentence that is either true or false. For example:

- $8 + 3 = 10$
- Any field of characteristic zero contains a copy of the rational numbers.
- The earth is 90% cheese.

NOT Propositions

The following are examples of things that aren't propositions:

NOT Propositions

The following are examples of things that aren't propositions:

- Chocolate ice cream is better than lemon sherbet.

NOT Propositions

The following are examples of things that aren't propositions:

- Chocolate ice cream is better than lemon sherbet.
(This is an opinion.)

NOT Propositions

The following are examples of things that aren't propositions:

- Chocolate ice cream is better than lemon sherbet.
(This is an opinion.)
- Tall green ideas swim fiscally.

NOT Propositions

The following are examples of things that aren't propositions:

- Chocolate ice cream is better than lemon sherbet.
(This is an opinion.)
- Tall green ideas swim fiscally.
(There is no reasonable way to interpret this except as poetry.)

NOT Propositions

The following are examples of things that aren't propositions:

- Chocolate ice cream is better than lemon sherbet.
(This is an opinion.)
- Tall green ideas swim fiscally.
(There is no reasonable way to interpret this except as poetry.)
- The whole number x is a perfect square.

NOT Propositions

The following are examples of things that aren't propositions:

- Chocolate ice cream is better than lemon sherbet.
(This is an opinion.)
- Tall green ideas swim fiscally.
(There is no reasonable way to interpret this except as poetry.)
- The whole number x is a perfect square.
(We'll return to this, but the main issue is that the truth of the statement depends on what x is.)

Logical Connectives

We commonly name individual propositions we are to refer to with letters, e.g. P , Q , R , and so on. These are sometimes called **primitive** or **atomic** propositions.

Logical Connectives

We commonly name individual propositions we are to refer to with letters, e.g. P , Q , R , and so on. These are sometimes called **primitive** or **atomic** propositions. But if we then want to be able to talk about things that are more interesting than just the basic truth of each atomic propositions, we need a way to put them together meaningfully.

Logical Connectives

We commonly name individual propositions we are to refer to with letters, e.g. P , Q , R , and so on. These are sometimes called **primitive** or **atomic** propositions. But if we then want to be able to talk about things that are more interesting than just the basic truth of each atomic propositions, we need a way to put them together meaningfully.

The first of these methods are the **logical connectives**. For example, the English word “and” is somewhat equivalent to the logical \wedge , where $P \wedge Q$ is true when both P and Q are true.

Beware Ambiguity

But we have to be careful about precisely what we mean.
Natural language is more subtle than the pure math that we need to use and often is quite ambiguous.

Beware Ambiguity

But we have to be careful about precisely what we mean. Natural language is more subtle than the pure math that we need to use and often is quite ambiguous. The word “or” is a perfect representative of this. In natural language it frequently suggests exclusivity: “Sign up for basic cable and you’ll receive a free month of HBO or Showtime”. But that isn’t the case in mathematical logic (or programming).

Beware Ambiguity

But we have to be careful about precisely what we mean. Natural language is more subtle than the pure math that we need to use and often is quite ambiguous. The word “or” is a perfect representative of this. In natural language it frequently suggests exclusivity: “Sign up for basic cable and you’ll receive a free month of HBO or Showtime”. But that isn’t the case in mathematical logic (or programming).

In addition, in natural language there are multiple words sharing a **denotation** but with different **connotations**.

↑
dictionary
definition

↑
other (unstated)
definition

Beware Ambiguity

But we have to be careful about precisely what we mean. Natural language is more subtle than the pure math that we need to use and often is quite ambiguous. The word “or” is a perfect representative of this. In natural language it frequently suggests exclusivity: “Sign up for basic cable and you’ll receive a free month of HBO or Showtime”. But that isn’t the case in mathematical logic (or programming).

In addition, in natural language there are multiple words sharing a **denotation** but with different **connotations**. E.g., in translating from English to math, the word “BUT” would usually translate to \wedge , but it frequently suggests surprise on the part of the speaker: “Joe Montana threw for five touchdowns but the 49ers lost”.

M

$M \wedge L$

Loss of Meaning

Minor trigger warning: I use being nauseous and its effects in an example here

What happened with the last example is that we lost some of the meaning when we translated into logic. This happens.

Loss of Meaning

Minor trigger warning: I use being nauseous and its effects in an example here

What happened with the last example is that we lost some of the meaning when we translated into logic. This happens. Similarly, “AND” often implies, in English, a time sequence or causality: “I ate at Rally’s and I puked”.

Loss of Meaning

Minor trigger warning: I use being nauseous and its effects in an example here

What happened with the last example is that we lost some of the meaning when we translated into logic. This happens. Similarly, “AND” often implies, in English, a time sequence or causality: “I ate at Rally’s and I puked”. In math \wedge is commutative so that $P \wedge Q$ means the same thing as $Q \wedge P$.

Loss of Meaning

Minor trigger warning: I use being nauseous and its effects in an example here

What happened with the last example is that we lost some of the meaning when we translated into logic. This happens. Similarly, “AND” often implies, in English, a time sequence or causality: “I ate at Rally’s and I puked”. In math \wedge is commutative so that $P \wedge Q$ means the same thing as $Q \wedge P$. But “I puked and I ate at Rally’s” would have a very different meaning.

Loss of Meaning

Minor trigger warning: I use being nauseous and its effects in an example here

What happened with the last example is that we lost some of the meaning when we translated into logic. This happens. Similarly, “AND” often implies, in English, a time sequence or causality: “I ate at Rally’s and I puked”. In math \wedge is commutative so that $P \wedge Q$ means the same thing as $Q \wedge P$. But “I puked and I ate at Rally’s” would have a very different meaning.

Similarly in other natural languages, there will be ambiguity in how one is to interpret the equivalent translations.

Truth Tables

In order to avoid this ambiguity we introduce the main tool of propositional logic: truth tables.

A **truth table** is way of explicitly telling us what the truth value of a **compound proposition** depending on the truth values of each of its atomic propositions (or **propositional variables**).

Truth Tables

In order to avoid this ambiguity we introduce the main tool of propositional logic: truth tables.

A **truth table** is way of explicitly telling us what the truth value of a **compound proposition** depending on the truth values of each of its atomic propositions (or **propositional variables**). For example the truth table for the logical connective \vee , the near-equivalent of the english “OR” would be

P	Q	$P \vee Q$
0	0	0
0	1	1
1	0	1
1	1	1

Note that this is what is sometimes called the **inclusive-or**.

Truth Tables Defined

A truth table for a logical expression is a table which has one column for each propositional variable (input) and one column for the expression as a whole (output). There should be one row for every possible combination of truth values for each of the inputs.

Truth Tables Defined

A truth table for a logical expression is a table which has one column for each propositional variable (input) and one column for the expression as a whole (output). There should be one row for every possible combination of truth values for each of the inputs. (For this class true will be represented by the number one and false by the number zero.)

Truth Tables Defined

A truth table for a logical expression is a table which has one column for each propositional variable (input) and one column for the expression as a whole (output). There should be one row for every possible combination of truth values for each of the inputs. (For this class true will be represented by the number one and false by the number zero.) The value in the output's column is the value of the expression given the variables' truth values.

P	Q	R	$f(P, Q, R)$
0	0	0	
0	0	1	
0	1	0	
0	1	1	
1	0	0	
1	0	1	
1	1	0	
1	1	1	

Truth Tables Defined

A truth table for a logical expression is a table which has one column for each propositional variable (input) and one column for the expression as a whole (output). There should be one row for every possible combination of truth values for each of the inputs. (For this class true will be represented by the number one and false by the number zero.) The value in the output's column is the value of the expression given the variables' truth values.

If there are n inputs, how many rows (not including the header) should there be in the truth table?

Truth Tables Defined

A truth table for a logical expression is a table which has one column for each propositional variable (input) and one column for the expression as a whole (output). There should be one row for every possible combination of truth values for each of the inputs. (For this class true will be represented by the number one and false by the number zero.) The value in the output's column is the value of the expression given the variables' truth values.

If there are n inputs, how many rows (not including the header) should there be in the truth table? 2^n , because there's two possibilities for each of the n variables and a choice for one doesn't affect the choice for the others.

The Commonly Used Connectives

So far we've talked about \wedge and \vee :

P	Q	$P \wedge Q$
0	0	0
0	1	0
1	0	0
1	1	1

P	Q	$P \vee Q$
0	0	0
0	1	1
1	0	1
1	1	1

The Commonly Used Connectives

So far we've talked about \wedge and \vee :

P	Q	$P \wedge Q$
0	0	0
0	1	0
1	0	0
1	1	1

P	Q	$P \vee Q$
0	0	0
0	1	1
1	0	1
1	1	1

The other common logical connectives are \neg (called “not”),

$\sim P$

P	$\neg P$
0	1
1	0

The Commonly Used Connectives

So far we've talked about \wedge and \vee :

P	Q	$P \wedge Q$
0	0	0
0	1	0
1	0	0
1	1	1

P	Q	$P \vee Q$
0	0	0
0	1	1
1	0	1
1	1	1

The other common logical connectives are \neg (called “not”),
the **conditional** \Rightarrow (“ P implies Q ”),

P	$\neg P$
0	1
1	0

P	Q	$P \Rightarrow Q$
0	0	1
0	1	1
1	0	0
1	1	1

The Commonly Used Connectives

So far we've talked about \wedge and \vee :

P	Q	$P \wedge Q$
0	0	0
0	1	0
1	0	0
1	1	1

P	Q	$P \vee Q$
0	0	0
0	1	1
1	0	1
1	1	1

The other common logical connectives are \neg (called “not”), the **conditional** \Rightarrow (“ P implies Q ”), and the **biconditional** \Leftrightarrow , which have the following truth tables:

P	$\neg P$
0	1
1	0

P	Q	$P \Rightarrow Q$
0	0	1
0	1	1
1	0	0
1	1	1

P	Q	$P \Leftrightarrow Q$
0	0	1
0	1	0
1	0	0
1	1	1

$P \Rightarrow Q$
and
 $P \Leftarrow Q$ ($Q \Rightarrow P$)

The Conditional

The conditional is worth special attention. The statement $P \Rightarrow Q$ means that if P is true, then Q is also true.

The Conditional

The conditional is worth special attention. The statement $P \Rightarrow Q$ means that if P is true, then Q is also true. Notice that the only way that it can be false is if P (called the **antecedent**) is true and Q (called the **consequent**) is false.

The Conditional

The conditional is worth special attention. The statement $P \Rightarrow Q$ means that if P is true, then Q is also true. Notice that the only way that it can be false is if P (called the **antecedent**) is true and Q (called the **consequent**) is false.

With some exceptions the English/math translations tend to be as follows:

if P then Q	$P \Rightarrow Q$
P if Q	$P \Leftarrow Q$
P only if Q	$P \Rightarrow Q$ or $Q \Rightarrow P$
P if and only if Q	$P \Leftrightarrow Q$

The Conditional

The conditional is worth special attention. The statement $P \Rightarrow Q$ means that if P is true, then Q is also true. Notice that the only way that it can be false is if P (called the **antecedent**) is true and Q (called the **consequent**) is false.

With some exceptions the English/math translations tend to be as follows:

if P then Q

$P \Rightarrow Q$

P if Q

$P \Leftarrow Q$

P only if Q

$P \Rightarrow Q$ or $Q \Rightarrow P$

P if and only if Q

$P \Leftrightarrow Q$

But not always. The conditional is especially troublesome to translate.

The Converse & the Contrapositive

Given the conditional proposition $P \Rightarrow Q$, there are two associated propositions worth defining:

The Converse & the Contrapositive

Given the conditional proposition $P \Rightarrow Q$, there are two associated propositions worth defining:

- The **converse**: $Q \Rightarrow P$, and

The Converse & the Contrapositive

Given the conditional proposition $P \Rightarrow Q$, there are two associated propositions worth defining:

- The **converse**: $Q \Rightarrow P$, and
- The **contrapositive**: $(\neg Q) \Rightarrow (\neg P)$

The Converse & the Contrapositive

Given the conditional proposition $P \Rightarrow Q$, there are two associated propositions worth defining:

- The **converse**: $Q \Rightarrow P$, and
- The **contrapositive**: $(\neg Q) \Rightarrow (\neg P)$

P : "no shoes"
 Q : "no service"

As an example consider the common sign in restaurants "No shoes, no shirt, no service." interpreted as "If you're not wearing shoes and a shirt then we will not serve you." (Or, simpler: "No shoes implies no service".)

The Converse & the Contrapositive

Given the conditional proposition $P \Rightarrow Q$, there are two associated propositions worth defining:

- The **converse**: $Q \Rightarrow P$, and
- The **contrapositive**: $(\neg Q) \Rightarrow (\neg P)$

P : "no shoes"
 Q : "no service"

As an example consider the common sign in restaurants "No shoes, no shirt, no service." interpreted as "If you're not wearing shoes and a shirt then we will not serve you." (Or, simpler: "No shoes implies no service".) What is its contrapositive?

"If you were served, you were wearing shoes"

The Converse & the Contrapositive

Given the conditional proposition $P \Rightarrow Q$, there are two associated propositions worth defining:

- The **converse**: $Q \Rightarrow P$, and
- The **contrapositive**: $(\neg Q) \Rightarrow (\neg P)$

As an example consider the common sign in restaurants “No shoes, no shirt, no service.” interpreted as “If you’re not wearing shoes and a shirt then we will not serve you.” (Or, simpler: “No shoes implies no service”.) What is its contrapositive? How about the converse?

“no service implies no shoes”

Why is $0 \Rightarrow P$ Always True?

The Agreement (not an actual agreement)

Suppose I tell you “If you study for ten hours a day, you will pass the class.” We can translate this sentence into logic as follows:

$P =$ “You study 10 hours a day.”	$Q =$ “You’ll pass the class”	$P \Rightarrow Q$
0	0	1
0	1	1
1	0	0
1	1	1

Why is $0 \Rightarrow P$ Always True?

The Agreement (not an actual agreement)

Suppose I tell you “If you study for ten hours a day, you will pass the class.” We can translate this sentence into logic as follows:

$P =$ “You study 10 hours a day.”	$Q =$ “You’ll pass the class”	$P \Rightarrow Q$
0	0	1
0	1	1
1	0	0
1	1	1

If we think of the original sentence as an agreement between us, we see that the agreement isn’t violated by the first two rows.

Why is $0 \Rightarrow P$ Always True?

The Contrapositive & Chained Implications

In addition, consider the following truth table:

P	Q	$P \Rightarrow Q$	$\neg Q$	$\neg P$	$(\neg Q) \Rightarrow (\neg P)$
0	0	1			
0	1	1	0		
1	0	0		0	0
1	1	1	0	0	

Why is $0 \Rightarrow P$ Always True?

The Contrapositive & Chained Implications

In addition, consider the following truth table:

P	Q	$P \Rightarrow Q$	$\neg Q$	$\neg P$	$(\neg Q) \Rightarrow (\neg P)$
0	0	1			
0	1	1			
1	0	0			
1	1	1			

Finally, consider that if you knew that $A \Rightarrow B$ and $B \Rightarrow C$, what else might you consider to be true?

Why is $0 \Rightarrow P$ Always True?

The Contrapositive & Chained Implications

In addition, consider the following truth table:

P	Q	$P \Rightarrow Q$	$\neg Q$	$\neg P$	$(\neg Q) \Rightarrow (\neg P)$
0	0	1			
0	1	1			
1	0	0			
1	1	1			

Finally, consider that if you knew that $A \Rightarrow B$ and $B \Rightarrow C$, what else might you consider to be true? That $A \Rightarrow C$ is one pretty reasonable thing.

Why is $0 \Rightarrow P$ Always True?

The Contrapositive & Chained Implications

In addition, consider the following truth table:

P	Q	$P \Rightarrow Q$	$\neg Q$	$\neg P$	$(\neg Q) \Rightarrow (\neg P)$
0	0	1			
0	1	1			
1	0	0			
1	1	1			

Finally, consider that if you knew that $A \Rightarrow B$ and $B \Rightarrow C$, what else might you consider to be true? That $A \Rightarrow C$ is one pretty reasonable thing. Or, in other words, you'd want $((A \Rightarrow B) \wedge (B \Rightarrow C)) \Rightarrow (A \Rightarrow C)$ to be always true.

Why is $0 \Rightarrow P$ Always True?

The Contrapositive & Chained Implications

In addition, consider the following truth table:

P	Q	$P \Rightarrow Q$	$\neg Q$	$\neg P$	$(\neg Q) \Rightarrow (\neg P)$
0	0	1			
0	1	1			
1	0	0			
1	1	1			

Finally, consider that if you knew that $A \Rightarrow B$ and $B \Rightarrow C$, what else might you consider to be true? That $A \Rightarrow C$ is one pretty reasonable thing. Or, in other words, you'd want $((A \Rightarrow B) \wedge (B \Rightarrow C)) \Rightarrow (A \Rightarrow C)$ to be always true. It turns out that the truth table for \Rightarrow is the only choice that makes that work.

Bigger than Atoms

So now that we have some connectives and a tool for investigating Boolean logic we can look at bigger things than individual atomic propositions (like I promised).

Bigger than Atoms

So now that we have some connectives and a tool for investigating Boolean logic we can look at bigger things than individual atomic propositions (like I promised). A **compound proposition** is a well-formed string (WFS) of atomic propositions, connectives, and parentheses. Where a string counts as “well-formed” if it is either

Bigger than Atoms

So now that we have some connectives and a tool for investigating Boolean logic we can look at bigger things than individual atomic propositions (like I promised). A **compound proposition** is a well-formed string (WFS) of atomic propositions, connectives, and parentheses. Where a string counts as “well-formed” if it is either

- 1 an atomic proposition (e.g., P or Q),

Bigger than Atoms

So now that we have some connectives and a tool for investigating Boolean logic we can look at bigger things than individual atomic propositions (like I promised). A **compound proposition** is a well-formed string (WFS) of atomic propositions, connectives, and parentheses. Where a string counts as “well-formed” if it is either

- 1 an atomic proposition (e.g., P or Q or φ),

Bigger than Atoms

So now that we have some connectives and a tool for investigating Boolean logic we can look at bigger things than individual atomic propositions (like I promised). A **compound proposition** is a well-formed string (WFS) of atomic propositions, connectives, and parentheses. Where a string counts as “well-formed” if it is either

- ① an atomic proposition (e.g., P or Q or φ),
- ② of the form $thing1 \wedge thing2$,

Bigger than Atoms

So now that we have some connectives and a tool for investigating Boolean logic we can look at bigger things than individual atomic propositions (like I promised). A **compound proposition** is a well-formed string (WFS) of atomic propositions, connectives, and parentheses. Where a string counts as “well-formed” if it is either

- ① an atomic proposition (e.g., P or Q or φ),
- ② of the form $thing1 \wedge thing2$, $thing1 \vee thing2$,

Bigger than Atoms

So now that we have some connectives and a tool for investigating Boolean logic we can look at bigger things than individual atomic propositions (like I promised). A **compound proposition** is a well-formed string (WFS) of atomic propositions, connectives, and parentheses. Where a string counts as “well-formed” if it is either

- 1 an atomic proposition (e.g., P or Q or φ),
- 2 of the form $thing1 \wedge thing2$, $thing1 \vee thing2$, $thing1 \Rightarrow thing2$,

Bigger than Atoms

So now that we have some connectives and a tool for investigating Boolean logic we can look at bigger things than individual atomic propositions (like I promised). A **compound proposition** is a well-formed string (WFS) of atomic propositions, connectives, and parentheses. Where a string counts as “well-formed” if it is either

- 1 an atomic proposition (e.g., P or Q or φ),
- 2 of the form $thing1 \wedge thing2$, $thing1 \vee thing2$, $thing1 \Rightarrow thing2$, $thing1 \Leftrightarrow thing2$,

Bigger than Atoms

So now that we have some connectives and a tool for investigating Boolean logic we can look at bigger things than individual atomic propositions (like I promised). A **compound proposition** is a well-formed string (WFS) of atomic propositions, connectives, and parentheses. Where a string counts as “well-formed” if it is either

- 1 an atomic proposition (e.g., P or Q or φ),
- 2 of the form $thing1 \wedge thing2$, $thing1 \vee thing2$, $thing1 \Rightarrow thing2$, $thing1 \Leftrightarrow thing2$, or $\neg thing1$;

Bigger than Atoms

So now that we have some connectives and a tool for investigating Boolean logic we can look at bigger things than individual atomic propositions (like I promised). A **compound proposition** is a well-formed string (WFS) of atomic propositions, connectives, and parentheses. Where a string counts as “well-formed” if it is either

- ① an atomic proposition (e.g., P or Q or φ),
- ② of the form $thing1 \wedge thing2$, $thing1 \vee thing2$, $thing1 \Rightarrow thing2$, $thing1 \Leftrightarrow thing2$, or $\neg thing1$; where $thing1$ and $thing2$ are well-formed strings,

Bigger than Atoms

So now that we have some connectives and a tool for investigating Boolean logic we can look at bigger things than individual atomic propositions (like I promised). A **compound proposition** is a well-formed string (WFS) of atomic propositions, connectives, and parentheses. Where a string counts as “well-formed” if it is either

- 1 an atomic proposition (e.g., P or Q or φ),
- 2 of the form $thing1 \wedge thing2$, $thing1 \vee thing2$, $thing1 \Rightarrow thing2$, $thing1 \Leftrightarrow thing2$, or $\neg thing1$; where $thing1$ and $thing2$ are well-formed strings,
- 3 (or similarly constructed strings with any connectives that we may define in the future), or

Bigger than Atoms

So now that we have some connectives and a tool for investigating Boolean logic we can look at bigger things than individual atomic propositions (like I promised). A **compound proposition** is a well-formed string (WFS) of atomic propositions, connectives, and parentheses. Where a string counts as “well-formed” if it is either

- ① an atomic proposition (e.g., P or Q or φ),
- ② of the form $thing1 \wedge thing2$, $thing1 \vee thing2$, $thing1 \Rightarrow thing2$, $thing1 \Leftrightarrow thing2$, or $\neg thing1$; where $thing1$ and $thing2$ are well-formed strings,
- ③ (or similarly constructed strings with any connectives that we may define in the future), or
- ④ of the form $(thing1)$ where $thing1$ is a WFS.

An Example

Consider the example

$$((Q \Rightarrow P) \wedge (((\neg P) \vee Q) \wedge (R \Rightarrow R))) \Leftrightarrow P.$$

An Example

Consider the example

$$((Q \Rightarrow P) \wedge (((\neg P) \vee Q) \wedge (R \Rightarrow R))) \Leftrightarrow P.$$

			A	B	C	D	
P	Q	R	$Q \Rightarrow P$	$((\neg P) \vee Q)$	$R \Rightarrow R$	$A \wedge (B \wedge C)$	$D \Leftrightarrow P$
0	0	0	1	1	1	1	0
0	0	1	1	1	1	1	0
0	1	0	0	1	1	0	1
0	1	1	0	1	1	0	1
1	0	0	1	0	1	0	0
1	0	1	1	0	1	0	0
1	1	0	1	1	1	1	1
1	1	1	1	1	1	1	1