

The Satisfiability Problem

(This slide through to “NP-complete” is optional and not covered in lecture or tested.)

While propositional logic is pretty straightforward, that doesn't mean that there's nothing but triviality within. For example, consider the satisfiability problem. Given a proposition in conjunctive normal form, determine whether or not there is a choice of truth values for the atomics that makes the proposition true (i.e. **satisfied**).

The Satisfiability Problem

(This slide through to “NP-complete” is optional and not covered in lecture or tested.)

While propositional logic is pretty straightforward, that doesn't mean that there's nothing but triviality within. For example, consider the satisfiability problem. Given a proposition in conjunctive normal form, determine whether or not there is a choice of truth values for the atomics that makes the proposition true (i.e. **satisfied**).

For example consider the proposition

$$(P \vee Q \vee R) \wedge (\neg P \vee \neg Q) \wedge (\neg P \vee \neg R) \wedge (\neg R \vee \neg Q).$$

Is it satisfiable?

P vs. NP

The satisfiability problem is probably the biggest open problem in computer science and is part of the whole “P versus NP” question. A problem that an algorithm can give an answer to in polynomial time is considered to be in “class P”. A problem whose answer can be **verified** in polynomial time is in “class NP”. So, for example, a Sudoku puzzle is in class NP.

P vs. NP

The satisfiability problem is probably the biggest open problem in computer science and is part of the whole “P versus NP” question. A problem that an algorithm can give an answer to in polynomial time is considered to be in “class P”. A problem whose answer can be **verified** in polynomial time is in “class NP”. So, for example, a Sudoku puzzle is in class NP.

The big question is: Are the two classes the same? Is there a polynomial solution **finding** algorithm for every problem which has a polynomial time **verifying** algorithm?

NP-complete

One idea that has been used to attack the P vs. NP problem is the idea of **NP-completeness**. An NP problem is NP-complete if any other NP-class problem can be modeled using the first type of problem (said model made in polynomial time). The satisfiability problem is NP-complete.

NP-complete

One idea that has been used to attack the P vs. NP problem is the idea of **NP-completeness**. An NP problem is NP-complete if any other NP-class problem can be modeled using the first type of problem (said model made in polynomial time). The satisfiability problem is NP-complete.

Interestingly, so is minesweeper. (See 'Minesweeper is NP-Complete.pdf' in the 'Extra Resources / Further Reading' module on Carmen.)

Sets

A **set** is a collection of distinct objects, each of which is called an **element** of the set. We tend to write sets within curly braces: {
... }.

Sets

A **set** is a collection of distinct objects, each of which is called an **element** of the set. We tend to write sets within curly braces: $\{ \dots \}$. Some examples:

- $\mathbb{N} = \{0, 1, 2, \dots\}$, the set of natural numbers,

Sets

A **set** is a collection of distinct objects, each of which is called an **element** of the set. We tend to write sets within curly braces: $\{ \dots \}$. Some examples:

- $\mathbb{N} = \{0, 1, 2, \dots\}$, the set of natural numbers,
- $\mathbb{Z} = \{0, \pm 1, \pm 2, \dots\}$, the set of integers,

Sets

A **set** is a collection of distinct objects, each of which is called an **element** of the set. We tend to write sets within curly braces: $\{ \dots \}$. Some examples:

- $\mathbb{N} = \{0, 1, 2, \dots\}$, the set of natural numbers,
- $\mathbb{Z} = \{0, \pm 1, \pm 2, \dots\}$, the set of integers,
- \mathbb{Q} , the set of rational numbers (a.k.a. fractions),
- \mathbb{R} , the set of real numbers,
- \mathbb{C} the set of complex numbers

Sets

A **set** is a collection of distinct objects, each of which is called an **element** of the set. We tend to write sets within curly braces: $\{ \dots \}$. Some examples:

- $\mathbb{N} = \{0, 1, 2, \dots\}$, the set of natural numbers,
- $\mathbb{Z} = \{0, \pm 1, \pm 2, \dots\}$, the set of integers,
- \mathbb{Q} , the set of rational numbers (a.k.a. fractions),
- \mathbb{R} , the set of real numbers,
- \mathbb{C} the set of complex numbers
- $\{ \textit{The Big Lebowski}, \textit{Six-String Samurai}, \textit{12 Monkeys}, \textit{Tangled} \}$

Sets

A **set** is a collection of distinct objects, each of which is called an **element** of the set. We tend to write sets within curly braces: $\{ \dots \}$. Some examples:

- $\mathbb{N} = \{0, 1, 2, \dots\}$, the set of natural numbers,
- $\mathbb{Z} = \{0, \pm 1, \pm 2, \dots\}$, the set of integers,
- \mathbb{Q} , the set of rational numbers (a.k.a. fractions),
- \mathbb{R} , the set of real numbers,
- \mathbb{C} the set of complex numbers
- $\{ \textit{The Big Lebowski}, \textit{Six-String Samurai}, \textit{12 Monkeys}, \textit{Tangled} \}$
- $\{ \{3, 10\}, \{2, 3, 17\}, \{ \frac{5k-1}{2} \mid k \in \mathbb{N} \} \}$

Defining Sets With a Rule

That last set in the last set, $\{\frac{5k-1}{2} \mid k \in \mathbb{N}\}$ or $\{\frac{5k-1}{2} \in \mathbb{R} \mid k \in \mathbb{N}\}$, was a bit different.

Defining Sets With a Rule

That last set in the last set, $\{\frac{5k-1}{2} \mid k \in \mathbb{N}\}$ or $\{\frac{5k-1}{2} \in \mathbb{R} \mid k \in \mathbb{N}\}$, was a bit different. What this means is that any fraction of the form $\frac{5k-1}{2}$ for any natural number k is in the set.

Defining Sets With a Rule

That last set in the last set, $\{\frac{5k-1}{2} \mid k \in \mathbb{N}\}$ or $\{\frac{5k-1}{2} \in \mathbb{R} \mid k \in \mathbb{N}\}$, was a bit different. What this means is that any fraction of the form $\frac{5k-1}{2}$ for any natural number k is in the set. The set could also be written as $\{-\frac{1}{2}, 2, \frac{9}{2}, 7, \frac{19}{2}, \dots\}$, but that would be a lot less clear.

Defining Sets With a Rule

That last set in the last set, $\{\frac{5k-1}{2} \mid k \in \mathbb{N}\}$ or $\{\frac{5k-1}{2} \in \mathbb{R} \mid k \in \mathbb{N}\}$, was a bit different. What this means is that any fraction of the form $\frac{5k-1}{2}$ for any natural number k is in the set. The set could also be written as $\{-\frac{1}{2}, 2, \frac{9}{2}, 7, \frac{19}{2}, \dots\}$, but that would be a lot less clear. The vertical line (which is also sometimes a colon, $\{\frac{5k-1}{2} : k \in \mathbb{N}\}$) can be read as “such that”.

Set Notation

- The elements of a set are required to be distinct, but the order of the elements is not important. For example $\{x, y\} = \{y, x\}$, and if $x = y$ then it is also the same as $\{x\}$.

A set with just one element is called a "singleton set".

Set Notation

- The elements of a set are required to be distinct, but the order of the elements is not important. For example $\{x, y\} = \{y, x\}$, and if $x = y$ then it is also the same as $\{x\}$.
I.e. $\{1, 2, 1, 2, 3\} = \{1, 2, 3\}$.

Set Notation

- The elements of a set are required to be distinct, but the order of the elements is not important. For example $\{x, y\} = \{y, x\}$, and if $x = y$ then it is also the same as $\{x\}$.
I.e. $\{1, 2, 1, 2, 3\} = \{1, 2, 3\}$.
- Repetition does not happen in sets. An object is either an element of a set or not; the set can't have multiple copies of an object.

Set Notation

- The elements of a set are required to be distinct, but the order of the elements is not important. For example $\{x, y\} = \{y, x\}$, and if $x = y$ then it is also the same as $\{x\}$.
I.e. $\{1, 2, 1, 2, 3\} = \{1, 2, 3\}$.
- Repetition does not happen in sets. An object is either an element of a set or not; the set can't have multiple copies of an object.
- The expression " $e \in S$ " states that e is an element of S .

Set Notation

- The elements of a set are required to be distinct, but the order of the elements is not important. For example $\{x, y\} = \{y, x\}$, and if $x = y$ then it is also the same as $\{x\}$.
I.e. $\{1, 2, 1, 2, 3\} = \{1, 2, 3\}$.
- Repetition does not happen in sets. An object is either an element of a set or not; the set can't have multiple copies of an object.
- The expression " $e \in S$ " states that e is an element of S .
- Two sets are equal if they contain exactly the same elements.

Set Notation

- The elements of a set are required to be distinct, but the order of the elements is not important. For example $\{x, y\} = \{y, x\}$, and if $x = y$ then it is also the same as $\{x\}$.
I.e. $\{1, 2, 1, 2, 3\} = \{1, 2, 3\}$.
- Repetition does not happen in sets. An object is either an element of a set or not; the set can't have multiple copies of an object.
- The expression " $e \in S$ " states that e is an element of S .
- Two sets are equal if they contain exactly the same elements.
- The set containing no elements is called the empty set and is written \emptyset or sometimes just $\{ \}$.

Set Notation

- The elements of a set are required to be distinct, but the order of the elements is not important. For example $\{x, y\} = \{y, x\}$, and if $x = y$ then it is also the same as $\{x\}$.
I.e. $\{1, 2, 1, 2, 3\} = \{1, 2, 3\}$.
- Repetition does not happen in sets. An object is either an element of a set or not; the set can't have multiple copies of an object.
- The expression " $e \in S$ " states that e is an element of S .
- Two sets are equal if they contain exactly the same elements.
- The set containing no elements is called the empty set and is written \emptyset or sometimes just $\{ \}$.
- The **cardinality** of a set is the number of elements it contains and is written $|S|$.

Comparing Sets

- The expression “ $S \subseteq T$ ” indicates that the set S is a **subset** of the set T . This means that every element of S is an element of T (i.e., $(e \in S) \Rightarrow (e \in T)$).

Comparing Sets

- The expression “ $S \subseteq T$ ” indicates that the set S is a **subset** of the set T . This means that every element of S is an element of T (i.e., $(e \in S) \Rightarrow (e \in T)$). For example,

$$\mathbb{N} \subseteq \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R} \subseteq \mathbb{C}.$$

Comparing Sets

- The expression “ $S \subseteq T$ ” indicates that the set S is a **subset** of the set T . This means that every element of S is an element of T (i.e., $(e \in S) \Rightarrow (e \in T)$). For example,

$$\mathbb{N} \subseteq \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R} \subseteq \mathbb{C}.$$

Note that $\emptyset \subseteq S$ for any set S (and also $S \subseteq S$).

Comparing Sets

- The expression " $S \subseteq T$ " indicates that the set S is a **subset** of the set T . This means that every element of S is an element of T (i.e., $(e \in S) \Rightarrow (e \in T)$). For example,

$$\mathbb{N} \subseteq \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R} \subseteq \mathbb{C}.$$

Note that $\emptyset \subseteq S$ for any set S (and also $S \subseteq S$).

- The expression $S \subset T$ indicates that $S \subseteq T$, but that $S \neq T$.

" S is a proper subset of T "

Comparing Sets

- The expression “ $S \subseteq T$ ” indicates that the set S is a **subset** of the set T . This means that every element of S is an element of T (i.e., $(e \in S) \Rightarrow (e \in T)$). For example,

$$\mathbb{N} \subseteq \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R} \subseteq \mathbb{C}.$$

Note that $\emptyset \subseteq S$ for any set S (and also $S \subseteq S$).

- The expression $S \subset T$ indicates that $S \subseteq T$, but that $S \neq T$. (In older sources \subset was frequently used in the way that we use \subseteq today, however.)

Comparing Sets

- The expression “ $S \subseteq T$ ” indicates that the set S is a **subset** of the set T . This means that every element of S is an element of T (i.e., $(e \in S) \Rightarrow (e \in T)$). For example,

$$\mathbb{N} \subseteq \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R} \subseteq \mathbb{C}.$$

Note that $\emptyset \subseteq S$ for any set S (and also $S \subseteq S$).

- The expression $S \subset T$ indicates that $S \subseteq T$, but that $S \neq T$. (In older sources \subset was frequently used in the way that we use \subseteq today, however.)
- As you might expect $S = T$ means that $S \subseteq T$ and $T \subseteq S$.

Combining Sets

- The **union** of two sets A and B (denoted $A \cup B$) is the set that contains any element that is in either set. E.g., if $A = \{1, 2, 4\}$ and $B = \{3, 4\}$ then $A \cup B = \{1, 2, 3, 4\}$.

Combining Sets

- The **union** of two sets A and B (denoted $A \cup B$) is the set that contains any element that is in either set. E.g., if $A = \{1, 2, 4\}$ and $B = \{3, 4\}$ then $A \cup B = \{1, 2, 3, 4\}$.
"In A OR in B"
- The **intersection** of A and B (denoted $A \cap B$) is the set containing all elements that are in both A and B . E.g., With A and B as above, $A \cap B = \{4\}$.
"In A AND in B"

Combining Sets

- The **union** of two sets A and B (denoted $A \cup B$) is the set that contains any element that is in either set. E.g., if $A = \{1, 2, 4\}$ and $B = \{3, 4\}$ then $A \cup B = \{1, 2, 3, 4\}$.
- The **intersection** of A and B (denoted $A \cap B$) is the set containing all elements that are in both A and B . E.g., With A and B as above, $A \cap B = \{4\}$.
- The **difference** of A and B (denoted $A - B$ or $A \setminus B$) consists of all elements that are in A , but not in B . For our sets,
 $A - B = \{1, 2\}$. $B - A = \{3\}$

Combining Sets

- The **union** of two sets A and B (denoted $A \cup B$) is the set that contains any element that is in either set. E.g., if $A = \{1, 2, 4\}$ and $B = \{3, 4\}$ then $A \cup B = \{1, 2, 3, 4\}$.
- The **intersection** of A and B (denoted $A \cap B$) is the set containing all elements that are in both A and B . E.g., With A and B as above, $A \cap B = \{4\}$.
- The **difference** of A and B (denoted $A - B$ or $A \setminus B$) consists of all elements that are in A , but not in B . For our sets, $A - B = \{1, 2\}$.
 $\mathbb{R}^2 = \mathbb{R} \times \mathbb{R}$
- The **Cartesian product** of A and B (denoted $A \times B$) is the set of ordered pairs $\{(a, b) | (a \in A) \wedge (b \in B)\}$. For our example, $A \times B = \{(1, 3), (1, 4), (2, 3), (2, 4), (4, 3), (4, 4)\}$. $|C \times D| = |C| \cdot |D|$

The Power Set

The **power set** of a set S (denoted $\mathcal{P}(S)$, or $\text{POW}(S)$) is the set of all subsets of S . I.e., $\mathcal{P}(S) = \{T \mid T \subseteq S\}$.

The Power Set

The **power set** of a set S (denoted $\mathcal{P}(S)$, or $\text{POW}(S)$) is the set of all subsets of S . I.e., $\mathcal{P}(S) = \{T \mid T \subseteq S\}$.

Consider $A = \{1, 2, 4\}$ from before. What is $|\mathcal{P}(A)|$?

$$\mathcal{P}(A) = \{\emptyset, \{1\}, \{1, 2\}, \{2, 4\}, A, \{1, 4\}, \{2\}, \{4\}\}$$

The Power Set

The **power set** of a set S (denoted $\mathcal{P}(S)$, or $\text{POW}(S)$) is the set of all subsets of S . I.e., $\mathcal{P}(S) = \{T \mid T \subseteq S\}$.

Consider $A = \{1, 2, 4\}$ from before. What is $|\mathcal{P}(A)|$?

$$|\mathcal{P}(A)| = |\{\emptyset, \{1\}, \{2\}, \{4\}, \{1, 2\}, \{1, 4\}, \{2, 4\}, A\}| = 8.$$

The Power Set

The **power set** of a set S (denoted $\mathcal{P}(S)$, or $\text{POW}(S)$) is the set of all subsets of S . I.e., $\mathcal{P}(S) = \{T \mid T \subseteq S\}$.

Consider $A = \{1, 2, 4\}$ from before. What is $|\mathcal{P}(A)|$?

$$|\mathcal{P}(A)| = |\{\emptyset, \{1\}, \{2\}, \{4\}, \{1, 2\}, \{1, 4\}, \{2, 4\}, A\}| = 8.$$

Notice that A has three elements and $\mathcal{P}(A)$ has eight. If $|B| = 10$, what do you think the cardinality (size) of $\mathcal{P}(B)$ is?

$$2^{10}$$

The size of the power set

$$\mathcal{P}(S) = 2^S$$

Lemma

If S is a finite set then $|\mathcal{P}(S)| = 2^{|S|}$.

The size of the power set

Lemma

If S is a finite set then $|\mathcal{P}(S)| = 2^{|S|}$.

Our first proof of this theorem is to notice that each subset of S , T

The size of the power set

Lemma

If S is a finite set then $|\mathcal{P}(S)| = 2^{|S|}$.

Our first proof of this theorem is to notice that each subset of S , T (i.e., each element of the power set),

The size of the power set

Lemma

If S is a finite set then $|\mathcal{P}(S)| = 2^{|S|}$.

Our first proof of this theorem is to notice that each subset of S , T (i.e., each element of the power set), can be distinguished from other subsets by which particular elements are in T .

The size of the power set

Lemma

If S is a finite set then $|\mathcal{P}(S)| = 2^{|S|}$.

Our first proof of this theorem is to notice that each subset of S , T (i.e., each element of the power set), can be distinguished from other subsets by which particular elements are in T . But this means that we can describe T by answering a question for each element of S : “Is this element in T or not?”

The size of the power set

Lemma

If S is a finite set then $|\mathcal{P}(S)| = 2^{|S|}$.

Our first proof of this theorem is to notice that each subset of S , T (i.e., each element of the power set), can be distinguished from other subsets by which particular elements are in T . But this means that we can describe T by answering a question for each element of S : “Is this element in T or not?”

If $|S| = n$ that's n questions each with one of two answers,

The size of the power set

Lemma

If S is a finite set then $|\mathcal{P}(S)| = 2^{|S|}$.

Our first proof of this theorem is to notice that each subset of S , T (i.e., each element of the power set), can be distinguished from other subsets by which particular elements are in T . But this means that we can describe T by answering a question for each element of S : “Is this element in T or not?”

If $|S| = n$ that's n questions each with one of two answers, so 2^n ways of answering all the questions, or 2^n subsets.

2^S is the set of functions from S to $2 = \{0, 1\}$

Venn Diagrams

A **Venn diagram** is a pictorial representation of **all** possible intersections between a finite number of different sets. They represents the sets as regions inside closed curves (for example, the interior of a circle) and elements as points in a plane. For example, if I wanted to confirm (*not prove*) that

$$(A \cup B) - C = (A - C) \cup (B - C)$$

then I could draw the picture for both sides and confirm that they are the same.

$$(A \cup B) - C = (A - C) \cup (B - C)$$




Figure: $A \cup B$




Figure: $A - C$




Figure: $B - C$

$$(A \cup B) - C = (A - C) \cup (B - C)$$




Figure: $A \cup B$




Figure: $A - C$




Figure: $B - C$




Figure: $(A \cup B) - C$







Figure: $(A - C) \cup (B - C)$


$$(A \cup B) - C$$

$$\stackrel{?}{=} (A - C) \cup B$$



$$(A \cup B)$$


$$(A - C)$$


$$(A \cup B) - C$$


$$(A - C) \cup B$$

$$A = \{2, 3, 4\}, B = \{4, 5\}, C = \{3, 4, 5, 6\}$$

$$(A \cup B) - C = \{2, 3, 4, 5\} - C = \{2\}$$

$$(A - C) \cup B = \{2\} \cup B = \{2, 4, 5\}$$

Sequences

A **sequence** is an ordered list of objects. You can think of it as a function from the natural numbers into the set the objects are from.

Sequences

A **sequence** is an ordered list of objects. You can think of it as a function from the natural numbers into the set the objects are from.

E.g.,

- $(a_n)_{n=0}^{\infty} = (3, 1, 4, 1, 5, 9, \dots)$

Sequences

A **sequence** is an ordered list of objects. You can think of it as a function from the natural numbers into the set the objects are from.

E.g.,

- $(a_n)_{n=0}^{\infty} = (3, 1, 4, 1, 5, 9, \dots)$
- $(b_n)_{n=1}^{\infty} = (\cos(n\pi))_{n=1}^{\infty} = (-1, 1, -1, 1, -1, \dots)$

Sequences

A **sequence** is an ordered list of objects. You can think of it as a function from the natural numbers into the set the objects are from.

E.g.,

- $(a_n)_{n=0}^{\infty} = (3, 1, 4, 1, 5, 9, \dots)$
- $(b_n)_{n=1}^{\infty} = (\cos(n\pi))_{n=1}^{\infty} = (-1, 1, -1, 1, -1, \dots)$
- $(f_n) = (1, 1, 2, 3, 5, 8, 13, \dots)$

Predicates

A **predicate** is like a proposition whose truth depends on the value of one or more variables.

Predicates

A **predicate** is like a proposition whose truth depends on the value of one or more variables. For example, “ n is a perfect square” is a predicate whose truth value depends on n ’s value. It is true for $n = 4$ or $n = 9$, but false for $n = 5$ or $n = \text{Frank}$.

Predicates

A **predicate** is like a proposition whose truth depends on the value of one or more variables. For example, “ n is a perfect square” is a predicate whose truth value depends on n ’s value. It is true for $n = 4$ or $n = 9$, but false for $n = 5$ or $n = \text{Frank}$.

Like propositions, predicates are generally named with a letter, but because of the dependence on variables, a function-like notation is used. For example, naming the above predicate P tells us that $P(4)$ and $P(9)$ are true but $P(5)$ and $P(3.14159)$ are false.

Predicates

A **predicate** is like a proposition whose truth depends on the value of one or more variables. For example, “ n is a perfect square” is a predicate whose truth value depends on n ’s value. It is true for $n = 4$ or $n = 9$, but false for $n = 5$ or $n = \text{Frank}$.

Like propositions, predicates are generally named with a letter, but because of the dependence on variables, a function-like notation is used. For example, naming the above predicate P tells us that $P(4)$ and $P(9)$ are true but $P(5)$ and $P(3.14159)$ are false.

In other words, predicates are simply functions that return Boolean values.

Quantifying Predicates

Having predicates by themselves is no real improvement on propositional logic. We need a way to be able to talk about multiple propositions (even an infinite number of them) at the same time. For this we have quantifiers

Quantifying Predicates

Having predicates by themselves is no real improvement on propositional logic. We need a way to be able to talk about multiple propositions (even an infinite number of them) at the same time. For this we have quantifiers

There are two common quantifiers: \forall and \exists .

The Universal Quantifier \forall

We use the **universal quantifier** when we want to assert that a predicate is true for every element in a given set. So

$$(\forall x \in S)[P(x)]$$

says that $P(x)$ is true for every x in the set S .

The Universal Quantifier \forall

We use the **universal quantifier** when we want to assert that a predicate is true for every element in a given set. So

$$(\forall x \in S)[P(x)]$$

says that $P(x)$ is true for every x in the set S .

For example $(\forall x \in \mathbb{R})[x^2 \geq 0]$ is a true sentence, but $(\forall x \in \mathbb{R})[x^2 \geq 42]$ is false.

The Universal Quantifier \forall

We use the **universal quantifier** when we want to assert that a predicate is true for every element in a given set. So

$$(\forall x \in S)[P(x)]$$

says that $P(x)$ is true for every x in the set S .

For example $(\forall x \in \mathbb{R})[x^2 \geq 0]$ is a true sentence, but $(\forall x \in \mathbb{R})[x^2 \geq 42]$ is false.

If the set is obvious (or the set of everything) then the set can be omitted and at times we can include extra information in the quantifier.

The Universal Quantifier \forall

We use the **universal quantifier** when we want to assert that a predicate is true for every element in a given set. So

$$(\forall x \in S)[P(x)]$$

says that $P(x)$ is true for every x in the set S .

For example $(\forall x \in \mathbb{R})[x^2 \geq 0]$ is a true sentence, but $(\forall x \in \mathbb{R})[x^2 \geq 42]$ is false.

If the set is obvious (or the set of everything) then the set can be omitted and at times we can include extra information in the quantifier. E.g., in $(\forall x > 0)[e^{\ln x} = x]$ the fact that $x \in \mathbb{R}$ is implied and

The Universal Quantifier \forall

We use the **universal quantifier** when we want to assert that a predicate is true for every element in a given set. So

$$(\forall x \in S)[P(x)]$$

says that $P(x)$ is true for every x in the set S .

For example $(\forall x \in \mathbb{R})[x^2 \geq 0]$ is a true sentence, but $(\forall x \in \mathbb{R})[x^2 \geq 42]$ is false.

If the set is obvious (or the set of everything) then the set can be omitted and at times we can include extra information in the quantifier. E.g., in $(\forall x > 0)[e^{\ln x} = x]$ the fact that $x \in \mathbb{R}$ is implied and the inequality in the quantifier is a short way of saying

$$(\forall x \in \mathbb{R})[(x > 0) \Rightarrow (e^{\ln x} = x)].$$