


MAE 11.2


HW 2

**TRIET**



1. Calculate theoretical (ideal) flame temperature for methane in stoichiometric ratio with enriched air (50% O<sub>2</sub>, 50% N<sub>2</sub> by volume). Pressure is constant at 15 atm and the initial temperature is 298 K.

Stoichiometric reaction:



Mole number ( $n$ ), heat of formation ( $h_{f,m}$ ) and specific heats ( $c_{p,m}$ )

	$n$	$h_{f,m} [\text{J/mol}]$	$c_{p,m} [\text{J/mol}\cdot\text{K}]$
CH <sub>4</sub>	1	-74 831	N/A
O <sub>2</sub>	2	0	N/A
N <sub>2</sub>	2	0	34.805
CO <sub>2</sub>	1	-393 546	58.292
H <sub>2</sub> O	2	-241 845	47.103
N <sub>2</sub>	2	0	34.805

Energy balance:

$$\sum_{\text{Reactants}} n \left( h_{f,m} + \int_{T_{\text{ref}}}^{T_i} c_{p,m} dT \right) = \sum_{\text{Products}} n \left( h_{f,m} + \int_{T_{\text{ref}}}^{T_{\text{final}}} c_{p,m} dT \right)$$

$$n_{\text{CH}_4} (h_{f,\text{CH}_4} + 0) + 0 + 0 = n_{\text{CO}_2} (h_{f,\text{CO}_2} + c_{p,\text{CO}_2} \Delta T) + n_{\text{H}_2\text{O}} (h_{f,\text{H}_2\text{O}} + c_{p,\text{H}_2\text{O}} \Delta T) + n_{\text{N}_2} (0 + c_{p,\text{N}_2} \Delta T)$$

$$n_{\text{CH}_4} \times h_{f,\text{CH}_4} = n_{\text{CO}_2} h_{f,\text{CO}_2} + n_{\text{H}_2\text{O}} h_{f,\text{H}_2\text{O}} + \Delta T (n_{\text{O}_2} c_{p,\text{O}_2} + n_{\text{H}_2\text{O}} c_{p,\text{H}_2\text{O}} + n_{\text{N}_2} c_{p,\text{N}_2})$$

$$\Delta T = \frac{n_{\text{CH}_4} \times h_{f,\text{CH}_4} - (n_{\text{CO}_2} h_{f,\text{CO}_2} + n_{\text{H}_2\text{O}} h_{f,\text{H}_2\text{O}})}{n_{\text{O}_2} c_{p,\text{O}_2} + n_{\text{H}_2\text{O}} c_{p,\text{H}_2\text{O}} + n_{\text{N}_2} c_{p,\text{N}_2}}$$



$$T_g - 298 = \frac{1 \times (-74.831) - [1 \times (-393546) + 2 \times (-241845)]}{1 \times 58.292 + 2 \times 47.103 + 2 \times 34.805}$$

$$T_g = 3910.68 \text{ [K]}$$

3. (a) Calculate AF at stoichiometric condition ( $AF_{st}$ ) for ethyl alcohol  $C_2H_5OH$  (aka ethanol) initially at  $550^\circ R$  burning in air at 20 atmospheres of pressure. AF is the ratio of mass flow of air to mass flow of fuel. Also, calculate  $FA = 1/AF$  for the same condition.
- (b) Calculate AF and  $\Phi = AF_{st} / AF = FA / FA_{st}$  for ethyl alcohol and 50% excess air at the same conditions.

Molecular weight of single element:

$$W_C = 12.011 \left[ \frac{g}{mol} \right] \quad W_H = 1 \left[ \frac{g}{mol} \right] \quad W_O = 16 \left[ \frac{g}{mol} \right] \quad W_N = 14 \left[ \frac{g}{mol} \right]$$




(a) Air to fuel ratio:

$$A/F = \frac{m_{Air}}{m_{Fuel}} = \frac{3 \times (2 \times 16 + 3.76 \times 2 \times 14)}{1 \times (2 \times 12.01 + 5 + 16 + 1)} = \boxed{8.9488}$$

Fuel to air ratio:

$$F/A = 1 \div A/F = 1 \div 8.9488 = \boxed{0.1117}$$

(b) Air mole:  $1.5 n_{air} = 1.5 \times 3 = 4.5$



Air to fuel ratio:

$$A/F = \frac{m_{Air}}{m_{Fuel}} = \frac{4.5 \times (2 \times 16 + 3.76 \times 2 \times 14)}{1 \times (2 \times 12.01 + 5 + 16 + 1)} = \boxed{13.4231}$$

Fuel to air ratio:



$$F/A = 1 \div A/F = 1 \div 13.4231 = \boxed{0.0745}$$

Equivalent ratio:

$$\Phi = \frac{A/F \text{ stoichiometric}}{A/F \text{ at 50\% excess air}} = \frac{8.9488}{13.4231} = \boxed{0.6667}$$

4. (a) Establish the equations which can be employed for the calculation of the equilibrium composition and the flame temperature when one mole of propane  $C_3H_8$  burns adiabatically at a constant pressure of ten atmospheres. The mixture is lean with 75% excess air. Both air and fuel enter at a temperature of 800°R. Consider the products to be  $CO_2$ ,  $CO$ ,  $H_2O$ ,  $H_2$ ,  $O_2$ , and  $N_2$  only. Write all the required equations with known quantities and parameters substituted into the equation. Identify the unknowns. Propane is gaseous at room temperature. Explain what would be different in the analysis if propane entered at a lower temperature in liquid form.
- (b) Use the computer software to calculate the final flame temperature and concentrations of the products with the gaseous propane fuel.
- (c) For the adiabatic situation with gaseous fuel described in Part a, establish the equations to solve for the theoretical (ideal) temperature and composition. What are the products in this case? Again, write the necessary equations, identify the known quantities, and identify the unknowns. Solve the equations for the final temperature and composition. Which of the two temperatures from 2b and 2c is larger? Why?

(a) Stoichiometric reaction:




$$C: 3 = a + c$$

$$H: 8 = 2b + 2d$$


$$O: 17.5 = 2a + b + c + 2e$$

N: Already balanced

Two dissociation:



$$K_1 = \frac{X_{CO} X_{O_2}^{1/2}}{X_{CO_2}} = P^{-1/2} K_{p1}(T_f)$$





$$K_2 = \frac{X_{H_2} X_{O_2}^{1/2}}{X_{H_2O}} = P^{-1/2} K_{p2}(T_f)$$

Energy conservation:

$$\sum_{\text{reactants}} n_m \left( h_{f,m} + \int_{T_{\text{ref}}}^{T_i} c_{p,m} dT \right) = \sum_{\text{products}} n_m \left( h_{f,m} + \int_{T_{\text{ref}}}^{T_f} c_{p,m} dT \right)$$

③ Stoichiometric reaction:



From textbook / internet:

	$h_{f,m} [\text{J/mol}]$	$c_{p,m} [\text{J/mol} \cdot \text{K}]$
$\text{C}_3\text{H}_8$	-103847	73.920
$\text{O}_2$	0	34.88
$\text{N}_2$	0	32.676
$\text{H}_2\text{O}$	-241845	41.103
$\text{CO}_2$	-393546	54.299
$\text{CO}$	-110500	28.56
$\text{H}_2$	0	14.5
$\text{O}_2$	0	34.88
$\text{N}_2$	0	32.676

Energy conservation:

$$\sum_{\text{reactants}} n_m \left( h_{f,m} + \int_{T_{\text{ref}}}^{T_i} c_{p,m} dT \right) = \sum_{\text{products}} n_m \left( h_{f,m} + \int_{T_{\text{ref}}}^{T_f} c_{p,m} dT \right)$$

$$T_i = 800 [^\circ\text{R}] = 444 [\text{K}] \quad T_{\text{ref}} = 298.15 [\text{K}]$$

$$\Rightarrow T_f = 1769.7475 [\text{K}]$$

Less than part ②  $T_f = 1765.7 [\text{K}]$

$\Rightarrow$  Excess conditions decrease  $T_f$