

One 16-mer spans three partners' breakpoints — and only one of the three is a junction any patient is reported to carry

The three FET-family donors are identical over the ten nucleotides before the breakpoint, and the acceptor exon is the same in all three. Every row is TARGET mRNA, sense; the reagent is the reverse complement of the shaded window.

target mRNA (sense, 5' to 3')		breakpoint	
EWSR1 e12::NR4A3 e3	A A T G G T T T G A T G A T A T G C C C T G C G	EWSR1 · reported in patients at this exon	
FUS e10::NR4A3 e3	A C T G G T T T G A T G A T A T G C C C T G C G	FUS · no exon-resolved report at all	
TAF15 e11::NR4A3 e3	A C T G G T T T G A T G A T A T G C C C T G C G	TAF15 · partner reported, this exon not	



reagent 5'-GGGCATATCATCAAAC-3' (antisense), 8 donor and 8 acceptor bases either side of the seam

Left of the breakpoint rule, donor exon; right, NR4A3 acceptor exon; boxed, the one position at which the three donors differ. Shaded box, the window this reagent targets. Online, donor bases are blue, acceptor bases green and the boxed position purple.

The reagent is 5'-GGGCATATCATCAAAC-3', the reverse complement of the shaded window (5'-GTTTGATGATATGCCC-3'). Transcribing the shaded letters orders the sense strand, which is a different molecule with no antisense activity.

Research use only, not for administration. The bases alone, ordered as unmodified DNA, are a different molecule; order from fusion-junction-aso-sequences.csv.