

Two junction-spanning gapmer reagents for *NR4A3* fusion transcripts in extraskelatal myxoid chondrosarcoma, selected from 190 designs against a wild-type parent screen that condemns 87

Version of 2026-08-20 · submission format · built from a1c5630, tree not clean at build time

Author. Tristan D. McRae

Independent researcher, unaffiliated. Correspondence: trimcrae@gmail.com ORCID: [0000-0002-1823-1451](https://orcid.org/0000-0002-1823-1451)

Running title. Junction gapmer reagents for EMC

Keywords. antisense oligonucleotide; gapmer; RNase-H1; fusion transcript; *NR4A3*; extraskelatal myxoid chondrosarcoma; off-target screening

Abstract

Extraskelatal myxoid chondrosarcoma (EMC) is an ultra-rare sarcoma defined in most cases by an in-frame fusion of a variable partner gene to *NR4A3*. That junction is present in no normal transcript, so an antisense gapmer could in principle cleave the fusion and spare both parents. This work is computational: nothing was synthesised or tested, and nothing here asserts efficacy, safety, delivery or clinical readiness. Every sequence named is a research reagent for laboratory investigation only and must not be administered to any person or animal. Of 190 junction-spanning 16-mers tiled at 5-6-5 across the 38 in-frame junctions of five modelled partners, 87 let a mature wild-type parent transcript pair their whole catalytic gap over ten or more contiguous base pairs, 61 of those against wild-type *NR4A3* itself. Two reagents are named at the two most frequently reported breakpoints: 5'-GGGCATATCATCAAAC-3' at *EWSR1* exon 12 and 5'-GGGCATATCTTGTGTG-3' at *TAF15* exon 6, both at the panel's top gap-level margin of three, with longest wild-type parent runs of eight and nine base pairs. Two fusion-positive EMC cell models and two engineered constructs at these junctions are named as test articles, and the design pipeline is released for breakpoints outside the panel.

1 • Background

EMC is defined in the large majority of cases by an in-frame fusion of *EWSR1* to *NR4A3*,¹ with *TAF15* a substantial minority and *TCF12* and *TFG* rare.² The disease responds poorly to conventional cytotoxic chemotherapy,⁸ and the tyrosine-kinase inhibitors trialled in it give disease control more often than response.⁷

The fusion junction is the one feature of an EMC tumour that exists at the RNA level and in no normal cell. An antisense gapmer tiled across it recruits RNase-H1 to cleave the transcript it pairs, and the six-nucleotide DNA gap at the centre of a 5-6-5 architecture is where that cleavage occurs. Junction-directed oligonucleotides are a thirty-five-year lineage, reported against *BCR::ABL1*,¹⁰ *EWSR1::FLI1*,¹¹ *FGFR3::TACC3*,¹³ *PML::RARα*,¹⁵ *TPR223::ERG*¹⁶ and *DNAJB1::PRKACA*.¹⁸ No such design is reported for any *NR4A3* fusion in the literature retrieved here, and that absence is the reason this work exists: EMC is rare enough that the design step has not been done, and rare enough that no group is likely to do it as a by-product of something else.

What a junction design has to survive is specific to its construction. Both halves of the oligonucleotide are parent-gene sequence, so each of the fusion's two parents matches roughly half of it. Half-identity falls far outside the

mismatch budget of a conventional off-target search, so a parent transcript is not returned as a near-match. RNase-H1, however, does not require the whole duplex. It requires the gap to be paired. Whether a wild-type parent pairs the catalytic gap contiguously is therefore a separate question from overall similarity, and it is the question this work puts to all 190 designs before any of them is recommended.

2 • The reagents

The two reagents named for synthesis are the best available designs at the two junctions with a published exon-resolved breakpoint and the highest reported prevalence: 5'-GGGCATATCATCAAAC-3' at *EWSR1* exon 12 joined to *NR4A3* exon 3, and 5'-GGGCATATCTTGTGTG-3' at *TAF15* exon 6 joined to *NR4A3* exon 3 (Table 1). Both hold the panel's top gap-level margin of three, meaning three junction-unique bases inside the catalytic gap on the shorter side of the breakpoint. Neither pairs a wild-type parent through the gap at the ten-base-pair criterion adopted here.

Both sit close to that criterion, and the number belongs beside the sequence rather than in a limitations paragraph. The *EWSR1* reagent's longest wild-type parent run is eight base pairs and the *TAF15* reagent's is nine, both against wild-type *TFG*. Where the criterion is set therefore decides how the two read: at a cut of eight both fall inside the class this work marks as not to be ordered, at nine the *TAF15* reagent alone does, and only at ten does neither. The panel moves with them, 143 of the 190 designs pairing a wild-type parent through the whole gap at a cut of eight, 98 at nine and 87 at ten. Ten is adopted as a convention and is not measured for this architecture.

One hazard of this design space belongs beside the sequences rather than in a methods note. Consecutive registers of one seam differ by a single-base slide and can carry opposite verdicts, so a design that pairs a wild-type parent through its whole catalytic gap sits one nucleotide from one that does not (Table 2). Neither member of such a pair may be substituted for the other, and an oligonucleotide should be ordered from the canonical sequence file rather than transcribed from a printed table.

Predicted transcriptome load separates the two. The *EWSR1* reagent carries 123 gap-paired sense-strand near-matches at ten times the default search depth, recounting to six gene loci, none of them on a parent transcript; the *TAF15* reagent carries eight such near-matches at five loci. The *EWSR1* reagent also carries a sense-strand near-match in wild-type *TAF15* precursor RNA at two mismatches, one of them inside the catalytic gap, spanning an intron-exon boundary. That site is the cost of the same ten shared donor bases that let one oligonucleotide span the *EWSR1*, *TAF15* and *FUS* breakpoints at once (Figure 1). The *TAF15* reagent carries no sense-strand precursor site at all. Neither load is a disqualification, and neither is a statement about safety: these are predictions from sequence search, not measured off-target activity.

Discounted by the breakpoint distribution of an 18-case series,²² the two junctions account for 68.4% of molecularly confirmed cases in a 58-case cohort,⁹ roughly two thirds. That figure prices which published junctions the two reagents address. It is not a coverage measurement, no patient having been screened with either sequence, and its interval is wide for the denominators rather than for the estimate, spanning 39.9% to 82.8% when each breakpoint fraction is taken to its own Wilson bound.

3 • Selection from a panel of 190 designs

The two reagents above are what survived a screen applied uniformly to the whole panel. Across the 38 in-frame junctions of five modelled partners, 190 junction-spanning designs were tiled and put through five specificity screens. Of those, 87 let one of six mature wild-type parent transcripts pair their entire catalytic gap over a contiguous duplex of at least ten base pairs, and 61 of the 87 do so against wild-type *NR4A3*. That is the single largest liability class in the panel, and 85 of the 87 are paired by one of the design's own two parent genes rather than

by an unrelated transcript. A second class is invisible to any screen over mature transcripts: 19 designs carry a sense-strand near-match in parent precursor RNA that pairs the gap in full and touches intronic sequence. Taken as a union rather than a sum, the two screens condemn 93 of the 190.

Lengthening the catalytic gap does not remove this liability, and the reason is arithmetic rather than empirical. Every base inside the gap comes from the donor exon or from the acceptor exon, so the junction-unique bases on the shorter side and the bases a wild-type parent can pair on the longer side tile the gap and sum to it. A longer gap therefore raises the margin available only by conceding parent-paired gap DNA at the design's own seam, and across the three geometries screened the count of designs pairing a parent at the ten-base-pair criterion is flat at 87, 88 and 87.

Three designs clear every screen applied here. None of the three sits at a junction any patient is reported to carry, which is what makes them mechanism controls rather than candidates. Selecting within each junction rather than across the panel is what makes the two named reagents available: 35 of the 38 junctions have a design that clears the parent screen, and all five junctions with a published exon-resolved breakpoint have one.

Two bounds on the cleanliness claim are load-bearing and are stated with it. The alignment screen is heuristic and stores at most 50 hits per query, so only 47 of the 183 filtered designs have hit lists short enough to assess for cleanliness at all; a count of clean designs is therefore a floor over that subset rather than a total over the panel. And search depth moves the result: of the nine designs that carry no sense-strand near-match at the default search ceiling, six are not clean at ten times it, three of the six having returned no near-match whatever at the shallower setting.

The ten-base-pair criterion is adopted rather than measured, and the comparison against null models does not resolve an excess specific to this disease. Scrambled sequences meet the parent screen at 6.2% against 45.8% for designs at real breakpoints, but chimeras built at real exon termini of the same two transcripts, at junctions almost never reported in a patient, meet it at 40.6%. Most of the liability is therefore what joining two exon termini of these genes gives, and across cuts of six to thirteen base pairs the excess of the observed rate over the strongest null changes sign four times. No cut in that range is a boundary the data picks out.

4 • Test articles

Five test articles carry a junction this panel designs against, and they divide into two sources with opposite limits.

Three are engineered constructs from a published functional study,²⁷ whose exon spans that paper states verbatim. Two of them, E-N and T-N*, carry exactly the two junctions the reagents above span, so both named reagents have a stated test article. Rebuilding the constructs is the faster route and its critical path contains no laboratory that has to answer an email. What it cannot buy is biological relevance: a complementary DNA over-expressed in a heterologous background is not the disease, so such an experiment speaks to junction-selective knockdown of the intended transcript and not to activity at endogenous expression from an endogenous locus.

The other two are patient-derived, identity-clean models reported with two EMC tumours,³⁷ USZ20-EMC1 (RRID:CVCL_C6MX) and USZ22-EMC2 (RRID:CVCL_C6MY). These are the only source of a fusion-positive EMC cell identified here. They are available on request from the originating laboratory with no repository deposit, and they are slow, at reported doubling times of five to six days as sarco-spheres passaged every two to three weeks, which constrains any exposure window. Their fusions are reported as *EWSR1* exon 13 and *TAF15* exon 6 joined to *NR4A3* exon 2 rather than exon 3. Reagents exist at both acceptors, so each line has one, and it is not in either case the same molecule as the reagent named above for the exon-3 acceptor. A reagent selected for one acceptor is not valid for the other.

One requirement is upstream of all of them. The breakpoint of the cell line or tumour sample used as the test article must be established at nucleotide resolution by RNA sequencing before any oligonucleotide is ordered. Every design here is specific to the exon pair it was tiled at, and none is valid for an unverified junction. Routine diagnosis does not supply the seam: break-apart *NR4A3* fluorescence in situ hybridisation detects a rearrangement irrespective of partner,⁶ so on its own it locates neither the partner nor the exon pair.

5 • The falsification experiment

The experimental design that would resolve the central uncertainty, an isogenic fusion-positive against fusion-negative comparison, has been published in an analogous fusion sarcoma. Fusion-specific antisense oligonucleotides against *NAB2::STAT6* in solitary fibrous tumour, evaluated against CRISPR-engineered isogenic fusion-positive and fusion-negative cells, reduced fusion expression by 58% and proliferation by 22% in vitro.³⁸

Three assay controls are required, and a knockdown assay alone distinguishes none of them. A positive control gapmer against an abundant housekeeping transcript in the same cells separates failed delivery from a reagent that reached its target and did not cleave it, and it must carry the same 5-6-5 β -D-oxy-locked-nucleic-acid phosphorothioate geometry as the test article, since uptake and endosomal release track chemistry class. A scrambled gapmer of the same chemistry and geometry, dinucleotide-preserving so that guanine content and the 5' run are held as well as base composition, separates the backbone-class component of toxicity; the scramble actually ordered must itself be put through the mature-parent screen before it is made, because 6.2% of scrambles pair a parent's whole catalytic gap at the ten-base-pair criterion and 1.8% do so against wild-type *NR4A3*. A fusion-negative isogenic comparator supplies the discrimination readout that neither of the other two gives, and no supplier of one is named here: such a pair has to be engineered. A fourth arm is available at no design cost: the three designs that clear every screen sit at junctions no patient is reported to carry, which makes them mechanism controls for gap-directed cleavage rather than candidates, and the extended report names them.

The decision threshold should be fixed before the experiment. Selectivity is the wild-type *NR4A3* half-maximal knockdown concentration divided by the fusion's, from a matched dose-response in the same wells, and the cut is 5.0, taken as a convention rather than measured for this comparison. A ratio of residual transcript at a single dose is not commensurate with it and must not be compared against the same cut: that ratio is bounded above by one divided by the fusion knockdown's complement, so at the 58% knockdown reported for the published experiment above it cannot exceed approximately 2.4 however selective the reagent is, and a cut of 5 would return falsification as arithmetic rather than as biology.

The replicate count follows from the variance rather than being asserted. At a replicate standard deviation of 0.35 on the natural-log scale, six independent biological replicates give about 80% power to falsify a true selectivity of 3 and three give about 30%, computed from a noncentral t. Above a standard deviation of about 0.65, no observed ratio at or above one can place the upper limit of a two-sided 95% interval below 5 at three replicates, so the test can fail only where the reagent is anti-selective. Such a test is void: it cannot fail, which is a different outcome from one that fails to falsify. The replicate count should be set from a pilot estimate with three as a floor and not a target, and the void gate applied to the upper confidence bound on the pilot's standard deviation rather than to its point estimate. The ranking is falsified only where the upper bound of the interval lies below the cut, never where a point estimate does.

Two limits of that threshold belong with any result. It is defined on wild-type *NR4A3*, which is the acceptor parent; no donor parent is read by the ratio, so a reagent can clear the cut while pairing a donor transcript through its whole catalytic gap. Wild-type transcript for the reagent's own donor parent, and for the parent carrying its longest duplex, should be measured in the same wells and reported beside the ratio against no cut. No cut is stated for a donor ratio because no retrieved measurement bounds the parent case, so a threshold on it would be a number with nothing

behind it; the donor measurement is required to be reported and is not required to clear anything. A result on the stated cut can therefore falsify the ranking and cannot falsify the rationale. And no multiplicity correction is imposed, because the family over which an error rate would be controlled is open-ended: the released procedure generates reagents against the same ranking without limit, so how many reagents have been tested against this cut, and in what order, is part of a result rather than context for it.

6 · Beyond the panel

The named reagents do not reach every patient, and the panel is bounded by what has been sequenced rather than by what can be designed. The procedure that produced the 190 designs is released unchanged with the artefacts, and it is the second deliverable of this work. Its input is the breakpoint at nucleotide resolution; an exon pair inferred from a break-apart assay is not sufficient. Given a declared exon pair it retrieves the parent transcripts, builds the modelled fusion, grades the pair for frame, tiles the junction-spanning gapmers, and runs the five screens, of which the two parent screens matter most, since pairing a parent through the whole catalytic gap surrenders the only advantage the modality has. A design is certifiable where all five screens could be run on it and it cleared all five; a design one screen cannot address is uncertifiable whatever the other four return. What the procedure yields is a candidate, not a validated reagent.

7 · Discussion

Designability is not the constraint in this disease. Junction-spanning designs exist at every in-frame *NR4A3* fusion junction modelled here, and 35 of the 38 have one that clears the parent screen. The constraint is discrimination between the fusion and its parents, and it is not resolved here. A fusion-junction design's most plausible wild-type liability is its own parent, reached either in the mature transcript or across a splice junction in precursor RNA, and both compartments are searched in this work before any molecule exists. The four reports of parental sparing cited here^{13,14,15,16} were all made on molecules already synthesised, and three of the four went further than cells. Every one of those readouts required the molecule to exist first. What other groups do at the design stage is not established here; no survey of published design pipelines was performed.

The premise that sparing wild-type *NR4A3* is worth a specificity cost deserves examination rather than assumption. *NR4A3* has two close paralogues and the family is functionally redundant where it has been tested,^{29,30} which cuts against the premise; the evidence is not decisive in either direction.

Several limits bound what any test of these reagents could show. Every screened count here is for one architecture, a 16-mer at 5-6-5. All five screens address hybridisation rather than cleavage, and none establishes that a predicted duplex forms or is cut. Counts are lower bounds, the alignment screen being heuristic and depth-dependent. Which exon pair a patient carries is not decidable from a break-apart assay, so no reagent named here is valid for a tumour whose seam has not been sequenced. The method-level novelty of this work is nil, junction-directed oligonucleotides being long established; what is new is the indication and the screen applied before synthesis. And systemic, antigen-dependent delivery of an oligonucleotide to a solid tumour remains unsolved, which is the gate this modality faces after any result reported here. Nothing in this work addresses it.

One constraint sits above all of the others and no reagent choice moves it. Every source of a test article named here ends at someone culturing cells, so the rate-limiting step is a laboratory rather than a line, a construct or an oligonucleotide.

8 • Methods

All analyses are computational, use public data, run in continuous integration at no compute cost, and commit their outputs. No laboratory work was performed. Full parameters, every screen's settings, the complete bounds on each claim and the per-design tables are in the extended report, which is the bioRxiv preprint of this work; the released artefacts are in the archive named under Data availability.

Canonical transcripts for the five partner genes and for *NR4A3* were obtained from Ensembl.⁴⁰ Junction-spanning 16-mer gapmers were tiled in a 5-6-5 β -D-oxy-locked-nucleic-acid/DNA/ β -D-oxy-locked-nucleic-acid geometry,³⁹ one design per register at which the breakpoint falls inside the six-nucleotide DNA gap, which admits five per junction. A design's gap-level margin is the count of junction-unique bases inside the gap on the shorter side of the breakpoint.

Five specificity screens were applied: an alignment screen against human RefSeq RNA, classifying each near-match by whether the catalytic gap is paired; an exhaustive transcript scan complete for substitutions within a one-mismatch budget; a screen of the parents' unspliced sequence; a mature-parent screen recording the longest contiguous duplex any of six wild-type parent transcripts forms through the catalytic gap; and a genome-wide screen over every position of GRCh38. A near-match is a transcript window pairing a design at 14 or more of its 16 positions. A design is liable where a wild-type parent pairs its whole catalytic gap over a contiguous run of ten base pairs or more, ten being adopted rather than measured. Null ensembles were built as scrambles of each design and as chimeras joining the same two parent transcripts at real exon termini, screened identically.

Tables

Every cell below is a column of *fusion-junction-aso-sequences.csv*, the canonical machine-readable record, except the test-article column of Table 1, which is a literature fact and carries its source in the caption. An oligonucleotide should be ordered from that file rather than transcribed from this page.

Table 1. The two reagents named for synthesis, with their parent-duplex label and their test article. Both are the best available design at their junction and both hold the panel's top gap-level margin. The parent-duplex column is the longest contiguous duplex any of six mature wild-type parent transcripts forms through the catalytic gap; neither reagent reaches the ten-base-pair criterion, and both sit close to it, so the length is printed rather than a pass mark. Test articles are the engineered constructs of Brenca et al. (PMID: 31020999), whose exon spans that paper states verbatim. The two patient-derived models of Bangerter et al. (PMID: 36316541) carry NR4A3 exon-2 acceptors and are therefore matched to different designs, not to these two. Nothing here has been synthesised or tested, and no sequence may be administered to any person or animal.

Table 1. The two reagents named for synthesis, with their parent-duplex label and their test article				
seam	reagent	gap-level margin	longest wild-type parent duplex through the gap	test article
<i>EWSR1</i> e12:: <i>NR4A3</i> e3	5'-GGGCATATCATCAAAC-3'	3	8 bp, wild-type <i>TFG</i>	E-N, engineered construct
<i>TAF15</i> e6:: <i>NR4A3</i> e3	5'-GGGCATATCTTGTGTG-3'	3	9 bp, wild-type <i>TFG</i>	T-N*, engineered construct

Table 2. Four near-identical designs at two seams, two orderable and two not. Each pair is two consecutive registers of one seam differing by a single-base slide, and the two members carry opposite verdicts: the condemned member pairs its whole catalytic gap against a wild-type parent gene at the ten-base-pair criterion, and the orderable member does not. Neither member of a pair may be substituted for the other, and neither is a reagent this paper names for synthesis. The pairing is read from the canonical file's own cross-reference column rather than asserted here.

Table 2. Four near-identical designs at two seams, two orderable and two not				
seam	design	verdict	gap-level margin	longest wild-type parent duplex through the gap
<i>TCF12</i> e7:: <i>NR4A3</i> e3	5'-GGCATATCAAGCGCTG-3'	DO NOT ORDER	2	11 bp, wild-type <i>TCF12</i>
<i>TCF12</i> e7:: <i>NR4A3</i> e3	5'-GGGCATATCAAGCGCT-3'	orderable (a single-base slide)	3	8 bp, wild-type <i>NR4A3</i>
<i>EWSR1</i> e12:: <i>NR4A3</i> e3	5'-CAGGGCATATCATCAA-3'	DO NOT ORDER	1	11 bp, wild-type <i>NR4A3</i>
<i>EWSR1</i> e12:: <i>NR4A3</i> e3	5'-AGGGCATATCATCAA-3'	orderable (a single-base slide)	2	7 bp, wild-type <i>TFG</i>

Figure legends

One 16-mer spans three partners' breakpoints — and only one of the three is a junction any patient is reported to carry

The three FET-family donors are identical over the ten nucleotides before the breakpoint, and the acceptor exon is the same in all three. Every row is TARGET mRNA, sense; the reagent is the reverse complement of the shaded window.

target mRNA (sense, 5' to 3')	breakpoint	
EWSR1 e12::NR4A3 e3	A A T G G T T T G A T G A T A T G C C C T G C G	EWSR1 · reported in patients at this exon
FUS e10::NR4A3 e3	A C T G G T T T G A T G A T A T G C C C T G C G	FUS · no exon-resolved report at all
TAF15 e11::NR4A3 e3	A C T G G T T T G A T G A T A T G C C C T G C G	TAF15 · partner reported, this exon not



reagent 5'-GGGCATATCATCAAAC-3' (antisense), 8 donor and 8 acceptor bases either side of the seam

Blue, donor exon; green, NR4A3 acceptor exon; boxed and purple, the one position at which the three donors differ. Shaded box, the window this reagent targets.

Every sequence row above is the TARGET mRNA read 5' to 3', not the oligonucleotide. The reagent is 5'-GGGCATATCATCAAAC-3', the reverse complement of the shaded window (5'-GTTTGATGATGCCC-3'). Transcribing the shaded letters orders the sense strand, which is a different molecule with no antisense activity.

The same paralogy that lets one reagent cover three fusions is why these designs are hard to discriminate from the parent transcripts: this reagent's gap-level margin is three junction-unique bases inside the six-nucleotide catalytic gap, the window RNase-H1 cleaves. Coverage is predicted from sequence and has not been measured.

Research use only, not for administration. The bases alone, ordered as unmodified DNA, are a different molecule; order from fusion-junction-aso-sequences.csv.

Figure 1. One 16-mer spans three partners' breakpoints. The junction windows of *EWSR1* exon 12, *TAF15* exon 11 and *FUS* exon 10 joined to *NR4A3* exon 3, aligned at the breakpoint. Blue, donor exon; green, acceptor exon; positions at which the three donors differ are boxed as well as coloured, for greyscale and colour-blind readers. The shaded box is the target window of 5'-GGGCATATCATCAAAC-3' with the 5-6-5 architecture below it. The three donors are identical over the ten nucleotides before the breakpoint, which is what makes one oligonucleotide junction-spanning at all three. Two of the three drawn junctions are reported in no patient: the exon-resolved *TAF15* breakpoints in this disease are at exon 6, and no exon-resolved *FUS* breakpoint has been published. This is a statement about sequence, not a claim that one reagent serves three patient groups. Coverage is predicted from sequence and has not been measured.

Declarations

Research use only, and not for administration to any person or animal. Every oligonucleotide sequence in this manuscript and its tables is a research reagent for laboratory investigation only. None is a medicine or a candidate drug, none has been synthesised or tested, and none may be administered to any human being or animal or supplied to anyone for that purpose. Ordering any of them from a commercial synthesis service is possible for anyone; doing so does not make it a treatment, and nothing in this manuscript should be read as licensing use in a patient. An oligonucleotide should not be ordered by copying it out of this article: the canonical record is `fusion-junction-aso-sequences.csv`. No sequence should be ordered at all until the breakpoint of the cell line or patient sample has been established at nucleotide resolution by RNA sequencing.

Ethics approval and consent to participate. Not applicable. No human subjects, human material or animals were involved.

Consent for publication. Not applicable.

Data and code availability. All code, graded artefacts and per-design tables are public and are deposited under [doi:10.5281/zenodo.22028916](https://doi.org/10.5281/zenodo.22028916). The extended report of this work, carrying every screen's full parameters and the complete bounds on each claim, is deposited as a preprint on bioRxiv.

Use of artificial intelligence. A large language model (Claude, Anthropic) was used throughout this work: to write and review the analysis code, to run the screens, to retrieve and check literature, and to draft and revise this manuscript. Every quantitative claim is tied by automated guard to the committed artefact that produces it. The author directed all work reported here and is responsible for its content.

Author contributions. T.D.M. is the sole author and directed all work reported here.

Funding. This work received no external funding and was self-funded by the author.

Competing interests. The author declares no financial competing interests: he holds no patent, patent application, equity or consultancy relating to any sequence or method described here.

Acknowledgements. None.

References

Numbering follows the extended report, so that a reference cited in both documents carries the same number in each. Metadata is read from retrieved bibliographic records.

1. Labelle Y, Zucman J, Stenman G, Kindblom LG, Knight J, Turc-Carel C, Dockhorn-Dworniczak B, Mandahl N, Desmaze C, Peter M. Oncogenic conversion of a novel orphan nuclear receptor by chromosome translocation. *Human molecular genetics*. 1995;4(12):2219-2226. PMID: 8634690. doi:10.1093/hmg/4.12.2219
2. Paioli A, Stacchiotti S, Campanacci D, Palmerini E, Frezza AM, Longhi A, Radaelli S, Donati DM, Beltrami G, Bianchi G, et al. Extraskeletal Myxoid Chondrosarcoma with Molecularly Confirmed Diagnosis: A Multicenter Retrospective Study Within the Italian Sarcoma Group. *Ann Surg Oncol*. 2021;28(2):1142-1150. PMID: 32572850. doi:10.1245/s10434-020-08737-7
3. Remiszewski P, Falkowski S, Szumera-Ciećkiewicz A, Spalek MJ, Rutkowski P, Czarnecka AM. From pathogenesis to the patient's bedside: a comprehensive review of extraskeletal myxoid chondrosarcoma. *Journal of cancer research and clinical oncology*. 2025;151(11):283. PMID: 41055792. doi:10.1007/s00432-025-06316-5
4. Stacchiotti S, Ferrari S, Redondo A, Hindi N, Palmerini E, Vaz Salgado MA, Frezza AM, Casali PG, Gutierrez A, Lopez-Pousa A, Grignani G, Italiano A, LeCesne A, Dumont S, Blay JY, Penel N, Bernabeu D, et al. Pazopanib for treatment of advanced extraskeletal myxoid chondrosarcoma: a multicentre, single-arm, phase 2 trial. *The Lancet Oncology*. 2019;20(9):1252-1262. PMID: 31331701. doi:10.1016/S1470-2045(19)30319-5
5. Stacchiotti S, Dagrada GP, Sanfilippo R, Negri T, Vittimberga I, Ferrari S, et al. Anthracycline-based chemotherapy in extraskeletal myxoid chondrosarcoma: a retrospective study. *Clinical Sarcoma Research*. 2013;3(1):16. PMID: 24345066. doi:10.1186/2045-3329-3-16
6. Huang SC, Lee JC, Hsu YC, Tsai JW, Kao YC, Hsieh TH, Chang YM, Chang KC, Wu PS, Chen PC, Chen CH, Chang CD, Lee PH, Tai HC, Liu TT, Wen MC, Li WS, Yu SC, Wang JC, Huang HY. Extraskeletal Myxoid Chondrosarcomas: The Uncommon Clinicopathologic Manifestations and Significance of TAF15::NR4A3 Fusion. *Modern pathology*. 2023;36(7):100161. PMID: 36948401. doi:10.1016/j.modpat.2023.100161
7. Skórski T, Szczylik C, Malaguarnera L, Calabretta B. Gene-targeted specific inhibition of chronic myeloid leukemia cell growth by BCR-ABL antisense oligodeoxynucleotides. *Folia histochemica et cytobiologica*. 1991;29(3):85-89. PMID: 1794439.
8. Toretsky JA, Connell Y, Neckers L, Bhat NK. Inhibition of EWS-FLI-1 fusion protein with antisense oligodeoxynucleotides. *Journal of neuro-oncology*. 1997;31(1-2):9-16. PMID: 9049825. doi:10.1023/a:1005716926800

9. Parker Kerrigan BC, Ledbetter D, Kronowitz M, Phillips L, Gumin J, Hossain A, Yang J, Mendt M, Singh S, Cogdell D, Ene C, Shpall E, Lang FF. RNAi technology targeting the FGFR3-TACC3 fusion breakpoint: an opportunity for precision medicine. *Neuro-oncology advances*. 2020;2(1):vdaa132. [PMID: 33241214](#). [doi:10.1093/noajnl/vdaa132](#)
10. Lee MS, An S, Song JY, Sung M, Jung K, Chang ES, Choi J, Oh DY, Jeon YK, Yang H, Lakshmi C, Park S, Han J, Lee SH, Choi YL. Cancer-Specific Sequences in the Diagnosis and Treatment of NUT Carcinoma. *Cancer research and treatment*. 2023;55(2):452-467. [PMID: 36265509](#). [doi:10.4143/crt.2022.910](#)
11. Ward SV, Sternsdorf T, Woods NB. Targeting expression of the leukemogenic PML-RAR α fusion protein by lentiviral vector-mediated small interfering RNA results in leukemic cell differentiation and apoptosis. *Human gene therapy*. 2011;22(12):1593-1598. [PMID: 21846246](#). [doi:10.1089/hum.2011.079](#)
12. Shao L, Tekedereli I, Wang J, Yuca E, Tsang S, Sood A, Lopez-Berestein G, Ozpolat B, Ittmann M. Highly specific targeting of the TMPRSS2/ERG fusion gene using liposomal nanovectors. *Clinical cancer research*. 2012;18(24):6648-6657. [PMID: 23052253](#). [doi:10.1158/1078-0432.ccr-12-2715](#)
13. Neumayer C, Ng D, Requena D, Jiang CS, Qureshi A, Vaughan R, Prakash TP, Revenko A, Simon SM. GalNAC-conjugated siRNA targeting the DNAJB1-PRKACA fusion junction in fibrolamellar hepatocellular carcinoma. *Molecular therapy*. 2024;32(1):140-151. [PMID: 37980543](#). [doi:10.1016/j.ymthe.2023.11.012](#)
14. Panagopoulos I, Mertens F, Isaksson M, Domanski HA, Brosjö O, Heim S, Bjerkehagen B, Sciort R, Dal Cin P, Fletcher JA, Fletcher CD, Mandahl N. Molecular genetic characterization of the EWS/CHN and RBP56/CHN fusion genes in extraskeletal myxoid chondrosarcoma. *Genes, chromosomes & cancer*. 2002;35(4):340-352. [PMID: 12378528](#). [doi:10.1002/gcc.10127](#)
15. Brenca M, Stacchiotti S, Fassetta K, Sbaraglia M, Janjusevic M, Racanelli D, et al. NR4A3 fusion proteins trigger an axon guidance switch that marks the difference between EWSR1 and TAF15 translocated extraskeletal myxoid chondrosarcomas. *J Pathol*. 2019;249(1):90-101. [PMID: 31020999](#). [doi:10.1002/path.5284](#)
16. Freire PR, Conneely OM. NR4A1 and NR4A3 restrict HSC proliferation via reciprocal regulation of C/EBP α and inflammatory signaling. *Blood*. 2018;131(10):1081-1093. [PMID: 29343483](#). [doi:10.1182/blood-2017-07-795757](#)
17. Beard JA, Tenga A, Chen T. The interplay of NR4A receptors and the oncogene-tumor suppressor networks in cancer. *Cellular signalling*. 2015;27(2):257-266. [PMID: 25446259](#). [doi:10.1016/j.cellsig.2014.11.009](#)
18. Bangerter JL, Harnisch KJ, Chen Y, Hagedorn C, Planas-Paz L, Pauli C. Establishment, characterization and functional testing of two novel ex vivo extraskeletal myxoid chondrosarcoma (EMC) cell models. *Human cell*. 2023;36(1):446-455. [PMID: 36316541](#). [doi:10.1007/s13577-022-00818-x](#)
19. Li Y, Nguyen JT, Ammanamanchi M, Zhou Z, Harbut EF, Mondaza-Hernandez JL, Meyer CA, Moura DS, Martin-Broto J, Hayenga HN, Bleris L. Reduction of Tumor Growth with RNA-Targeting Treatment of the NAB2-STAT6 Fusion Transcript in Solitary Fibrous Tumor Models. *Cancers*. 2023;15(12):3127. [PMID: 37370737](#). [doi:10.3390/cancers15123127](#)
20. Kauppinen S, Vester B, Wengel J. Locked nucleic acid (LNA): High affinity targeting of RNA for diagnostics and therapeutics. *Drug discovery today. Technologies*. 2005;2(3):287-290. [PMID: 24981949](#). [doi:10.1016/j.ddtec.2005.08.012](#)
21. Dyer SC, Austine-Orimoloye O, Azov AG, et al. *Ensembl* 2025. *Nucleic acids research*. 2025;53(D1):D948-D957. [PMID: 39656687](#). [doi:10.1093/nar/gkae1071](#)

Appendix A. Correction and supersession register

This appendix records corrections that reached the analyses behind this manuscript, so that a reader comparing it against earlier versions of the work can see what changed and why. The full chronology is in the working record deposited with the archive.

Appendix A1 — Withdrawn reagents

An earlier draft of this work recommended two reagents that a deeper re-screen subsequently withdrew. They are named here as withdrawn rather than dropped silently. Six of the nine designs that carried no sense-strand near-match at the default search ceiling lost that property at ten times the ceiling, three of them having returned no near-match at all at the shallower setting. The reagents named in the present manuscript are those that survive the deeper screen.

Appendix A2 — Exon-numbering correction

An earlier version of these analyses placed the acceptor junction using coding-exon indices and concatenated coding sequence to coding sequence, which discarded the 5' untranslated region that *NR4A3* transcript exon 3 carries and which a fusion transcript retains. The corrected model is mRNA-level. Exon numbers in this manuscript are transcript exon indices counted from the transcript 5' end, including non-coding exons; the two conventions differ for *TCF12*, *TFG* and *NR4A3*. An acceptor exon number read under the wrong convention selects a different reagent.

Appendix A3 — Terminology

The term *cleavage risk* was replaced by *gap-paired sense-strand match* on 2026-08-19, because the earlier term named a catalytic outcome for what is a sequence observation. No screen in this work predicts cleavage; each grades hybridisation only.