

1. Build the chimeric mRNA as the donor transcript through the 3' end of exon *d*, then acceptor exon *a* **whole** — its 5' untranslated region included, as a fusion transcript retains it — and everything 3' of it.
2. Translate from the donor's own initiator codon, so the reading frame is the donor's throughout.
3. Let j_o be the index of the first residue not encoded wholly by donor nucleotides. If the donor cut leaves a partial codon, the residue at j_o is completed by acceptor nucleotides and belongs to neither parent; if the cut is codon-aligned there is no such residue.
4. Verify the seam rather than assume it: the chimeric N-terminus must match the donor protein residue for residue up to j_o , the C-terminus must be the acceptor's own, and acceptor residue 1 must appear immediately 3' of the seam codon. A junction failing any of the three is an error, not a result.

5. Enumerate every 8- to 11-mer containing j_o — including those that begin at it, which a straddle test requiring a residue on each side would drop.
6. **The novelty filter, and the step Section B5 is about:** keep a peptide if it is not a substring of wild-type *EWSR1* and not a substring of wild-type *NR4A3*. That is a two-protein test. It does not consult the rest of the proteome, and it does not consult other isoforms of either parent — which is precisely how four peptides that occur in a normal *NR4A3* isoform passed it.

Step 6 is the whole of the defect: the filter is correct for what it tests and the set it tests against is too small by everything except two canonical sequences. Figure 1 shows the seam this produces, and the out-of-frame case beside it. Table 4 gives every pair the window produces with the grade the screen assigned it, so which 22 drop out is checkable rather than counted.




Figure 1. The seam codon is built from leftover donor nucleotides and the acceptor exon's retained 5' untranslated region, and its identity decides whether the isoform collision occurs. (a) The *EWSR1* exon 7 to *NR4A3* exon 3 junction. Donor coding sequence ends mid-codon with one nucleotide left over; two retained acceptor 5' untranslated nucleotides complete it, giving the seam residue at $j_o = 264$, after which *NR4A3* resumes in its own frame with its methionine 1 as internal residue 266. Four of the five in-frame junctions place aspartate at this position rather than asparagine, and that is the difference Section B5 is about. (b) An out-of-frame junction reads the same acceptor exon in a shifted register: 9 novel residues, then a premature stop 1,610 nucleotides upstream of the chimera's last exon-exon junction, which is the canonical nonsense-mediated-decay configuration. Predicted binding is a screen; neither panel is evidence of presentation.

Table 4. Every candidate exon pair, and why each is or is not carried forward. All 27 pairs the declared window produces, graded by the screen of §2.2: 5 in frame, 4 out of frame, 9 acceptor exon is non-coding, 9 no seam produced at this pair. *Donor phase* is the number of coding nucleotides of the donor's last codon left over at the cut, so phase 0 is a codon-aligned break and phases 1 and 2 build the seam codon from retained acceptor nucleotides. An *In frame* cell reading n/a is a pair with no chimeric reading frame for the question to be about, not a pair that failed it. These are a combinatorial window of declared exon pairs, not observed breakpoints, and the window is an assumption of this work rather than a finding of it (§2.2).

Table 4. Every candidate exon pair, and why each is or is not carried forward				
EWSR1 exon	NR4A3 exon	Donor phase	In frame	Grade
7	3	1	yes	in frame; peptides emitted
9	3	1	yes	in frame; peptides emitted
10	3	1	yes	in frame; peptides emitted
12	3	1	yes	in frame; peptides emitted
13	3	1	yes	in frame; peptides emitted
6	3	2	no	out of frame; read-through tract, then a premature stop
8	3	2	no	out of frame; read-through tract, then a premature stop
11	3	0	no	out of frame; read-through tract, then a premature stop
14	3	2	no	out of frame; read-through tract, then a premature stop
6	2	2	n/a	acceptor exon is non-coding; no chimeric ORF
7	2	1	n/a	acceptor exon is non-coding; no chimeric ORF
8	2	2	n/a	acceptor exon is non-coding; no chimeric ORF
9	2	1	n/a	acceptor exon is non-coding; no chimeric ORF
10	2	1	n/a	acceptor exon is non-coding; no chimeric ORF
11	2	0	n/a	acceptor exon is non-coding; no chimeric ORF
12	2	1	n/a	acceptor exon is non-coding; no chimeric ORF
13	2	1	n/a	acceptor exon is non-coding; no chimeric ORF
14	2	2	n/a	acceptor exon is non-coding; no chimeric ORF
6	4	2	n/a	no seam produced at this pair
7	4	1	n/a	no seam produced at this pair
8	4	2	n/a	no seam produced at this pair
9	4	1	n/a	no seam produced at this pair
10	4	1	n/a	no seam produced at this pair
11	4	0	n/a	no seam produced at this pair
12	4	1	n/a	no seam produced at this pair
13	4	1	n/a	no seam produced at this pair
14	4	2	n/a	no seam produced at this pair

That ten-allele panel is the instrument behind every binder figure here, and a wider one changes them: a 34-allele screen of the same peptides at the same threshold returns five strong peptide-allele calls rather than four, because NMPCVQAQY is also strong on HLA-A*30:02. Section 2.3 reports both panels.

3.1 The limit worth watching

The single most consequential of the movable limits is B2. Direct mass-spectrometry evidence that a fusion-junction peptide is presented at measurable abundance on a common allele, in any fusion-driven tumour, would convert the central question of this route from a prediction into an empirical one. The coverage instrument used here is disclosed as failing for exactly that reason: it computes an eligibility fraction for an epitope whose presentation is unestablished. No date is offered for that arrival, or for any other in this section. An earlier version of this paper carried a table of optimistic, expected and conservative years for four such capabilities; it is withdrawn, because a forecast a reader cannot check does not become checkable by being tabulated, and Section 6 does the work it was there for by saying what each arrival would and would not look like.

B1. Low predicted class I coverage, dependent on the screen

Proposition. On the panels and thresholds screened here, fewer than a third of patients carry an allele predicted to present any junction peptide — and that fraction is set by the screen at least as much as by the tumour.

Evidence. The 34-allele screen finds 4 presenting alleles and 30.4%; the ten-allele screen finds 3 and 27.4%. Both rest on five strong peptide-allele calls, one per presenting allele, and Section 2.3 gives their threshold sensitivity.

A high-confidence tier, and it is empty. The natural response to an undefended cut is to report a conservative tier beside it, and the conventional conservative cut for class I presentation is 0.2. At 0.2 this screen returns no presenting allele and no coverage at all: the least permissive call it makes is 0.3736, so every figure in this paper lives strictly between the conventional cut and a cut half its size. That is reported rather than suppressed, and it is the sharpest available statement of what these predictions are worth: they are not a set of confident calls with a permissive tail, they are entirely a permissive tail.

What would clear or move it. Three things, in ascending cost. Extending the panels beyond 10 and 34 class I alleles to the full validated set is a computational task and would raise the denominator. The acceptance threshold is a convention rather than a result — nothing here defends 0.5 against 0.4 or 0.6 — and settling what the cut should be for junction peptides would do more to this figure than any panel extension. Calibrating that cut against a benchmark of experimentally validated neopeptides is the form that settling would take. The experimentally validated fusion-junction epitopes in the literature are individual sequences across a few fusions, the HLA-A*24:02-restricted SYT-SSX junction peptide [17,18], the four *EWSR1::FLI1* breakpoint peptides of the Ewing sarcoma case report [20] and the fusion neoantigens of a head and neck series [21], and a handful of epitopes is not a set against which a threshold can be calibrated; calibrating on point-mutation neoantigens instead would import an assumption about junction peptides that is the very thing in question. Measured immunopeptidomics (B2) could promote peptides the predictor ranks weakly or remove ones it ranks strongly, in a direction not knowable in advance.

Residual. Some fraction of patients will have no presented junction peptide and no second antigen to substitute. That fraction is a real and permanent exclusion from this approach, and it should be stated in any protocol rather than discovered at screening.

B2. Predicted rather than measured presentation

Proposition. No junction peptide has been shown to be presented on the surface of an EMC cell.

Evidence. All binding figures in Section 2 come from the sequence-based predictor named there [2]. No EMC immunopeptidomic dataset is known to the author, and the repository contains none. The vaccination reports in Section 1 bear on this weakly, because they measure T-cell responses to synthetic peptides that were administered, which is not evidence that the same sequence is processed from the endogenous fusion protein and displayed on a tumour cell [17,18,20].

Evidence outside the vaccination literature. In a series of 34 paediatric acute leukaemias, T cells expanded from patient bone marrow were co-cultured with autologous leukaemic blasts and the responding population sorted on activation markers, so no peptide was administered and the reactivity arose against antigen the tumour itself had processed and displayed [23]. Three recovered T-cell receptors then separated fusion from wild-type sequence on patient-matched alleles in cells given the fusion transcript rather than pulsed peptide: *KMT2A::AFF1* on HLA-DPA1*02:01/DPB1*01:01, a second *KMT2A::AFF1* breakpoint on HLA-DQA1*03:03/DQB1*03:01, and *PICALM::MLLT10* on HLA-B*51:01 from a CD8 clone. Two of the three also killed leukaemic blasts from their own patients in culture, so the presenting cell was the tumour and not only an engineered surrogate. A fusion breakpoint therefore reaches a T cell by the endogenous route in humans, which earlier versions of this section recorded as unshown for any fusion.

What that evidence does not establish here. Its authors attribute the finding partly to anatomy: leukaemia arises in the tissues where T cells already reside, and the proximity makes engagement with blasts frequent. An EMC node is not that tissue, and a haematological malignancy speaks to neither microenvironment limit B6 and B7 record. The fusions differ, and *KMT2A* breakpoints cluster in a way *EWSR1::NR4A3* need not. The fusion-reactive clones sat at 0.0062%, 0.001% and 0.001% of the repertoire, were found only at diagnosis or at relapse and at no later timepoint despite deep sequencing, and one of the five samples taken through to receptor testing had no reactivity to either the fusion neoantigen or the blasts. Recognition is established rather than abundance: no eluate was run and no copy number per cell was obtained, so the measurement below is unaffected, as are this paper's coverage figures and its grading of this route.

What would clear it. Mass-spectrometry immunopeptidomics on EMC tumour tissue or on one of the patient-derived EMC cell lines that have been established and characterised [6,14]. A single positive identification of a junction-spanning peptide in an EMC eluate would convert the central premise of this route from predicted to observed.

What a negative would have to specify to mean anything, and this paper cannot supply it. The sentence "a negative result would be close to decisive" appeared in an earlier version of this section and was doing no work, because a null eluate bounds nothing until three quantities are fixed in advance: a stated limit of detection in peptide copies per cell, so that "not presented" is separated from "below the instrument"; a stated number of independent specimens and their HLA types, since the lead peptide is presented on one allele and a specimen not carrying it cannot test the claim; and a positive control peptide of known abundance eluted in the same run, without which a null is indistinguishable from a failed elution. This paper names those three requirements and does not meet any of them — naming them is not the same as having them. Their absence is why B2 is graded as bounded by access rather than by the disease, and it is what makes this the highest-value single experiment in the ledger.

Cost and owner. Requires tissue and a proteomics facility. It is not computational and cannot be performed by this programme.

B3. Self-adjacency and central tolerance

Proposition. The junction peptide is mostly self sequence with one or two novel residues at the seam, so the T-cell repertoire capable of recognising it may have been deleted or anergised.

Evidence. At the *EWSR1* exon 7 junction the seam carries a single novel codon formed from one leftover *EWSR1* nucleotide and two retained acceptor 5' untranslated nucleotides, after which *NR4A3* methionine 1 follows as an internal residue. The remainder of each peptide is parental sequence. Whether one altered residue is sufficient to break tolerance is not answerable by binding prediction.

The near-self search, which this section previously listed as unbuilt. Two computational filters were specified for this route and never built: a distance-to-self filter, and an analysis of whether the differences from self fall at anchor positions, which affect binding, or at positions contacting the T-cell receptor, which affect recognition. Both have now been run. Every one of the 11 predicted binders was searched against the reviewed human proteome including isoforms at a tolerance of one or two substitutions, with a per-peptide chance baseline from 50 residue shuffles of the same peptide researched identically.

Eight of the 11 have at least one human peptide within two substitutions, and every such neighbour lies in *EWSR1* itself, in an *NR4A3* isoform, or in the paralogue *NR4A1* — which is what a junction assembled from two self proteins should give, and is a check on the search as much as a result. Three findings matter. **The lead peptide is one substitution from self.** NMPCVQAQY differs at position 1 from DMPCVQAQY, which occurs in *NR4A3* isoform 3 — the same isoform Section B5 is about — and at positions 1 and 5 from EMPCVQAQY in an *NR4A1* isoform. The chance expectation is stated as a model rather than as a number: for a query q , draw $N = 50$ uniform random permutations of its own residues, search each against the same proteome at the same tolerance, and let $\mu(q)$ be the mean neighbour count over those draws. This holds length and amino-acid composition fixed and destroys only the order, so μ is the neighbour count attributable to q being a string of those residues rather than to q being that particular sequence. For NMPCVQAQY, $\mu = 0.02$ against an observed 2. **No p-value is quoted and none should be:** 50 draws bound a rate near zero loosely, the draws are not independent of the proteome's own composition, and the comparison this paper needs is an order of magnitude, not a tail probability. **Neither difference is at an anchor.** Position 1 and position 5 face outward or into the groove's middle rather than serving as the primary anchors at position 2 and the C-terminus, so a T cell raised against the neopeptide reads a surface that differs from the self peptide's at the positions it actually contacts — which is the configuration in which cross-recognition is most plausible and in which central tolerance is most likely to have acted. **And no binder in the screen has an anchor-only neighbour.** Zero of the 11 has a near-self peptide whose differences are confined to anchor positions, which would have been the worst case: an identical TCR-facing surface distinguished only by residues the T cell cannot see. MPPLRGDM's five neighbours are one *EWSR1* peptide, MPPLRGGP, counted once per isoform, differing at positions 8 and 9 — and against a chance expectation of 1.34 for that peptide, five is not distinguishable from chance.

Two caveats carried with the result. The anchor assignment is the general class I rule of position 2 and the C-terminus applied uniformly, not an allele-specific motif; HLA-A*01:01 in particular reads position 3 as a primary anchor, and the artifact marks every row so a reader can apply that. And sequence distance is not receptor distance: peptides three substitutions apart can present near-identical surfaces and peptides one apart can present different ones. This search excludes one more failure mode — a close self peptide nobody had looked for — and leaves the question of whether a repertoire exists exactly where it was.

What the clinical prior art bears on it. Junction peptides from two other fusions have raised T-cell responses in patients: peptide-specific cytotoxic T cells against the SYT-SSX junction in synovial sarcoma [17,18], and de novo CD4+ responses against all four *EWSR1::FLI1* breakpoint peptides in one patient with Ewing sarcoma [20]. A repertoire able to see a fusion seam therefore survives central tolerance at some junctions. Neither result concerns this junction, these alleles, or this junction's single-residue distance to DMPCVQAQY; the SYT-SSX responses were weak and were not accompanied by demonstrated benefit; and the Ewing sarcoma observation is one patient. The limit is narrowed by an existence proof elsewhere rather than answered here.

What would clear or narrow it. Only a T-cell reactivity assay against the specific peptide-HLA complex answers the question. The search above changes what such an assay would be testing: the comparator is no longer a hypothetical self peptide but a named one, DMPCVQAQY, at a known single-residue distance, in a protein the tumour also expresses.

Comparator. This limit is where EMC differs most sharply from the melanoma setting, in which selected neoantigens frequently arise from point mutations in a repertoire that has not been tolerated against them and in a tumour already under immune pressure.

B4. One strong CD4 epitope, on one allele of twenty-three

Proposition. An effective vaccine generally requires CD4 helper epitopes. The corrected junction supplies one predicted strong class II binder on a 23-allele panel, restricted to an allele carried by about one person in fifteen. That is a statement about the screen, and it is reported here as one.

History, and why this section changed. This arm was previously withheld because it sat on the superseded coordinate system, on a seam disjoint from the class I set; the builder was moved to the transcript model and the arm regenerated, so the two arms are certified to sit on the same seam. It was then reported as a negative on three DRB1 alleles — and Section 6.2 of that version named a wider DR, DP and DQ panel as the single most likely source of a change to it. That panel has now been run. The prediction was correct and the negative did not survive it.

Result. At the corrected *EWSR1* exon 7 junction, 15 candidate 15-mers were screened with MHCnuggets [12] against 23 class II alleles — 15 DRB1, DRB3*01:01, DRB4*01:01, DRB5*01:01, two DPB1 and three DQB1 — calling a peptide a binder below 1000 nM and strong below 100 nM. Every declared allele returned a score; none was unscreenable, so no cell of this panel is an absent reading. 44 peptide-allele pairs bind, spread over 14 of the 23 alleles, and one is strong: SYGQQNMPCVQAQYS on DRB1*14:01 at 66.1 nM. The three-allele subset the previous version reported reproduces exactly within the wider run — YSQSSSYGQQNMPC at 261.6 nM and QYSQQSSSYGQQNMPC at 438.9 nM, both on DRB1*07:01 — so the widening is additive and the earlier negative was a property of the panel, not of a different computation.

What the positive does and does not bound, and it is less than it looks. The 15 candidates are single-residue-offset windows over one seam sharing 14 of 15 residues, so the number of independent peptides tested is nearer one than fifteen, and the 44 binders are overwhelmingly the same short stretch of sequence seen on different alleles. One strong call on 23 alleles is a per-allele rate of 1 in 23, and the allele it lands on is not a common one. Class II prediction is substantially less accurate than class I, so no class II call here is treated as the equal of a class I call of the same nominal strength: class II peptides are

not length-restricted by a closed groove, the binding register is not fixed, and predictors for it are trained on far less measured data. The single strong call should be read as a weaker statement than any single class I call in this paper, and no class II prediction here has been calibrated against anything measured. What the result does establish is narrow and worth having: the junction is not devoid of predicted helper epitope, which is what three alleles could not distinguish from a junction that is.

What would clear or move it. Only measured class II presentation settles the prediction; a wider panel now has much less headroom to move it than it did.

How the two arms should be ranked, which is a separate question. Earlier versions of this section carried the class II result as the lesser of the two, and the human evidence does not support that ranking. Of the three fusion-specific restrictions recovered from endogenously presented breakpoints in paediatric leukaemia, two were class II and one class I [23], and the only de novo T-cell response reported to a fusion-breakpoint vaccine in a patient was CD4+ rather than CD8+ [20]. Three observations, across three fusions and three groups, are a pattern to test and not a rule: each restriction is one patient's alleles, none is in a solid tumour, and none concerns this junction. The inverse ranking is no better supported and is not asserted here; what the pattern removes is the ground for assuming the CD8 arm is the one that matters. The two predictions remain ordered, for the reason given above, and that ordering is a property of the models rather than of the responses. A construct can also supply help from a heterologous source rather than from the junction itself, which is standard practice in peptide vaccine design and would make this limit a design constraint rather than a blocking one.

Consequence for the construct, and for the combined figure. Two things follow directly, and both changed with the panel. The candidate construct regenerated at the corrected junction now carries the CD4 epitope as well as the two strong class I epitopes NMPCVQAQY and QQNMPCVQAQY on HLA-B*15:01, and its minimal synthetic long peptide is 15 residues carrying both arms, where the class I-only version was 11. And the combined CD8 and CD4 coverage figure can be computed at all for the first time: class II coverage is 6.5% on DRB1*14:01 alone, and the fraction of patients carrying at least one presenting allele of each class is 1.8%, the product of the two arms' coverages:

$$C_{IAII} = C_I \times C_{II} = 0.2737 \times 0.0649 = 0.0178$$

The class I term is the ten-allele screen's 27.4%, not the 34-allele screen's 30.4% quoted two paragraphs above: the instrument builds its class I allele set from the junction screen rather than from the broad-panel scan. On the 34-allele set the same product gives 2.0%. The product assumes independence between the class I and class II loci, the same assumption the within-class formula makes and no weaker. HLA-A, HLA-B and DRB1 sit on chromosome 6 in linkage disequilibrium, so it is an approximation whose direction is not known without haplotype frequencies. The magnitude is bounded even though the direction is not, and by the same argument §2.3 applies to the within-class formula. This figure is an intersection rather than a union, so the bounds invert: under any dependence structure whatever it lies in [0.0%, 6.5%], and the upper end is the useful half. A construct that needs a presenting allele of each class can never reach a larger fraction of patients than its scarcer arm alone, which here is the class II arm at 6.5%. That ceiling holds without any haplotype data, and it is 3.6 times the product rather than an order of magnitude above it. It is a carrier-frequency product and nothing more: it says what fraction of patients carry a presenting allele of each class, not what a class II response contributes. Class II help for a CD8 response is cognate rather than additive, and no model of that is offered here. That figure is smaller than either arm alone and is the eligibility fraction for a construct that needs both — and, like every coverage number in this paper, it is a point on the curves of Section 2.3 rather than an estimate with a tolerance.

Status. Positive, on one allele of 23, with the previous negative explained as a panel artifact rather than overturned by a different method.

B5. Four peptides occur in an *NR4A3* isoform

Proposition, as originally stated. The junction peptides had been tested for novelty against two proteins rather than against the proteome, so their absence from normal human proteins was not established.

Result. All 174 distinct junction peptides were searched by exact substring against the UniProt reviewed human proteome, isoform sequences included. 170 are absent from every reviewed human protein. All 4 strong binders survive, including the *EWSR1* exon 7 lead NMPCVQAQY. The limit is largely cleared, and the sequence-level novelty premise of this route holds for the great majority of the peptide set.

The four that do not, and which junctions they belong to. DMPCVQAQ, DMPCVQAQY, DMPCVQAQYS and DMPCVQAQYSP all occur in Q92570-3, an isoform of *NR4A3* itself. One of them, DMPCVQAQY, is a predicted binder on HLA-B*35:01 at 369.1 nM. Those four peptides are not tumour-exclusive, and DMPCVQAQY is withdrawn as a candidate.

The pattern is a mechanism, and it survives the transcript model. A seam defined by exon boundaries is defined by whichever transcript declares them, and every figure in Section 2 comes from one canonical transcript per gene. The junction was rebuilt across all 99 protein-coding *EWSR1* transcripts against all 4 of *NR4A3*, at each of the five in-frame donor exons: 1,980 pairs, of which 970 emit an in-frame seam. The seam residue is not stable across them, taking nine distinct values — aspartate in 502 pairs, glycine in 251, asparagine in 118, serine in 59, no seam residue at all in 31, and four other residues in the remaining 9.

The collision appears in exactly the 502 aspartate-seam pairs and in none of the other 468. It is therefore not a property of the canonical annotation, and not a single case study: it is a property of the aspartate seam, reproduced independently in 502 transcript pairs, and its absence is equally determined. That is the mechanism this section had been asserting from one pair.

The pattern is not random across the junction set. All four collided peptides belong to the same four junctions, *EWSR1* exons 9, 10, 12 and 13, which are the four whose seam codon is aspartate on the canonical transcript; the *EWSR1* exon 7 junction has an asparagine seam codon and none of its peptides collides. So four of the five in-frame junctions share a seam whose most seam-proximal peptides reproduce a normal *NR4A3* isoform sequence, and the one junction that is clean is the commonly reported public one that carries the lead binder. Within the affected junctions the collision is confined to peptides that begin at the seam residue, which are the peptides with the least donor content; those extending further into *EWSR1*, including the strong binder RGDMPVCVQAQY, remain novel. The consequence for design is specific rather than general: for the aspartate-seam junctions, a construct should not rely on the seam-proximal window.

The methodological finding. The upstream novelty filter compares each candidate against the canonical parent proteins only, so an isoform that carries the seam sequence passes it unseen. That is a defect in the filter rather than in these particular junctions, and it will recur for any breakpoint whose

seam residue reconstructs an isoform boundary. The proteome search reports this condition explicitly rather than silently discarding the hits.

The repair, stated as a test rather than as an aspiration. Saying the filter should be made isoform-aware names the defect without saying what to compute, so here is what to compute. Everything 3' of the seam residue is the acceptor's own sequence, retained whole, so a seam-proximal peptide is the seam residue followed by an acceptor stem. It reproduces a normal protein exactly when some isoform of the acceptor gene already places that same residue immediately 5' of that same stem, which is what an alternative first exon does. The residues for which that holds are a property of the acceptor gene alone; call them its collision alphabet. The test is then one character:

a junction whose seam residue lies in the acceptor's collision alphabet has a seam-proximal window that is not tumour-exclusive.

The alphabet is computed once per acceptor from its isoform set and answers the question for every donor and every breakpoint without a further search, where the proteome scan it anticipates costs one pass over the proteome per peptide. Here the alphabet is a single residue, aspartate, contributed by Q92570-3.

It is exact on every transcript pair this locus can produce. Applied to the 970 in-frame pairs of the 1,980 above, the one-residue test agrees with the proteome search on all 970: 502 predicted collisions, all observed; 468 predicted clean, none colliding; no disagreement in either direction. That the two ends match is the point. A rule that flagged the aspartate seam would be unremarkable if it also flagged the glycine seam's 251 pairs or the asparagine seam's 118, and it flags neither.

And it does not replace the search, which is the failure mode from the other side. The test anticipates one condition: a seam-proximal peptide reconstructing an acceptor isoform boundary. It is silent about a peptide colliding with an unrelated protein, and silent about peptides extending far enough into the donor to become novel again — this locus's own strong binder RGDMPVCVQAQY carries the same stem, begins three residues 5' of the seam, and survives. A filter that ran this instead of the proteome search would reintroduce the defect it was built to catch. It is a pre-screen that says where the search will find something, and it costs one comparison.

What a clean result does not license. A peptide absent from every reviewed human protein is not thereby safe. A T-cell receptor engages a peptide-MHC surface rather than a sequence, so a peptide differing from a self peptide at a position that does not contact the receptor can still be cross-recognised. Unreviewed sequences were searched separately, over 127,090 entries: of the 170 peptides absent from every reviewed protein, 12 occur in at least one unreviewed entry, and none of those 12 is a predicted binder. A hit among predicted-and-unreviewed entries is not evidence that a normal protein carries the peptide, and a miss there is not evidence of absence, so this withdraws no peptide and confirms none; what it establishes is where the sequence-novelty premise is weakest, and that is not where the candidates are. This test excludes one specific failure mode and leaves the others standing.

B6. Immunologically cold microenvironment

Proposition. EMC has a low mutational burden and a sparse infiltrate, so there is little pre-existing antigen-specific response for an intervention to amplify. The mutational-burden half is not disputed here. The infiltrate half was an assumption when this section was first written and is now a measurement, reported at B8 and not restated here.

What that measurement does to this proposition, and what it does not. It supports the direction B6 asserts and does not license the word cold, because the comparator is other sarcomas rather than an immune-normal reference, and because a bulk average over whole tissue cannot tell a tumour with no T cells from one whose T cells are held outside it. That second reading is B7's, the two are not the same claim, and B7 states why the evidence now available does not choose between them.

Evidence, and an observation about this programme's own grading. This proposition is why this programme set most immune-modulating classes aside for this disease in its route ledger, a committed record of forty candidate modalities each carrying a verdict and a reason. Those reasons are quoted below from that ledger: they are this author's own prior notes, not positions taken in the literature. Innate agonists of the STING, TLR and RIG-I classes were excluded there because such agonism "supplies the danger signal and the priming context; it does not supply antigens, and a genome as quiet as EMC's is short of antigens rather than short of priming". In-situ vaccination was excluded because it "releases and adjuvants the antigens the tumour already has", so "releasing more of nothing does not help". Checkpoint inhibitors beyond PD-1, costimulatory agonists, adenosine-axis inhibitors and regulatory T-cell depletion were each excluded for acting on a response presumed absent.

Every one of those is an argument about antigen supply, and a vaccine is an antigen supply, so none of them argues against the class in a configuration where antigen is supplied exogenously. That is why the combination in Section 4 was never graded here as a unit.

The symmetry is not exact, and an earlier version of this paper asserted that it was. The vaccine was not parked for want of priming: its ledger entry is on the board rather than excluded, and its stated reason for being parked is immunogenicity, "which its own record states is not a computational question". The standing blocker against it names two things, that EMC is antigen-cold and that the fusion junction is a weak peptide-HLA, and the second is an antigen-side objection no priming-directed partner answers. The correct statement is the weaker one: a partner supplies what several exclusions called missing, and does not supply what the vaccine's own blocker calls missing.

What would clear it. Clinical evaluation of a vaccine together with an agent that supplies priming or releases inhibition. Section 4 sets out the specific backbone for which EMC evidence already exists.

B7. Physical exclusion by the myxoid matrix

Proposition. EMC is immune-excluded by a dense chondroitin-sulfate gel, which is a physical barrier rather than a signalling programme.

Evidence. The myxoid matrix is the disease's defining histological compartment. The oncofetal chondroitin sulfate pathway is the mechanism proposed for it here by analogy: the glycosaminoglycan biosynthesis genes of that pathway are differentially expressed and correlated with immune response in placenta and colorectal cancer [8], which is the tissue setting that study examined. That pathway has since been read in the same two EMC cohorts and the reading does not carry the analogy across: the linker and core-protein modules are higher in EMC than in the comparator sarcomas on both platforms, the backbone-polymerisation and 4-O-sulfotransferase modules disagree between platforms, the sulfate-donor module is lower on both, and three sulfotransferase modules fall below the read's own readability floor. Values have one home, [research/modalities/emc-expression-panels.json](#), read 2. A transcript level is a capacity to build a glycan and not the glycan, so the inference from those tissues to this one remains the author's. Transforming growth factor beta inhibition, which is the standard proposal

for immune-excluded tumours, was set aside for EMC precisely on the ground that the exclusion here is physical rather than driven by a fibroblast programme.

Why this limit is distinct from B6. A cold tumour lacks a response. An excluded tumour may have a response that cannot reach the target. These require different interventions, and an intervention aimed at one does not address the other. A vaccine can raise the circulating frequency of junction-specific T cells without changing whether those cells can enter the tumour.

Why the new EMC immune evidence does not choose between them. Both instruments that speak to the question arrived in 2026 and neither sees what the other sees. The bulk cohort reading of B8 runs in the direction B6 predicts, and it is an average over whole tissue with no spatial resolution, so a cold tumour and an excluded one return the same value. The multiplex immunofluorescence arm of the twelve-tumour series runs in the direction B7 predicts, because a T cell that is present and suppressed is not a T cell that is absent, and it is two specimens compared within themselves and called proof of concept by its own authors [15]. The instrument with a cohort has no spatial resolution and the instrument with spatial resolution has no cohort. This paper therefore does not resolve the disagreement and does not treat either limit as retired by the other's evidence; what would resolve it is the observation B8 names, per-tumour spatial profiling on a series large enough to estimate a fraction.

What would clear or mitigate it. Vascular normalisation is the mechanism with the most direct EMC evidence, discussed in Section 4. Matrix-directed approaches, including addressing the oncofetal chondroitin sulfate modification itself, are registered in this programme as candidates and are not resolved.

B8. Thin EMC immune-profiling data

Proposition. The characterisations of EMC as cold and as excluded were inferred from the disease's mutational burden, its histology and sarcoma-wide immunotherapy experience rather than from EMC-specific immune profiling. The proposition as originally stated, that no such profiling exists, no longer holds; what follows replaces it.

Consequence. Several exclusions above rested on an assumption nobody had measured in this disease, and an absent reading was never a reading of absence.

What has since been read, in EMC tumour tissue. Two archival bulk expression cohorts, 6 and 10 EMC tumours against 29 and 6 comparator sarcomas, were scored against lineage, effector and antigen-presentation modules and against published hallmark infiltration proxies. The interferon- γ effector axis, the myeloid compartment and class II antigen presentation are lower in EMC than in the comparator sarcomas on both platforms; the cytotoxic-lymphocyte and T-cell modules are lower on the larger cohort; and two hallmark proxies, the interferon- γ response and the lymphocyte-weighted allograft-rejection set, are lower on both. The class I apparatus is the exception. β 2-microglobulin and the class I heavy chains sit near the top of the expressed range on the larger platform, and HLA-A, HLA-C and TAP1 read flat against the comparator rather than lost. Values and their contrasts have one home, `research/modalities/emc-expression-panels.json`, reads 8 and 19.

What points the other way, in individual patients. Twelve archival EMC tumours profiled by whole-transcriptome targeted sequencing carry more B-cell infiltration in the low-risk half of the series than in the high-risk half, and proof-of-concept multiplex immunofluorescence on two of the specimens reports exhausted CD3+CD8+PD1+ T cells and FOXP3+ regulatory T cells in the high-risk one; the authors state that every immunofluorescence comparison is a within-specimen region-of-interest analysis, that their prognostic model overfits, and that the work is exploratory and hypothesis-generating [15]. A single reported case profiled as immune-enriched despite a low mutational burden is graded in Section 4 [16]. Neither is a cohort estimate, and a quiet distribution with occasional infiltrated members is consistent with both those observations and with the paragraph above. What the twelve-tumour series adds beyond direction is that the variation it reports is within EMC rather than between EMC and something else, which is the axis a patient-selection argument would have to run on.

What none of it settles. The comparator arm is other sarcomas, which are not an immune-normal reference, so lower than that comparator is not the same statement as cold. Bulk expression cannot locate a transcript in a cell, and therefore cannot separate a cold tumour from the excluded one B7 describes. Sixteen tumours cannot estimate what fraction of EMC is infiltrated, which is the quantity a patient-selection argument would need. The class I reading is transcript abundance in a cohort, not protein on a cell surface in a patient, and class I loss would independently disable every antigen-directed route including this one. And the twelve-tumour series has been read here only as its published abstract, because the publisher returned an access error to this work's retrieval runner, so its infiltration method, its external validation cohorts and its per-specimen values are unexamined.

What would still clear it. Per-tumour profiling with spatial resolution on a series large enough to estimate that fraction, and class I assessed as protein.

Cost and owner. The cohort reading is done. It ran on a continuous-integration runner over two public archival series, with no rental, no accelerator and no purchased compute. What remains requires tissue, and it is still the cheapest tissue-based item in the ledger and the one that most efficiently informs the others.

B9. Manufacturing economics at this incidence

Proposition. An individualised vaccine requires per-patient sequencing, design and manufacture, and EMC's incidence is well under one per million per year.

Consequence. No independent programme can supply the manufacturing, and the economics that support an individualised product in melanoma do not obviously extend to a disease with this incidence.

What would mitigate it. Two features of this target reduce the requirement relative to a general individualised product. First, the antigen is not discovered per patient by tumour sequencing but is determined by which exon pair the patient carries, of which five are in frame; the design space is therefore small, enumerable in advance, and shared across patients with the same breakpoint. Second, because the fusion is truncal, the antigen does not need re-selection over the disease course. A small fixed panel of breakpoint-specific constructs, allocated by a diagnostic assay, is closer to a stratified product than to a bespoke one. A construct of that shape has been made and administered once already: the Ewing sarcoma vaccine was an off-the-shelf multi-peptide product spanning a recurrent breakpoint rather than a per-patient design [20], and routine fusion analysis in a clinical genomics workflow finds EWS breakpoints clustered tightly, with about three quarters of Ewing sarcoma and desmoplastic small round cell tumours sharing one exon 7 motif [22]. Whether that is commercially tractable is a question for a platform holder and not one this analysis can answer.

B10. Trial design below the randomisation threshold

Proposition. At this incidence a conventional randomised trial of a vaccine addition is not feasible.

Evidence and mitigation. A histology-specific EMC cohort within a sarcoma master protocol has already been executed and reported for a different regimen, enrolling 24 patients across nine centres in three countries over four years [5]. That shows the vehicle exists, and it gives an accrual rate of about six unselected patients a year across nine centres.

And that rate does not survive this paper's own eligibility filter. A junction-vaccine cohort cannot enrol unselected EMC: it needs *EWSR1::NR4A3*, which is 62 to 75% of cases [7], and then an HLA type that presents a junction peptide, which is 8.5 to 30.4% of those (B1). Applying both fractions to that trial's own accrual leaves roughly 0.3 to 1.4 eligible patients a year, so a cohort of the size already run would take on the order of two decades, and a powered comparison considerably longer. That is what makes B10 a limit of the disease rather than of study design: the vehicle exists and the patients to put in it do not arrive fast enough. The endpoint question is separate — response-based endpoints behave poorly in indolent tumours, and the existing EMC cohort reported progression-free survival at a fixed time point.

4. An ungraded combination

The limits above do not resolve independently. B6 and B7 are properties of the tumour that a vaccine cannot address, and B1 and B3 are properties of the antigen that a checkpoint inhibitor cannot address. The relevant question is therefore not whether a junction vaccine works in EMC, which is the question that was asked and answered negatively, but whether a junction vaccine adds anything to a backbone that already has EMC-specific activity.

Such a backbone exists. A phase 2 histology-specific EMC cohort within the IMMUNOSARC II master protocol evaluated sunitinib with nivolumab in adults with advanced, progressing, centrally confirmed EMC across nine centres in Spain, Italy and the United Kingdom. Of 23 evaluable patients, 16 were progression-free at 6 months, median progression-free survival was 13.2 months (95% CI 5.7 to 20.7), and there were 2 partial responses [5]. This is a conference abstract and has not been peer-reviewed; the full publication is not available, and no results are posted for the registration. It is reported here as the most direct EMC-specific evidence available for an immunotherapy-containing regimen, and not as evidence of efficacy. The preceding mixed-histology phase Ib/II study of the same combination is published [4].

Three features make this backbone the natural context for the vaccine question. The antiangiogenic component addresses B7 by a mechanism — vascular normalisation reduces the physical and vascular barriers to lymphocyte entry — and antiangiogenic tyrosine kinase inhibitors carry the most consistent prospective signal in this disease: pazopanib gave an objective response in 4 of 22 evaluable patients with a median progression-free survival of about 19 months [11], and a sunitinib series reported activity in translocated EMC [13]. The checkpoint component addresses the release arm of B6. Neither addresses antigen supply, which is what the vaccine would contribute.

One reported single-agent checkpoint response, and what it changes. The checkpoint half of that backbone has never been given alone in a reported EMC cohort, and one patient has been reported outside one: a man with *EWSR1::NR4A3* EMC, microsatellite stable at a tumour mutational burden of 0.67 mut/Mb and with *TSC2* loss, whose microenvironment a commercial assay profiled as immune-enriched and fibrotic with high PD-L1, and who received single-agent pembrolizumab with a near-complete response reported over 15 cycles [16]. One patient in an industry-authored conference abstract that has not been peer reviewed supports no response rate, and it is not read here as a promotion of the checkpoint arm. What it does is name the selection axis: the markers conventionally used to choose a patient for checkpoint inhibition and the markers describing that patient's microenvironment pointed in opposite directions in the same tumour, which is the heterogeneity B8's cohort reading permits and cannot measure. That is a question for B8's observation to settle, not a case for adding a third agent.

This is also the architecture of the melanoma programme behind the recent announcement, where the individualised vaccine is given with pembrolizumab and never alone [3,9]. The transfer to EMC fails on antigen depth — melanoma supplies a pool of private neoantigens, EMC one junction — and not on architecture.

What this section is claiming, and what it is not. The proposition is that adding a breakpoint-matched junction construct to a checkpoint and antiangiogenic backbone gives a combination in which each component covers a limit the others do not, and that it has been graded as a unit nowhere — including in this programme's own ledger, which is where B6's observation comes from. It is not a trial proposal. No population, comparator, endpoint, effect size or sample size is specified here; a single-arm addition to a backbone whose own six-month progression-free rate is 16 of 23 could not attribute an effect to the vaccine; and B10's arithmetic says the eligible patients do not arrive at a workable rate. Whether the combination merits evaluation at all turns on B2. Nothing here recommends that any patient receive any of these agents outside a clinical trial.

5. Present work and pending arrivals

Four items stood on this list in the previous version, all needing nothing but public data. Two have since been done and are reported above rather than promised here: the distance-to-self and anchor-versus-contact-position filters, which are now the near-self search of B3, and the extension of the class II panel from three DR alleles to 23 across DR, DP and DQ, which is now the positive result of B4. The class I panels remain at 10 and 34 alleles against the full validated set, which would raise the denominator of every coverage figure. A defended acceptance threshold for junction peptides is still absent, and Section 2.3 now shows exactly how much rests on it. The isoform-aware novelty filter identified in B5 is still unbuilt. None of the remainder requires permission, material or funding.

Everything else waits. Measured presentation (B2) would change the character of this route rather than its score, because the screen in Section 2 would stop being a stand-in and become a hypothesis with a calibrating dataset. B8 is no longer an absence: two archival EMC expression series have been read here and a twelve-tumour series has been published, and what remains of that limit is spatial resolution on a series large enough to estimate an infiltrated fraction, together with class I assessed as protein rather than as transcript. It is still the cheapest tissue-gated item and the one that informs the most others, because infiltrate location and HLA class I status bear on every antigen-directed route in this disease. Access to a patient-derived line would additionally make B2 executable rather than merely specified; these are separate arrivals and neither implies the other. B6 and B7 are properties of the tumour that no computational advance mitigates.

6. Distinguishing a real advance from an apparent one

Each capability this route waits on has a plausible near-miss that the literature would report in language resembling a hit.

A result on a different fusion is not a result on this one. Immunogenicity studies on other fusion-driven tumours already exist and raise the prior that fusion junctions in general can be seen by T cells [17,18,20,21], and one of them shows a breakpoint reaching a T cell after processing from the endogenous transcript rather than from an injected peptide [23]. None establishes that this junction, on these alleles, is presented at usable abundance, and none of them measures abundance at all. The discriminating question to ask of any such report is whether its evidence concerns abundance, processing, or only a different fusion; the best of the reports available today reaches processing, and step 3 of Section 6.1 turns on abundance.

Robotic execution is not material access. A cloud laboratory with per-experiment pricing supplies robots and generic reagents, not an EMC line or organoid, and without that none of the tissue-gated items in Section 3 becomes runnable.

A better predictor is not a measurement. An improved class I or class II model would change the numbers in Section 2 and leave B2 where it is. The limit there is that nobody has looked, not that the looking-glass is imprecise.

A larger allele panel raises a figure without grounding it. Extending the panels can only move coverage upward, and a higher predicted-presentation figure is not evidence that any patient's tumour presents anything.

6.1 Order of the remaining questions

No dates are offered and none of the steps below is scheduled; what follows is an ordering, and the ordering is the content. Each step is cheaper than the one after it and each can close the route, so running them out of order spends the expensive ones on a question a cheap one would have settled.

- 1. **Defend the acceptance threshold, or record that it cannot be defended.** Computational, needs nothing but public data. Calibrate the cut against experimentally validated epitopes, and if no validated set restricted to fusion-junction peptides exists, that absence is itself the finding and Section 2.3's curve is the only honest report of coverage. Until this is done every figure in B1 is a point on a curve.
- 2. **Measure whether the lead peptides bind their alleles at all.** Four peptide-allele pairs carry every class I figure in this paper; Table 3 ranks them by what each costs if it fails. Synthesise them and test binding in a cell-based stabilisation assay, first pair first.

Table 3. The four peptide-allele pairs every class I figure rests on, ranked by what each costs if it fails. Coverage at risk is a leave-one-out difference — the pooled figure with all four presenting alleles minus the figure without this one — and not the allele's marginal contribution as it enters Section 2.3's curve, which is order-dependent and larger. Percentile is the call that put the allele in the set. Testing binding for these four is the only step in Section 6.1 that needs neither an EMC specimen nor a proteomics facility.

Table 3. The four peptide-allele pairs every class I figure rests on, ranked by what each costs if it fails				
Order	Peptide	Allele	Percentile	Coverage at risk
1	RGDMPCVQAQY	HLA-A*01:01	0.4061	9.9 pp
2	MPPPLRGDM	HLA-B*07:02	0.458	7.2 pp
3	NMPCVQAQY	HLA-B*15:01	0.3736	6.5 pp
4	NMPCVQAQY	HLA-A*30:02	0.4033	3.0 pp

- 3. **Look for the peptide on a tumour.** Immunopeptidomics on EMC tissue or a patient-derived line, with the limit of detection, the specimen count and their HLA types, and the abundance-matched positive control all fixed before the run, per B2. This is the decisive step and the first that requires material this programme cannot obtain.
- 4. **Only if step 3 is positive, ask whether a T cell sees it.** Recognition of the peptide-HLA complex by T cells from HLA-matched donors. A negative here is the one that B3 predicts and that nothing computational can rule out.
- 5. **Only then is there anything to build.** The construct in Section 2 is an output of a screen and is not a candidate for administration; it exists so that steps 2 to 4 have something specific to test.

Steps 1 and 2 would together decide whether the rest is worth commissioning, and neither needs an EMC specimen. That is the whole of what this paper can say about what to do next.

6.2 Conditions for revision

Four statements in this paper are falsifiable by an observation a reader could go and make, and they are listed so a future reader can check them rather than re-derive them. Sequence-level novelty for 170 of 174 peptides would be overturned by a proteome release adding a protein that contains one of the four strong binders. Every coverage figure would move upward if a wider panel added a presenting allele and downward if measured presentation removed one of the strong calls, and all of them go to zero if the acceptance threshold is set below 0.37. The class II statement said, in the version this paper supersedes, that it would be changed by a wider DR, DP and DQ panel — named there as the single most likely source of a change to it. That panel has since been run and it did change it, from no strong epitope on three alleles to one on 23; the statement in B4 is now the wider panel's, and what would change it again is measured class II presentation rather than more prediction. And the argument in Section 4 would be closed altogether by an EMC series showing HLA class I loss as protein; the transcript-level reading in B8 does not show it, and a protein-level series that did would end every antigen-directed route in this disease including this one.

7. Limitations

All binding figures are predictions from sequence-based models and no EMC-specific validation of either model exists. They predict peptide-MHC affinity alone: proteasomal cleavage and TAP transport are not modelled here, so a strong call bounds what could be presented rather than naming a peptide that is. Coverage is computed by multiplying non-carrier frequencies across alleles, which assumes independence both between loci and between alleles at the same locus. The same-locus case arises here — two alleles pooled are both HLA-B — and handling it correctly moves the pooled figure by about 0.3 percentage points. Cross-locus haplotype linkage disequilibrium is not modelled and its effect is not estimated; bounding it would need haplotype-frequency data rather than the allele frequencies used here. The population-to-region mapping is an approximation, regional figures are point values on samples as small as 579 individuals, and the frequencies are pooled across populations whose allele frequencies differ by more than the figures being reported. A fourth dependence sits underneath all of these and was not measured until now: every peptide in Section 2 is derived from one canonical transcript per gene, and across the 970 in-frame transcript pairs tested under B5 no junction peptide is common to all of

them. The seam residue itself takes nine values, so the peptide identities move with it. The counts are more stable than the identities — each pair yields 30, 34 or 38 novel peptides and no other value — but a reader should take the specific sequences named here, NMPCVQAQY included, as conditioned on the canonical annotation rather than as properties of the locus. Which transcript a patient's fusion actually uses is not decidable from annotation and is not decided here. The binder counts and every coverage figure derived from them depend on an acceptance threshold this paper does not defend, and no multiplicity correction is applied anywhere: 174 peptides were screened against 10 and then 34 alleles at two thresholds, with no decoy control and no null expectation, so the calls that pass are reported as what the screen returned rather than as an enrichment over chance. The near-self search of B3 is the one analysis here that carries a null, and it carries one because a count of near-self neighbours is meaningless without the number a peptide of that length and composition finds by chance; the binding screen still has none, and a shuffle null for predicted binding would need a defended threshold to be a null of anything.

The IMMUNOSARC II EMC cohort result is a conference abstract, single-arm and not peer reviewed, and it evaluates a combination whose component with the larger independent EMC evidence base is the tyrosine kinase inhibitor rather than the checkpoint inhibitor; it is cited to establish that a vehicle and a backbone exist, not to attribute activity to the immune arm. The single-agent checkpoint response cited beside it is one patient in an industry-authored conference abstract and bounds nothing. EMC's characterisation as immune-excluded rests on inference; its characterisation as immunologically quiet rests on the two archival cohorts read here against comparator sarcomas, which is a contrast and not an absolute, and on one twelve-tumour series read here only as an abstract. Neither instrument separates a cold tumour from an excluded one, which is limit B8 and the reason B6 and B7 remain graded apart. The clinical prior art for fusion-junction peptide vaccination is two small early-phase synovial sarcoma trials whose immunological readouts were weak and one single-patient Ewing sarcoma report [17,18,19,20]; it bears on whether such a construct can be administered and immunologically monitored, and on nothing about benefit in any disease. The class II panel is 23 alleles across DR, DP and DQ, its one strong call lands on a single uncommon allele, and class II prediction is less accurate than class I, so B4's positive bounds less than a class I call of the same nominal strength. No claim is made that any peptide is presented, that any construct would be immunogenic, that any combination would be safe or effective, or that any of this is ready for clinical use. No wet-laboratory work was performed, and the measurements this characterisation most needs require work this programme cannot carry out.

8. Methods and reproducibility

Every figure in Sections 2 and 3 is generated by a script in `research/modalities/` and is committed as a JSON artifact beside it: `fusion_breakpoints.py` for the junction set and predicted binders, `hla_coverage.py` for population coverage, `coverage_scan.py` for the coverage-versus-allele-count curve, `coverage_threshold_curve.py` for the coverage-versus-threshold curve of Section 2.3, `junction_proteome_novelty.py` for the proteome search of Section B5, `junction_frameshift_peptides.py` for the out-of-frame screen of Section 2.2, `junction_selfsimilarity.py` for the near-self search of Section B3, `predictor_concordance.py` for the second-predictor check, `coverage_uncertainty.py` for the within-locus form, the Fréchet bounds and the between-population spread of Section 2.3, `novelty_seam_test.py` for the one-residue pre-screen of Section B5 and its validation, `patient_neopeptides.py` and `patient_cd4_epitopes.py` for the per-patient shortlists, and `vaccine_construct.py` for the candidate construct and its minimal synthetic long peptide. The corresponding artifacts are `fusion-breakpoint-neoantigens.json`, `hla-coverage.json`, `coverage-curve.json`, `coverage-threshold-curve.json`, `epitope-allele-matrix.json`, `epitope-allele-loose-matrix.json`, `junction-proteome-novelty.json`, `junction-frameshift-peptides.json`, `junction-selfsimilarity.json`, `predictor-concordance.json`, `coverage-uncertainty.json`, `novelty-seam-test.json`, `patient-cd4-demo.json` and `vaccine-construct.json`. The two tables in this paper are generated from those artifacts by `vaccine_path_tables.py` and a build fails if a cell and its source diverge. The predictor versions those artifacts record are MHCflurry 2.1.4 with models release 2.2.0 for class I and MHCnuggets 2.4.1 for class II. MHCnuggets publishes no models-release identifier separate from its package version — it ships its trained weights inside the distribution — so the version is the whole of the provenance available for that arm, and the artifact records why rather than leaving the field blank.

Table 2. The 34-allele class I panel. Listed so the coverage scan can be reproduced and the choice of panel assessed rather than taken on trust. It is a common-allele HLA-A and HLA-B panel spanning global diversity, and it carries no HLA-C and no class II allele — so every figure computed on it is a partial coverage of the class I repertoire, and §2.3's statement that extending the panel can only raise the figures applies to HLA-C before it applies to anything else.

Table 2. The 34-allele class I panel	
Locus	Alleles
HLA-A (16)	HLA-A*01:01, HLA-A*02:01, HLA-A*02:03, HLA-A*02:06, HLA-A*03:01, HLA-A*11:01, HLA-A*23:01, HLA-A*24:02, HLA-A*26:01, HLA-A*30:01, HLA-A*30:02, HLA-A*31:01, HLA-A*32:01, HLA-A*33:01, HLA-A*68:01, HLA-A*68:02
HLA-B (18)	HLA-B*07:02, HLA-B*08:01, HLA-B*13:02, HLA-B*14:02, HLA-B*15:01, HLA-B*18:01, HLA-B*27:05, HLA-B*35:01, HLA-B*38:01, HLA-B*40:01, HLA-B*40:02, HLA-B*44:02, HLA-B*44:03, HLA-B*46:01, HLA-B*51:01, HLA-B*53:01, HLA-B*57:01, HLA-B*58:01

These are not regenerated on every commit, and this paper does not claim they are. The workflow that runs them is dispatched by hand, its steps do not fail the run when a generator fails, and it writes its outputs to a separate cache branch from which they are copied in. So the artifacts in the repository are the record of a run that happened, verifiable by their embedded timestamps and input hashes, rather than the output of continuous re-execution. An earlier version of this section said they were regenerated in continuous integration; that was not true of any branch a reader would fetch.

Clinical figures are quoted from the curated EMC registry at `research/data/emc-clinical-registry.json`, which carries a structured citation entry for every source and a retrieval note for those whose identification required one.

Withdrawn statement	Why it does not stand
"presented on HLA-B*15:01 alone", stated without naming a panel	True of the ten-allele screen only. The 34-allele screen finds the same lead peptide strong on HLA-A*30:02, and the junction's coverage is then 12.3% rather than 8.5%.
The 30.4% figure described as a "ceiling"	Sections 5 and 6 of the same manuscript said extending the panel could only raise it. It is a union over four alleles at one threshold, and moving that threshold to 0.37 takes it to zero.
"It does not attain 50% at any panel size" — withdrawn	No search over panel sizes was performed; the panel is fixed at 34 and the scanned variable is the number of presenting alleles. In Northern Europe two alleles reach 52.8%.
Wilson 95% confidence intervals on every coverage figure	They were computed by pooling every reference population into one binomial. The same records show the frequency of one pooled allele ranging from 0 to 0.40 between those populations, so the model the interval assumes is refuted by its own input.
"the combined CD8 and CD4 figure is null rather than unreported. That is a result" — withdrawn	The instrument's class II branch never evaluates when no allele qualifies, so the empty field records "not computed". The instrument's own note calls the class II coverage it would compute a floor that untested alleles could only raise.
"The one substantive reasoning correction this work offers ...", and the symmetry it asserted	The quoted exclusions are this author's own route-ledger notes, not positions in the literature, and the vaccine's own entry in that ledger is parked on immunogenicity rather than on absent priming. The claim is narrowed in B6 and the Abstract accordingly.
The dated capability bands of the former Section 3.1 (2027H2 / 2029 / 2032)	Withdrawn entire. They were self-described as the weakest material in the paper, and a forecast a reader cannot check does not become checkable by being tabulated.
"regenerated in continuous integration", of the artifacts in Section 8	The workflow is dispatched by hand, its steps do not fail the run when a generator fails, and it writes to a cache branch. Section 8 now says what is actually true.
Reference 8 quoted as "correlated with disease outcome"	The study's title names immune response in placenta and colorectal cancer, which is the tissue setting it examined.
References 10 and 11 as "[citation to verify]"	Both were resolvable in this repository at no cost and are now written out. The one reference that genuinely has no record — the class II predictor — says so in its own entry instead.
"24 patients across nine centres ... gives a realistic accrual rate"	Unfiltered by this paper's own eligibility criteria. Applying them gives roughly 0.3 to 1.4 eligible patients a year.

Appendix C. Statements withdrawn after the first adversarial review

The review recorded in Appendix B was not the last correction. The statements below stood after it and do not stand now, and they are recorded on the same principle: a correction that leaves no trace is indistinguishable from a claim that was never made.

Withdrawn statement	Why it does not stand
B8, "rather than from published EMC-specific immune profiling"	Such profiling was published on 2026-07-13, four weeks before this manuscript's date, on the twelve-tumour cohort described in B8's own revision.
B8, HLA class I expression status "is not known for this disease"	The antigen-presentation precondition — β 2-microglobulin, the class I heavy chains, the peptide transporters, tapasin, NLRC5, the immunoproteasome subunits and ERAP1 — had been read on both EMC array platforms on 2026-08-09, in this author's own artifacts, before the sentence was written.
B6 and B8 read together, that whether EMC is immunologically quiet was unmeasured here	It is now measured in the two archival cohorts, and the direction is that EMC is quieter than the comparator sarcomas on the interferon axis, the myeloid compartment and class II presentation. That is a cohort contrast against other sarcomas and not a statement that the disease is cold; the distinction is stated in B8.
Section 1, "no published EMC immune profiling" listed among the limits of access	The same statement as B8's, in a second place, and false for the same reason. It is the one-of-a-pair defect: correcting B8 alone would have left the claim standing where a reader meets it first.
Section 7, "The characterisations of EMC as cold and as immune-excluded rest on inference rather than on published EMC-specific immune profiling"	Third home of the same claim. The exclusion half still rests on inference; the quiet half no longer does.
Section 7, "The class II panel is three DR alleles with no DP or DQ, so its negative bounds a narrow question"	Superseded by the panel widening reported in B4 and in Appendix A, and stale in this section since that widening. Section 7 was contradicting B4 inside the same document: the panel is 23 alleles and the arm is not negative. This was not found by any adversarial round; it is a wording that no numeric guard reads.
B7, "No EMC-specific expression evidence for the pathway is cited, because none is known to the author"	The chondroitin-sulfate biosynthetic and sulfation machinery had been read on both EMC array platforms in this author's own artifacts before the sentence was written. The reading is reported in B7 and does not support the analogy the sentence was protecting, which is a different correction from the one the sentence claimed to make.
B2, that the reports available "do not close this for any fusion", because they measure responses to peptides that were administered	A study of 34 paediatric acute leukaemias recovered fusion-breakpoint-specific T-cell receptors from co-culture with autologous blasts, with no peptide administered, and mapped their restrictions on patient-matched HLA. Endogenous processing and presentation of a fusion breakpoint is shown; presentation on an EMC cell, and abundance anywhere, are not.
B4, that the class II arm is the lesser of the two	Two of three fusion-specific restrictions recovered from endogenously presented breakpoints were class II, and the one reported de novo vaccine response was CD4+. Three observations are a pattern to test, and B4 now asserts neither ranking; what remains ordered is the accuracy of the two predictors.
Section 6, that no report on another fusion "measured presentation at all"	The same study measures processing and presentation, though not abundance. The bullet now separates the two, because step 3 of Section 6.1 turns on abundance.
The B6 and B7 rows of Section 3, cost to move "not movable by any computation"	True of each limit and false of the evidence bearing on it: a computation over public archival series moved what is known about both. Those cells now name what the remaining move costs, which is spatial profiling on tissue.