
Coastal Processes

Tristan Salles

Jan 11, 2023

CONTENTS

1 Ocean Data Query	3
1.1 Coastal Ocean Observing Systems	3
1.2 Why do we need ocean observing tools?	6
1.3 National & Global Ocean Observing Infrastructure	6
1.4 Access to ocean data	9
1.5 Modeling & Analysis	9
1.6 Summary	11
2 Analysing Wave Climate	13
2.1 Wave Climate from the Australian Bureau of Meteorology.	13
2.2 Characterising wave climate	16
2.3 Extreme wave climate	20
2.4 Wave climate variability	23
2.5 Hands-on examples	26
3 Bibliography	29
3.1 References	29

Author: Tristan Salles



THE UNIVERSITY OF
SYDNEY

Course content

For this component of the unit, we will use:

1. a *learning website* where you will find all information relative to the lectures themselves: it contains some **notes**, **videos** and **exercises** for each weeks,
 2. a *web-based programming platform* based on **Jupyter notebooks** that contains all the materials for the **exercises** and **practicals**.
-

This part of the course focuses on coastal systems analysis using both ocean data from a variety of sources (hindcast and forecast data, satellite observations, radar, moored buoys) and numerical models of shoreline changes, wave dynamics and coral reef evolution.

The emphasise will be on:

- **How to query and analyse numerical and ocean data?**
 - **What can we learn from ocean and wave models on coastal evolution?**
 - **How to use these models and what are their limitations?**
-

Note: Today, numerical modelling and ocean data query techniques are routinely applied by governmental agencies, companies and research organisations to tackle complex coastal problems. It is based on advanced physical models and engineering approaches designed to describe and observe the connections between ocean dynamics and coastal evolution.

Over the last 20 years, major improvements in our understanding of coastal processes have been related to (1) easier access to observational dataset and (2) improved coastal numerical models. **It is now critical for any graduates to have a sense of how to access these dataset efficiently, how to process them using simple coding tools and how to interpret them.**

During this course, we will learn how to do that using web-enabled open source technologies that you will be able to reuse in the future.

Lectures content

Important: We will do **computer-based exercises during both lectures and practicals** so you will need to bring an electronic device (preferably a laptop but a tablet with internet access could work). As there will be a lot to cover, it is also recommended to go through each chapter prior to the lecture to be well prepared and able to get the most of it during classes.

- *Ocean Data Query*
- *Analysing Wave Climate*
- *Bibliography*

CHAPTER ONE

OCEAN DATA QUERY

1.1 Coastal Ocean Observing Systems

Note: Now, more than ever, there is a need for regional to global observing systems that can provide accurate real-time data and forecasts on coastal ocean conditions.

The ocean plays a role in everyone's life. It affects weather and climate patterns around the globe, hosts an abundance of wildlife that support fishing industries and provide food for the world, serves as a highway for vessels that deliver everyday materials, and supports economies as a tourism destination.

The coastal ocean is the part of the earth system where land, water, air, and people meet together. Populations, businesses, and infrastructure are increasing along coastlines, which are all susceptible to changing coastal ocean conditions.

Coastal ocean observing systems (COOS) are necessary for advancing our understanding on the state of the coastal ocean worldwide and its impact on matters of societal importance. These systems integrate a network of people, organisations, technologies, and data to share advances, improve research capabilities, and provide decision-makers with access to information and scientific interpretations.

1.1.1 Data acquisition



Important: Data, observations, and models integrated into the COOS come from a variety of platforms, including, for example, **moorings**, high-frequency (HF) **radars**, underwater **gliders** and profilers, **satellites**, and **ships**. The resulting data are used to better understand, respond to, and prepare for **short-term** events such as oil spills, harmful algal blooms, and fish kills, **longer term** changes in our oceans resulting in acidification, hypoxia, and sea level rise, and in everyday decisions related to maritime operations, public health, and management of healthy ecosystems.

1.1.2 A bit of history

Matthew Fontaine Maury (1806/1873) was an American astronomer, Confederate Navy officer, historian, oceanographer, meteorologist, cartographer, author, geologist, and educator.

He was nicknamed **Pathfinder of the Seas** and **Father of Modern Oceanography and Naval Meteorology** and later, **Scientist of the Seas** for his extensive works in his books, especially *The Physical Geography of the Sea* (1855), the first such extensive and comprehensive book on oceanography to be published. *Maury made many important new contributions to charting winds and ocean currents, including ocean lanes for passing ships at sea.*


As a sailor, Maury noted that there were numerous lessons that had been learned by ship masters about the effects of adverse winds and drift currents on the path of a ship. The captains recorded the lessons faithfully in their logbooks, but they were then forgotten. At the US Naval Observatory, Maury uncovered an enormous collection of thousands of old ships' logs and charts in storage in trunks dating back to the start of the US Navy. He pored over the documents to collect information on winds, calms, and currents for all seas in all seasons. His dream was to put that information in the hands of all captains.

Maury became convinced that adequate scientific knowledge of the sea could be obtained only by international cooperation. He proposed for the United States to invite the maritime nations of the world to a conference to establish a **universal system** of meteorology, and he was the leading spirit of a pioneer scientific conference when it met in *Brussels in 1853*.

As a result of the Brussels Conference, a large number of nations, including many traditional enemies, agreed to cooperate in the sharing of land and sea weather data using uniform standards. It was soon after the Brussels conference that Prussia, Spain, Sardinia, the Free City of Hamburg, the Republic of Bremen, Chile, Austria, and Brazil, and others agreed to joined the enterprise.

Within a few years, nations owning three fourths of the shipping of the world were sending their oceanographic observations to Maury at the Naval Observatory, where the information was evaluated and the results given worldwide distribution.

Note: Maury's idea set the scene for what is now a *Global Ocean Observing Infrastructure!* Over the 19th and early 20th century a lot of measurements were made by diverse communities for their **own needs** (scientists, fishermen, commercial navigators...). However as more and more data was collected by divers communities, data was shared only among small communities and was not properly archived and it wasn't done in a organised way. In situ archeology is a hard job providing questionable databases!



1.2 Why do we need ocean observing tools?

Until recently, the ocean was viewed as *being an unlimited resource to be exploited* (source: *US National Academies of Science report on Economic Benefits of Oceanographic Research, 1964*). Subsequent science, including that from national and international observing efforts, has changed what we know about the ocean.

Nowadays, we recognise that **the ocean is a finite and shared resource** that needs to be managed regionally, nationally, and worldwide.

Societal threats and challenges

There are a number of significant societal threats and challenges facing humans due to the changing ocean:

- The increased frequency and intensity of coastal storms and resulting storm surges will affect our coastal communities and disrupt commerce, nationwide.
- Sea level changes are threatening critical infrastructure worldwide.
- Harmful algal blooms and oxygen-deficient dead zones threaten water supplies, fisheries, and coastal recreation.
- Ocean acidification is negatively impacting coral reefs and shellfish harvesting.
- The increasing size and number of vessels calling on ports present challenges for our already inadequate maritime infrastructure, and pose potential environmental risks as well.
- The world's growing population will increasingly rely on the ocean for food, but fishing must be done sustainably.

To be able to understand and manage the ocean, we need meaningful measures of the ocean's state.

1.3 National & Global Ocean Observing Infrastructure

Observing systems are expensive; Australia invests billions of dollars in civil Earth observations to ensure that the nation's decision-makers and managers have the information they need about climate and weather, disaster events, land-use change, ecosystem health, natural resources, and many other characteristics of the planet.

The ocean is a harsh environment in which to operate an observing system, from corrosion due to salinity to bio-fouling. The costs of maintaining instrumentation in this environment, accessing remote locations, and establishing sufficient communications with deployed technologies are significant.

IMOS had as its prime focus improving scientific understanding of ocean conditions, but the information it generates is increasingly being used by government agencies and other users to inform decisions. The main activities of **IMOS** are based around:

- Deploying, maintaining, and developing advanced observations technologies
- Providing free and open access to data in support of a wide range of users
- Advancing modeling
- Focusing on education

In addition to providing observations and data to a large and growing number of research projects, student projects, and academic courses, **IMOS** is now recognised as an essential partner in large, multi-institutional research programs across multiple sectors. It has contributed to 180 postgraduate projects, over 400 journal publications, and 250 research projects.

1.3.1 Coastal ocean observing systems development

Most large-scale coastal ocean observing systems are funded through national governments for their own interests, often with different foci, but the world's oceans are connected, therefore partnering is the key to success. As an example, the U.S. Integrated Ocean Observing System ([IOOS](#)), **Australia's Integrated Marine Observing System (IMOS)**, or European Ocean Observing System ([EOOS](#)) are progressing in their respective regions and are working together to observe and compile ocean information in a way that is easily accessible to scientists and managers.

IMOS advances have benefited from an evolving set of ocean observing efforts. The envisioned concept was a coordinated national and international network of observations, data management, and analyses that systematically acquired and disseminated data and information on past, present, and future states of the oceans. The coastal ocean observing efforts are implemented via regional programs distributed around coastal regions.

Each program is designed to assess and predict the effects of weather, climate, and human activities on the state of the coastal ocean, its ecosystems and living resources, and on the world's economy.


Note: The Australian [IMOS](#) was formed in 2007, with equipment deployed from the next year onward. Scientific 'nodes' were formed broadly around state boundaries with both nationally unified overarching science goals and local priorities. The **NSW-IMOS** is an example of a successfully implemented ocean observing system along the coast of southeastern Australia. The current observational array is designed around pertinent science questions, leveraged existing data streams, and opportunities for further oceanographic research.

[IMOS](#) currently has a portfolio of **13 Facilities** that undertake systematic and sustained observing of Australia's marine environment, across scales (from open ocean, onto the continental shelf, and into the coast), and across disciplines (physics, biogeochemistry, and biology and ecosystems).

1.3.2 Ocean observing technologies

It is a complex task to measure the ocean in ways that deliver useful products for people. For example, to deliver a five-day weather forecast for any local region, meteorologists must sample the whole planet. **Satellites** are key tools that provide multispectral images, atmospheric soundings, and sea surface characteristics needed for these forecasts. While satellites enable us to see through the atmosphere, they can only measure the surface of the ocean, and this does not provide the detail needed for accurate weather prediction. Therefore, we also need a complement of *in situ* measurements that extend our reach to the depths of the oceans at all relevant scales of phenomena. *In situ* refers to systems that measure on or under the surface of the ocean in continuous and event-driven modes, automatically and by humans.

[IMOS](#) is a global leader in ocean observation. It has infrastructures that operate on continental scales, field observing technologies, shares best practices for a broad suite of variables, and generate masses of data. [IMOS](#) is partner with [IOOS](#) and [OTN](#) (this later is a global ocean research and technology development platform, in the area of animal tagging and tracking). To understand animal movements and survival, you also have to understand how the animals respond to changes in environmental conditions. Thus, the animal tracking effort is tightly integrated with the observation of the physical and chemical environment, and all networks depend on common platforms for oceanographic observations. Other observing technologies include Argo, moored buoys, high-frequency radar (HFR), and autonomous underwater vehicles known as gliders.



1.4 Access to ocean data

All the national systems that participate in the Global Ocean Observing System (**GOOS**) and the Group on Earth Observations (**GEO**) subscribe to the principal of **free and open data**. It is **policy** within Australia that all observational data collected with governmental funding support are made freely and openly available, which makes sense from scientific, disaster response, and economic perspectives.

Important: One challenge **IMOS** faces is the need to support operational systems, often with unique data formats, while simultaneously working to create mutually compatible data access systems and services, and common data formats and metadata standards, in order to facilitate access to this public resource through the internet.

IMOS provides marine data such as *temperature, salinity, currents, wind speed/direction, waves*, and other primarily physical observations for model assimilation through the GTS (Global Telecommunication System). For broader access, **IMOS** uses three standards to convey the information in an interoperable manner:

- Open-source Project for a Network Data Access Protocol (**OPeNDAP**),
- Sensor Observation Service (**SOS**) and Open Geospatial Consortium (**OGC**) Standard, and
- Web Map Services (**WMS**) OGC Standard.

1.5 Modeling & Analysis

To access the Notebooks exercises using the following link:



Prediction of future conditions is critical to deliver the full benefits of an ocean observing system. The Australian ocean and coastal modeling community is a partnership between multiple universities and national organisations (CSIRO, IMOS, GA, BOM to cite a few) that is being asked to provide greater resolution models that cover not only traditional physical

water circulation, which remains a critical need, but also expand outputs to include inundation forecasting and ecosystem modeling.

In May 2014, IMOS published the **IMOS Strategy 2015-25**. In this plan, one can read that:

Note: IMOS has a concerted focus on making **data available** and **seeing that it is used**. IMOS has worked with the coastal modeling community to make sure that available data are informing numerical models. Going forward, IMOS will use its infrastructure to be a coordinating entity to advance the assimilation and further development of coastal modeling. IMOS has begun a structured engagement with the coastal and ocean modeling communities through development of joint products (e.g., in ocean reanalysis), national workshops, and targeted infrastructure investment at the model-data interface (such as virtual laboratories).

1.5.1 Hands-on examples

As we just saw, access to quality data is **essential to understand marine processes**.

Over the last 20 years, **ocean data portals** have emerged and are routinely used to better understand the complexity of the ocean and its interactions with climate and life. These portals facilitate seamless access to marine data/services and promote the exchange and dissemination of ocean-related information.

Important: The information that is stored, processed, and exchanged, is at the heart of modern marine science. Ocean scientists routinely perform *data crunching* to understand a particular system and need to *access* and *query* extensive lists of dataset. **Understanding how these data are stored, their origin and how to quickly retrieve particular information from them are crucial skills that you will need in your job!**

Wave height measurements taken every day by a buoy offshore Sydney are data. A graph showing the evolution of the significant wave height over time, at a given place, is information. The fact that the number of extreme storms hitting Australian's coast increases as a result of climate change is knowledge. These three notions are very closely linked.

1.5.2 Exercises for the lecture

Roughly speaking, here is how you should use them:

- A piece of data provides a basic description, typically numerical for our purposes, of a given reality.
- Drawing on the collected data, information is obtained by organising and structuring data so as to derive meaning.
- By understanding the meaning of information, we obtain knowledge.

Note: One of the great challenges for Ocean Data users is to understand **where** and **how** to find technologies that make it possible to evaluate, validate, verify, and rank information to help them in their jobs. This involves understanding how the ocean data providers are organised, the main standards, vocabularies and formats which are used by the community as well as the best approach for accessing and querying these information routinely.

Important: Before starting the exercise, take some times to familiarise yourself with the environment. You will see in the `IntroNotebook` ipython notebook in your main repository a link to an introduction notebook that I highly recommend to do!

After following the video and the introduction notebook, you will open the workspace in **binder** and from the bottom of the IntroNotebook notebook you will click on the **Ocean Data Query** link. It will open a folder containing exercises and practicals:

- OceanData1.ipynb and OceanData2.ipynb are 2 exercises that complement what you learned during this lecture,
- Oceanforecast.ipynb and Waverider.ipynb are Python notebooks that you will use for your practicals.

Loading and checking IMOS NetCDF dataset

Exercise 1.1

In this first example (OceanData1.ipynb), we will work with the **IMOS Portal** using **Python** via **Jupyter Notebooks**. There are several advantages of using Python as a general data analysis language and the notebook environment is a versatile tool that is designed to be interactive, user-friendly, open-source and sharable.

We will see how to load NetCDF data into a Python environment, and show how to use the data once loaded.

Querying and analysis Coastal Ocean Radar dataset

Exercise 1.2

In this second example (OceanData2.ipynb), the Australian Coastal Ocean Radar Network (**ACORN**) facility comprises a coordinated network of HF radars delivering real-time ocean surface velocity data. We will export the dataset file (**NetCDF**) for a given region and then we will plot the velocity field at a given time in a latitude/longitude grid.

1.5.3 Exercises for the Practical

You will run a series of examples that will introduce some aspects of marine data querying.

- Offshore Sydney wave buoy data from Australian Integrated Marine Observing System (IMOS) and
- Different dataset from the Bureau of Meteorology, CSIRO as well as forecast model outputs from NOAA.
- Extract Ocean Radar dataset for Turquoise Bay from IMOS and plot them on a map.

Download the practical documentation from Canvas website and answer the questions using: the `Waverider.ipynb` and `Oceanforecast.ipynb` notebook.

1.6 Summary

Working together, agencies like **IMOS** or **IOOS** help to manage our oceans by measuring them and connecting observations to people. Yet, ocean observing systems are only as good as our ability to observe and accurately model ocean systems.

Finding the resources to sustain coastal observations over relevant time scales is a big challenge.

Important: As coastal data users - we need to understand how ocean dataset are obtained, managed and stored. We also need to know how to efficiently load these dataset, query and analyse them.

As a growing population continues to stress our planet, quality observations will increase in importance. But it is not good enough to measure the planet, we also need to work to ensure this information comes into play in our communities, our economies, and in management decisions.

CHAPTER
TWO

ANALYSING WAVE CLIMATE

Overview of this chapter content

Waves are a defining characteristic of water-bodies worldwide, transporting energy that impacts numerous physical and ecological processes. Since World War II, there has been a substantial increase in scientific understanding of wave formation, propagation and resulting impacts on coastal and marine features.

Initial wave studies were restricted to data from voluntary ship observations, which were limited in accuracy and sample size, particularly during extreme conditions. The development of technology such as wave buoys has provided accurate, hourly data of wave height, period and direction. This data enables almost immediate knowledge of the **sea state**, which is the present wave conditions.


2.1 Wave Climate from the Australian Bureau of Meteorology.

Wave records in Australia

In Australia, waves have been officially recorded since 1974 with the deployment of the first Waverider buoy in Port Kembla, New South Wales (Hemer et al. 2007). This was the start of an increasing network of *buoys*, *High-Frequency radars* and other wave measurement technology leading to significant knowledge of wave dynamics around the Australian coastline. A further technological leap in wave monitoring was achieved in 1985, with the launch of the **GEOSAT** satellites with a mounted *altimeter*. As a remote sensing technique, altimeters provide wave data observations with unparalleled spatial and temporal resolution (Ribal & Young 2019). Long-term analysis of wave conditions can lead to the characterisation of a wave climate, which is the modal wave conditions of an area (Godoi et al. 2016).

Variation from the modal wave climate can occur due to *extreme events* such as large **storms**, **Tropical Cyclones** (TC) or **tsunamis**. For Australia, it is critical to understand these extreme conditions and to analyse their role in ecological disturbance, as well as to predict present and future vulnerability, particularly with anthropogenic climate change likely to increase the frequency of intense TCs along our coasts.

The wave climate of a region is further impacted by **climate oscillations**, which are semi-regular cyclical changes that have well-defined effects on regional and global weather patterns. These fluctuations typically influence atmospheric temperature, sea surface temperature, wind or precipitation. Oscillations operate on different timescales and can therefore occur together, enhancing or diminishing the effects of each event (Godoi et al. 2019). A key example is the El Nino-Southern Oscillation (**ENSO**), which plays an integral role in climate modulation over the Pacific Ocean and surrounding continents. Both the Australian modal and extreme wave climates can be substantially influenced by these climate oscillations, primarily due to changes in wind patterns and changes to extreme event frequency and intensity.



2.1.1 Wave formation

Ocean waves are formed by different mechanisms and as a result, can have a wide range of properties. In this course, we focus on **ocean surface gravity waves** (henceforth referred to as waves), also known as *wind-generated waves*.



Important: At the atmosphere-ocean interface, kinetic wind energy is transferred to the ocean, which with sufficient wind, generate waves (Holthuijsen 2007). Several factors influence wave formation, including the *strength* and *duration* of the wind, and the *fetch*, the distance over open ocean which the wind blows.

When wind blows over only a short distance, wind-waves are formed. In Australia, these locally-generated waves typically have a short period of **1-8 s**, and travel slower than the prevailing wind. In contrast, swell-waves are generated by distant storms and have propagated out of the wave-generation area. Swell has a longer period of **8-30 s**, and is only minimally influenced by local wind conditions. In many instances, these two wave types occur simultaneously and can travel in opposite directions, creating chaotic seas.

2.1.2 Wave Parameters

Waves can be characterised by parameters such as *wave period*, *height*, *power* and *direction*. Distinctions between different wave classes are based on these measurements and are used to describe both individual waves and the wave climate of a region.

The vertical distance between the maximum and minimum surface elevation over one wave period is referred to as **wave height** (Holthuijsen 2007). To define the overall wave height of many waves, typically the significant wave height is calculated (H_s). Mathematically, H_s is the mean of the highest third of waves in a given sample period (Holthuijsen 2007); traditionally, it is the height a trained observer would see when attempting to estimate average wave height, as humans would not see or consider the smallest waves.




Wave period is the time for one full wavelength to pass a given point, from the beginning to the end of the wave (Holthuijsen 2007). Usually, the zero-crossing period is used (T_z) that measures the time for a wave to go below the mean elevation, rise above, then return to mean elevation.

As waves propagate across the ocean surface, energy is also transferred, referred to as **wave power** (P) or **wave energy flux**. The P of each wave is proportional to the T_z and the square of H_s , with higher values producing more powerful waves (Airy 1841). This parameter is important as it combines both H_s and T_z , providing an overall understanding of the wave conditions and the energy it transports. For example, a wave with a H_s of 1 m and T_z of 5 s will have less power than a wave with an H_s of 1m and T_z of 8 s.

Propagation direction (θ) is the direction from which waves are coming from. In certain regions a particular wave direction can indicate different swell characteristics, for example, large waves from the South-East in Queensland, Australia, indicate strong Southern Ocean swell or Extra-Tropical Cyclone swell, whilst large waves from the north indicate Tropical Cyclone waves.

2.2 Characterising wave climate

The modal wave climate of a region is determined through a long-term analysis of several wave parameters, including H_s , T_z , P and θ .



There are numerous techniques that can be used to measure these parameters. However, most methods do not measure all parameters: a deliberate choice must be made, with consideration to the advantages and disadvantages of each technique. Frequently, the method chosen is customarily calibrated with other techniques or used in combination.

Common methods include:

- wave buoys,
- wave hindcast models and
- satellite altimeters.

2.2.1 Wave buoys

Wave buoy data is routinely considered ground truth, as it is a physical measurement of individual waves instead of remote sensing (Hemer et al. 2007). Buoys also provide the most reliable and accurate records, particularly for extreme values. Despite this, variations in sampling, calibration and computational methods can lead to significant errors by both over and under-estimating wave parameters.

Note: Bender et al. (2010) revealed buoys can overestimate H_s by 26%, and overestimate during hurricane peak by up to 56%, leading to significant and highly misleading errors.

Buoys are also spatially limited, measuring only waves that propagate directly through the site, leading to a restricted understanding of regional wave climates. Furthermore, many locations do not have buoys in operation, thereby inhibiting the analysis of wave climate through buoys alone.

2.2.2 Wave hindcast models

Wave hindcast models use **reanalysed wind fields** to investigate **past** waves. Wind speed and duration are required parameters that are calculated alongside fetch and water depth to determine H_s , T_z and θ . This provides data sets that are used globally to establish wave climate and can evaluate conditions at different temporal and spatial scales.


Tip: A widely used wave hindcast is the *National Oceanic and Atmospheric Administration* (NOAA) **WaveWatch III model** (Tolman 2009) as shown in the introduction to this lecture.

However, significant problems can arise through error with forced wind condition. For example, extreme events such as TCs can cause abrupt, localised changes to wind speed and direction. Since models such as **WaveWatch III** have relatively coarse spatial-temporal resolution, insufficient energy from wind is inputted leading to significant underestimation of wave conditions during these extreme events. The accuracy of the hindcast model is also dependent of additional source term parameters and generation, propagation and dissipation equations.

Note: In Australia, this is a significant problem in reef areas, where wave propagation is modified substantially by reefs. With coarse spatial detail, identification and accounting for reefs are reduced, thus leading to potentially incorrect wave conditions in areas sheltered by reefs.

Therefore, whilst wave hindcast models can provide excellent information of wave conditions, particularly in areas where there are no buoys, inherent limitations regarding spatial detail and reliance on source terms means that it is currently unsuitable for wave climate characterisation for reef or areas with complicated bathymetry.

2.2.3 Wave-sensing technology



A remarkable wave-sensing technology with high spatial-temporal density is the radar altimeter, widely established as a pinnacle remote sensing technique to determine wave climates globally (Ribal & Young 2019).


Radar altimeters are mounted on satellites and measure a footprint of the ocean directly under the satellite, between 5-7 km wide. When the water surface is calm and flat, the reflection of the radar pulse back to the altimeter is almost instantaneous. In contrast, when waves are present the pulse is first reflected at the crest of the wave, then progressively reflected as the pulse reaches the wave trough.

Important: The altimeter interprets this signal to determine wave height and wind speed. Therefore, this remote sensing technique **does not provide individual wave statistics**, but rather **returns the average value over the footprint**, up to 7 km wide.

In comparison to a buoy, altimeters provide excellent spatial coverage, with observations along the ground track every second, approximately every 5-7 km. Since the launch of the first altimeter in 1985, (GEOSAT), altimeters have been used to determine wave climate. With the increased number of altimeters in orbit, the global coverage and temporal density are increasing, resulting in a technology highly suited to characterising wave climate, particularly in areas with no buoys, complicated bathymetry and in remote locations.

Example of wave analyse from satellites measurements

Visualising Australian Ocean Surface Wave by extracting information from synthetic aperture radar (**SAR**) satellites with a **Jupyter Notebook**. **Demo in BinderHub**.



Wave period

For waves that have been locally generated by wind, the local wind speed and wave height, both of which are accurately measured by altimeters, can be used to determine T_z . More accurate results are achieved when the *wave age* is calculated first, which is the length of time wind has been acting on a wave.

First, the wave age is calculated through:

$$\epsilon = 3.25 \frac{H_s g}{U_{10}}$$

where H_s is the significant wave height, g the acceleration by gravity and U_{10} the wind speed. Wave age ϵ can then be used to estimate T_z :

$$T_z = (((\epsilon - 5.78)/(\epsilon + (U_{10}/H_s \times (U_{10}/H_s) + H_s)))) + (H_s + 5.70)$$

Wave energy, group velocity & power

Mean wave energy density (E) (J/m^2) is calculated by:

$$E = \frac{1}{8}\rho g H_s^2$$

with ρ the density of seawater (set to 1027 kg/m^3). Wave group velocity (C_g) in deep water conditions is approximated with:

$$C_g = \frac{g T_z}{2\pi}$$

And wave power P can, therefore, be estimated through:


$$P = E C_g$$

which is the wave energy flux per metre of wave-crest (W/m).

2.3 Extreme wave climate

Globally, extreme waves are generated by:

- cyclones,
- tsunamis,
- rogue waves and
- large storms.




Above figure depicts the estimated 1-in-100 year significant wave height globally, with larger waves predominate in the higher latitudes due to extended fetch and frequent storms, whilst equatorial regions experience smaller extreme waves (Takbash & Young 2019).

Cyclones are a broad category of weather systems that can cause extreme waves, characterised by strong winds around a low pressure centre. TCs are formed over tropical or sub-tropical regions, whilst Extra-Tropical Cyclones form in the mid- or high-latitudes.

Note: **East Coast Lows (ECL)** are one such Extra-Tropical Cyclone that is generated near southeastern Australia, from either tropical or mid-latitude controls (Dowdy et al. 2019).


Tropical cyclones (TC) are low pressure systems that periodically develop over warm oceans. Sea surface temperature is a key driver of TCs, with a positive temperature anomaly leading to increased intensity and kinetic energy of the TC, as well as enhanced storm surges.

Strong winds and heavy rain can extend hundreds of kilometres from TC centres, and can last from days to weeks. The consequent destructive wind, rainfall, waves and storm surges are highly dangerous, causing extensive risk to life and millions of dollars in property damage. For example, severe TC Justin caused the death of over thirty people due to storm surges and large waves (source: *Bureau of Meteorology*).



TCs are major drivers of coral reef disturbance and destruction, with reefs close to a cyclone experiencing a decrease in hard coral cover, taxonomic richness and coral density that can last for decades. Lagoon flooding, increased sediment load, decrease in local salinity and pH levels are further impacts which negatively affect coral growth and recovery.

Note: Along with severe storms, cyclone waves are key producers of coral rubble. Together with carbonate sand, coral rubble constitutes the majority of reef volume. Over time and subsequent extreme events, the coral rubble progrades and can form rubble spits and islands. Despite the catastrophic nature of these impacts, TC are spatially bound, and thus reefs will typically go a number of years before another major disturbance tracks through the same area.



ECLs are a different type of cyclone that has significant effects on the Eastern Australian coastline. Formed by a temperature gradient between cold air in the upper atmosphere and warm Tasman Sea air, ECLs generate extreme winds, precipitation and large waves (Dowdy et al. 2019). They typically develop in the winter months close to the New South Wales coast, however, impacts can spread to southern Queensland and Victoria. Whilst typically not as severe as TCs, gusts over 170 km/hr and waves of over 14 m have been recorded, resulting in dangerous maritime conditions and coastal destruction.

2.4 Wave climate variability

Seminar on global wind & wave climate by Young et al. (2019)

The broadcasted seminar below investigates: global ocean wind and wave climate and ocean extremes. It describes changes in ocean winds and waves over the last 30 years and projections for future changes out to 2100. It also looks at projections for sea-level rise and the role waves play in determining coastal flooding. The results presented use measurements from a unique dataset of more than 20 satellite missions which have been combined to produce a single long-term global database of wind speed and wave height.

Various meteorologically driven changes in atmosphere-ocean coupling can substantially alter surface wind fields and, as a result, influence wave climate on both regional and ocean basin scales (Godoi et al. 2016).

Many studies have shown that the Pacific Ocean wave climate is altered by the **El Nino Southern Oscillation** (ENSO) and the **Southern Annular Mode** (SAM), however, the response to these can be variable in both space and time. Several regions, including islands in the Pacific, are projected to be more at risk from a changing wave climate than risks from sea level (Hemer et al. 2011). Thus, an understanding of the current variability of the wave climate is an important step to understanding the influence of global climate processes and potential links to climate change (Godoi et al. 2018).

2.4.1 Seasonal changes

Along eastern Australia, the strength of south-east trade winds fluctuate throughout the year and occur predominantly during the austral winter during *April–October* (dry season). From *November–March* (wet season) the trade winds lessen and can even reverse, linked to the Australian Monsoon and the location of the **Intertropical Convergence Zone** (ITCZ) (Hemer et al. 2011). During the wet season, the ITCZ moves closer to Australia, decreasing the strength of trade winds. Periodically, increased strength north-west winds occur due to the presence of the Australian Monsoon. This could have a significant impact on wave climate, with **decreased modal wave heights** during the wet season, however with **small periods of higher waves** due to intense storm and precipitation events during monsoons.


2.4.2 Long-term trends

Several regions of the global ocean have statistically significant positive and negative trends in H_s . An analysis of 33 years of altimeter data by Young & Ribal (2019) (see figure above) determined that extreme 90th percentile waves in the Southern Ocean are increasing by 1 cm/year, and in the North Atlantic by 0.8 cm/year. This is often correlated to an increase in extreme wind strength in the region.

Note: However we can see some differences, as a matter of fact, despite mean local wind speed increasing globally, the mean H_s show less explicit trends in all regions. Several areas, such as the Southern Ocean, have a slight positive increase, whilst other regions have slight negative trends. This implies that upper percentile trends are increasing faster than mean trends.

Many regions, such as Southern Australia, can be dominated by swell rather than local wind sea, and therefore increases in mean local wind speed may not have direct effects on the local wave climate. Instead, increasing wind speed in the higher latitudes may increase swell size, which propagates into the middle latitudes.


Important: Long-term decadal trends in wave height are occurring at different rates around the globe, with mean and extreme waves exhibiting different behaviours.



2.4.3 Climate oscillations

The **ENSO** phenomenon is the largest and most influential mode of climate variation that operates on a seasonal-to-interannual timescale. It is a complex ocean-air coupling in the equatorial region of the Central and Eastern Pacific Ocean that has significant influence over global climate.

- The positive **La Niña** phase leads to **reduced atmospheric convection** over the Pacific Ocean, causing **increased precipitation**, a **greater number of TCs**, and **cooler maximum temperatures** in Australia. There are stronger than normal trade winds, which could lead to **larger wave heights**.
- In contrast, the negative **El Niño** phase leads to a shift in atmospheric circulation, with a **weakening or reversal of the dominant south-easterly trade winds** and **decreased frequency of TC** for Australia.



Note: There is a substantial connection between the **two ENSO extreme phases** to inter-annual wave height and direction variability (Hemer et al. 2011). This typically results in **atypical coastal erosion** around the Pacific Ocean, with different locations experiencing diverse effects).

For example, the West Coast of the United States experiences an **increase in winter wave energy during El Niño phases**, leading to substantial beach erosion and coastal flooding. The region also experiences a more dominant southerly wave direction during the El Niño phase, which drives abnormal erosion in beaches normally protected from severe swell.

Understanding ENSO

In contrast, the **La Niña phase is linked to higher wave energy on the Eastern Australian coastline and more extreme storms.**

In New Zealand, increased H_s occurs during both of the ENSO extreme phases, resulting in increased coastal and marine operational risk alongside potential changes to biological and physical marine processes.

Note: Climate change is predicted to bring more frequent and extreme ENSO events of both phases, which could lead to atypical wave conditions. Therefore, it is imperative to understand the historical impacts of ENSO on wave climate to predict the future conditions for marine and coastal vulnerability globally.

The **SAM**, also known as the **Antarctic Oscillation**, is the principal mode of variability in the Southern Hemisphere extra-tropics and high latitudes. The SAM controls the north and south movement of the westerly wind belt that circulates Antarctica and is measured by the difference in zonal mean sea level pressure between 40S and 65S. SAM phases can significantly influence wave height and directional variability, influencing the entire Pacific Ocean.

- During the **positive phase**, the wind belt contracts towards Antarctica and is often **correlated to stable, dry conditions in Australia**.
- In the negative phase, the belt expands north and can lead to **increased storm frequency and precipitation in Australia and New Zealand** (Godoi et al. 2016).

Understanding the Southern Annular Mode (SAM)

Thus, climate oscillations can significantly modulate the wave climate around the globe, leading to diverse effects in different regions. Oscillations can also enhance or detract the effects of other oscillations, leading to complex interactions and influence on both atmospheric and wave climate.

2.5 Hands-on examples

To access the Notebooks exercises using the following link:



As we discussed, satellite radar altimeters can be used to determine significant wave height and wind speed. Analysis of past records bring new insights into inter-annual, seasonal and decadal variations of regional wave climates.

For this part of the course and to illustrate what we've seen today, we will use **RADWave** a Python package that provides a mechanism to access altimeter datasets through web-enabled data services (**THREDDS**).

We will work with the **Australian Ocean Data Network** database that spans from 1985-present and that has already been calibrated and validated.

With this tool we will query a range of spatial and temporal scales altimeter parameters in specific geographical regions and calculate:

- significant wave heights,
- periods,

- group velocities,
- average wave energy densities and
- wave energy fluxes.

Using **Jupyter Notebooks** as last week, you will:

- Extract data from different regions along Australia and evaluate wave modal conditions and seasonal changes in climate variability. Then calculate long-term trends and associated modulation by large-scale climate oscillations.
- Estimate the wave heights along Tropical Cyclone tracks.

RADWave Package by Smith et al. (2020)

CHAPTER
THREE

BIBLIOGRAPHY

3.1 References