

Spray

- Spray every **10 minutes**, Hold bottle **~1 inch away**, Use **5 sprays** each time
- If there is oozing, **spray more frequently** until it stops
- Place tissue on the forehead to prevent dressing from getting wet. Refrigerate the spray

Bandage

- **Do NOT change today.** If excessively moist, you may change the outer dressing
- Keep the pads in place

After Sedation (Important)

You may feel fine, but your body is still recovering and may be **slower than usual**.

- **Phone use:** Limit texting and work, you may not remember conversations clearly.
- **Driving / Car ride:** Avoid long drives if possible, Drive smoothly (limit stop-and-go), you **will get car sick easier**. You may feel **nauseous or lightheaded more easily**.

Lightheadedness (Very Common)

After long procedures, blood can temporarily pool in the legs → less blood flow to the brain → Lightheadedness, Nausea, Feeling faint.

Prevention: (1) Stay well hydrated (drink one cup of fluids hourly) (2) Wear compression socks (3) Avoid sudden movements

If you feel (A) lightheaded + nausea (B) lightheaded or (C) simply nauseous:

- **Immediately place your head down (low position)**
- **Bend your knees** → This helps blood flow back to your brain
- Stay there until symptoms resolve

If nauseous:

- **Do NOT rush to the bathroom**
- This can worsen symptoms and increase risk of fainting.
- **Immediately place your head down + Bend Knees until symptoms go away**

Hydration

- Drink **water or juice every hour** until bedtime. Use the restroom before going to sleep

Getting Out of Bed (3-Step Method)

1. **Sit up slowly**
 - Sit at edge of bed for a few minutes
 - Drink water or juice
2. **Stand up slowly**
 - Remain standing for 1–2 minutes
3. **Assess symptoms**
 - If lightheaded or nauseous → **lie back down immediately**
 - Sit when using the restroom

What's in Your Bag

- Staff will review all items with you
- **Pillow covers are important — use them**

From Dr. Sean:

My philosophy is to go above and beyond for every patient. We perform only one hair transplant procedure per day, allowing our entire team to focus our time, attention, and energy on achieving the best result possible for you.

We also encourage you to document your hair transplant journey and share your honest experience and progress. Your updates can be extremely helpful to others considering hair restoration.

Where to share your journey:

1. **Reddit** (sub reddit Hairtransplants) — Great for sharing photos, including immediately after surgery, and continuing to post updates as your hair grows.
2. **HairTransplantNetwork.com** — An excellent hair-transplant community. Members especially appreciate detailed post-operative photos and ongoing progress updates.
3. **Google Reviews** — A great place to share your overall experience with our office, team, and procedure.
4. **Yelp Reviews** — Another great place to share your overall experience with our office, staff, and your hair transplant journey.

IMPORTANT INSTRUCTIONS: PLEASE READ CAREFULLY A FEW TIMES. THIS IS IMPORTANT.
Dr. Sean Behnam's CELL PHONE: 310-237-8760

FUT Staple Removal Date: _____

DO NOT BUMP YOUR HEAD !!

- Getting in and out of the car.
- Cabinet door. This is a major culprit. Please be aware of this.

What's in your bag?

- **Spray Bottle:** Make sure it is in mist mode.
- **Pillow Cases:** It goes over your pillow. There will be some spotting overnight.
- **Tape Strips for forehead:** For those who had work done on the front of the scalp.
- **Medications:** Antibiotics, prednisone and Minoxidil.
- **Kerlix wraps:** To wrap your head
- **Tongue Depressor:** Used to apply ACell to donor area
- **ACell:**

Important things to remember for the first night:

- Spray routinely every 10 minutes when awake. Please set your alarm clock for 2AM and 4AM to spray as well. If you're tired, there's no need to get up at these times. IF you're coming back the next day for the 2nd procedure, no need to get up during the night and please sleep well.
- Keep the spray on the mist mode, not shooting mode. There is typically some oozing around the grafts and the spraying keeps it clean. The more you prevent blood clots, the less scabbing you will have later. If there is an area that blood is clotting, spray more on that area. Keep it very very clean. We do not want blood clots. Sometimes grafts exude a yellow sebum, spray it off. Get one to two inches close to the grafts.
- It's a long day. You have received medications. Sometimes, your body is catching up. Try not to make sudden moves.
 - Don't jump out of bed/chair. You need to do these in steps, otherwise you may get lightheaded.
- Pain Medications: California law has become very restrictive with codeine pain pills and are now written on an as-needed basis. Please discuss with Dr. Sean. For those with FUE procedure, pain is minimal and usually the majority may not need anything more than Tylenol. You can take two doses...

...of Tylenol 500mg once you get home or in the hotel. You may repeat it every 8 hours if necessary. After 48 hours, may add Advil (Ibuprofen). Please note to eat before taking Ibuprofen. No drinking of alcoholic beverages while taking Tylenol.

(Those who had FUE may skip this paragraph) For those who had the strip (FUT) procedure done, the office may have already communicated regarding prescription of Tylenol with Codeine. Please take one dose of codeine as soon as you get home. Repeat it again one hour before you go to bed. Please make sure you consume food every time prior to taking codeine so as not to get your stomach upset. If one codeine is too much for you (you're sensitive to it), take half a tablet.

- **Text Dr Sean (310-237-8760)** in the evening just to check in.
- **Sleeping the first night:** If you have had work done on the front of the scalp, I would like you to sleep at a 45 degree angle. You may want to place a few pillows behind your back. Use a neck pillow around your neck.
- It is OK to apply pressure to the back of your scalp. It's good. The pressure will cause clotting.
- If you need to bend over, squat so as to keep your head straight.
- **Antibiotic:** Please make sure to take your antibiotic when you get home. Most of the time, it is given prior to you leaving the office.
- **Nausea:** If you're sensitive to medications, please ask Dr Sean for anti-nausea medications.
- Eat a very good dinner and drink adequate fluids.
- Keep the compression socks on if it's not too tight.
- **Make sure you sit down when using the restroom** for the first 24 hours.
- Take it easy for the remainder of the day. **DO NOT** get sweaty and hot. **DO NOT** do any heavy lifting or strenuous exercise. For those traveling please make sure you're not lifting your luggage. **DO NOT** lean over to tie your shoes or pick up objects for the first 48 hours, as this may provoke bleeding or swelling (you may bend at the knees).

If you are planning to take any sleeping medication, please inform Dr. Behnam before you take it to make sure there are no interactions between the medications.

Bleeding: Bleeding is rare in the donor area. But if bleeding occurs, apply firm, steady pressure for 20-30 minutes. Repeat if necessary. Call Dr. Behnam immediately at **310-237-8760**. If bleeding persists, locate the exact bleeding point and apply pressure directly to the area with an ice pack. If you are out of town, go to the nearest emergency department, ask the attending physician to suture the area. If he/she has any questions, they should call Dr. Behnam at **(310) 237-8760** any time of the day or night.

Preparation for next day

- Have juice or water next to your bed.

When getting out of bed the next morning, follow the two step rule. Sit by the edge of the bed for one minute. Make sure you're not lightheaded. Drink water or juice. Then, get up slowly.

Summary for first night of the procedure:

- Spray every 10 minutes
- Pain Control if needed
- Sleep at 45 degrees, eat well, sleep well

Chart 1

Handwritten dates: [W 8/19](#) | [Th 8/20](#) | [F 8/21](#) | [Sa 8/22](#) | [Su 8/23](#) | [M 8/24](#)

Action	Day 0 (procedure night)	Day 1	Day 2	Day 3	Day 4	Day 5
Spray	e/v 10min	e/v 20min	e/v 30min	Optional e/v 1hr	Optional e/v 1hr	Optional e/v 1hr
Compression Socks	Yes	Yes				
Sitting down while Urinating	Yes	Yes				
Not Skipping meals	Yes	Yes				
Sleeping 45 degrees	Yes	Yes				
Tape over Forehead	Yes	Yes	Yes**	Yes	Yes	No*
Massage Forehead	No	Yes	Yes	Yes	Yes	No*
Ice Forehead	No	Yes	Yes	Yes	Yes	No*
Bend at knee level	Yes	Yes				

Action	Day 0	Day 1	Day 2	Day 3	Day 4	Day 5
2 Step out of bed	Yes	Yes				

* Swelling risk less by day five, can remove tape and stop massaging and icing

** May remove tape once a day for 10 minutes to let the forehead breathe. Massage the forehead while tape is off to prevent swelling.

We Will buy your after-photos from you (no face shown)!

We love to purchase your after-photos (no face shown), a comb-through your hair (anonymous) or a simple two minute testimonials. Photos \$500, Video \$750. This will really help others. Contact Dr. Sean for details.

If you like to document your journey on youtube/IG/Vlog with photos and videos, we will assist you with resources and pay for them.

Day Number 1 (the next day):

Be in communication with Dr Sean.

Breakfast/Lunch/Dinner: Don't skip any meals. Have a good breakfast before taking medications or changing the bandage.

Spraying the graft area (applies to both FUE and FUT Strip procedure): Keep spraying every 20 minutes. If there is an area that blood is clotting, spray more on that area. Keep it very very clean. Let it air dry. You can stand over the sink and spray it. The cleaner it is, the sooner you can go to work. You may apply a napkin or paper towel over your forehead to prevent the bandage from getting wet.

Tape on your forehead (applies to both FUE and FUT Strip procedure): If you had grafts placed above your forehead, then you must go home with tape on your forehead. Typically **swelling develops** around the eyes 4-5 days after the procedure. If you do develop swelling, this is normal and it will dissipate. If you wear this special tape provided on your forehead for 4 days and nights, you will not develop swelling or might reduce the swelling.

Icing the forehead (applies to both FUE and FUT Strip procedure): In order to minimize the swelling further, please ice the forehead three times a day, 5-10 minutes each time. Do not ice over the grafts. You can ice it until the tape is off after 4 days.

Massaging the swelling to the temples: You can minimize the swelling further by massaging the swelling from the middle of the tape to the temple areas. You can perform this massage a few times an hour.



Medications: Chart 2

Handwritten dates: Th-Fr 8/20-21 | Sa-M 8/22-24 | 8/25-29 Tu-Sa | 8/30-9/2 Su-W

Medication	Days 1-2	Days 3-5	Days 5-10	Days 10-14
Antibiotics	Yes	Yes	Yes	Yes if: - poor healing - ingrown hair history - prior infection
Prednisone	Yes	Yes	No More	
Tylenol, Codeine*	Yes as needed	As needed	As needed	As needed
NSAIDs (Advil, Ibuprofen)	NO	Yes as needed	As needed	As needed

* ...or while taking Tylenol.

Prednisone: This is a steroid and it is used as an anti-inflammatory. Please take it as directed in the morning with food until all is finished. Some people may not get them. Those above age 55 will not get them or get reduced doses.

- **Antibiotics:** Antibiotics prevent ingrown hair. For most people, it's Cephalexin. Take it twice daily (8AM and 8PM) until it's finished. If you are given Doxycycline, please make sure you eat first to prevent nausea. Also avoid dairy as it deactivates it. Infection is really rare. If in the unlikely event you have a fever, redness or pain, call Dr. Sean immediately. Some may get stomach upset with antibiotics, and in this case may take probiotics along with it as well.
- **Pain Medication:** You may take 2 doses of Tylenol with it as well (you can take up to 4,000 mg of Tylenol a day). For FUE procedure, most patients don't require much pain medications by the next day, Tylenol is sufficient. Whenever taking pain medications, please eat some food along with it to prevent an upset stomach.

ADVIL AND OTHER NSAIDS: Please ask Dr. Sean. Some patients may need to wait for 48 hours to take NSAIDS as they increase bleeding at the graft sites.

FUE, ACELL and the DONOR area: The donor area is the area where the grafts were removed. It is at the back and sides of the scalp. Please remove the bandage mid-day. You don't have to wash the donor area the 1st day as it may still be raw and tender. However, it would be advised for you to take ACELL ointment and apply it to the donor area and re-wrap it again. There is a regimen on how to use ACELL to get the best healing results. Please go to any pharmacy and buy a one-pound vaseline.

Apply ACell (one tablespoon) to the donor area three times a day, 8:00 AM, 12:00 PM, and 5:00 PM. After everytime you apply, re-wrap the donor area with the special kerlix provided. Two hours afterwards, remove the bandage and apply LOTS of vaseline over the ACell. There is no need to wash the ACell off first. Just keep applying over it. Keep doing this until you finish the ACell. Each jar of ACell should last about 4-5 days. As days pass and the tenderness decreases, you may rub the ACell into the holes with your fingers. This is important. When applying ACell, rub against the grain, from bottom to top. This ensures the ACell gets into the little holes. For those receiving two ACell jars, you may disregard the vaseline. Use only ACell.

Sleeping position: Sleep at 45 degree angle as well for the 2nd night. If you can't get good sleep, contact Dr. Sean. This will be your last night sleeping at 45 degrees.

Washing the donor and recipient: Not required.

Summary for Day 1 (Handwritten: *Th 8/20*)

- Spray e/v 20min
- Massage and ice forehead
- ACell x3/day
- Eat well, drink well
- Antibiotics + prednisone

- Pain: Tramadol or Codeine and Tylenol for pain, No NSAIDS
- Compression socks on
- Sleep 45 degrees

Day Number 2: **F 8/21**

- Refer to chart 1 and 2 above.
- Spray every 30 minutes
- Massage forehead a few times an hour
- Ice forehead 3 times a day, 5 minutes at a time. **NOT** the grafts
- ACell x3/day
- Eat well, drink well
- Take Antibiotics + prednisone
- Pain management: Tylenol for pain. **NSAIDS** like Advil is now ok. You may take two Advils (ibuprofen) and two Tylenol every 8 hours.
- Compression socks not necessary any more
- Sleep 45 degrees not necessary
- 2 step out of bed not necessary

Shampooing the Recipient Area : Chart 3

Action	Day 0-1	Day 2-11	Day 12	Day 13	Day 14	Day 15-30	Day 31
Rinse w/ shampoo	NO	Yes (no touching)				Yes	
Baby Oil overnight			Yes	Yes	Yes	As needed	
Shampoo + conditioner x15min			Yes	Yes	Yes	As needed	
Massage 5 min			Yes	Yes	Yes	As needed	
Regular showering							Yes

Handwritten note: 7/8/21 - 8/30

Shampooing the recipient area (The grafts) from Day 2 - Day 11

Note that until day 12, you may shampoo the grafts but **no need to touch or dab them**. Hard pressure may break the connection between blood vessels and the grafts. Nevertheless, you can wash them by rinsing to keep the area clean once a day or once every other day.

Please use clean unused plastic cup. Pour lukewarm water into the cup. We have sterile cups in the office. Add a small amount of Johnson's & Johnson's baby Shampoo. Shake the cup and gently rinse over the grafts. Then rinse clean water over the grafts to get rid of the residue. Let the hair air-dry.

Washing the donor area Day 2 onward

You may use Johnson's baby shampoo and gently massage the donor area. There is no problem massaging the donor area and cleaning it well. You do not have to scrape off the ACell.

For the donor area, you may rinse off the shampoo or even stand under water pressure as long as the water pressure doesn't hit the grafts and the water is not too hot and steamy. For those with crown work, rinsing the donor area using the cup is ideal.

Nutrition: There are various options

Good hair nutrition includes (A) protein concentrate plus (B) supplemental vitamins. Vitamins such as Biotin, Vitamin and Gelatin (protein).

Good supplemental proteins: Nutricost Whey protein Concentrate (Amazon), Premier Protein (Costco), Vega brand (vegetarian/Amazon), Orgain (wholefoods)

Vitamins: Options include Happy Head supplement (amazon) or you can get the individual vitamins on Amazon (vitamin D/ Gelatin/Biotin/saw palmetto).

Happy Head Supplement (Amazon) has Biotin, Vitamin D, Gelatin, Keratin, Saw palmetto, Vitamin D and A at exact dosage. Happy Head supplement can be purchased at Amazon.

There are TWO options for taking these:

Option #1) Nutricost whey protein plus Happy Head supplement

1. Morning routine: 1 scoop of Nutricost whey protein + one Happy Head capsule
2. Evening: one Happy Head capsule

OR

Option #2)

1. Morning routine: Protein in morning plus one gelatin 600 mg, one Vitamin D, Biotin and saw palmetto

1. Evening: one gelatin 600mg

Please follow these routines for 6 months.

Day Number 3:

- Refer to chart 1 and 2 and 3 above.
- Spraying is optional, every few hours.
- Massage forehead few times an hour
- Ice forehead 3 times a day, 5 minutes at a time. **NOT** the grafts
- ACell x3/day
- Take Antibiotics + prednisone
- Pain management: Ibuprofen and Tylenol. May use Tramadol or Codeine if necessary.
- Rinse the grafts with baby shampoo (as explained above on Day 2), although this can be done every other day.
- Clean and wash the donor area (explained on day 2)

Day Number 4:

- Refer to chart 1 and 2 and 3 above.
- No more Spraying.
- Massage your forehead a few times an hour.
- Ice forehead 3 times a day, 5 minutes at a time. Not the grafts
- ACell x3/day
- Take Antibiotics + prednisone
- Pain management: Ibuprofen and Tylenol. You may use Tramadol or Codeine if necessary.
- Rinse the grafts with baby shampoo every other day.
- Clean and wash the donor area every other day.

Itching:

Itching is common in both the recipient and donor areas. It's a sign that the nerves are waking up. For the recipient area you may use the spray bottle and spray the area if necessary with light padding. For the donor area, you may also pad.

Sometimes the itching can be significant, especially in the donor area. You may get Hydrocortisone 1% over the counter from any pharmacy and apply it twice a day in the donor area. If this doesn't work, ask Dr. Sean to send in stronger topical anti itch cream to your pharmacy. If the itch in the recipient area is still significant, please let Dr. Sean know.

Day Number 5:

- Refer to chart 1 and 2 and 3 above.
- You may remove the forehead tape gently.
- No need for forehead massage, tape or ice.
- ACell x3/day until it runs out.
- Take Antibiotics + last prednisone
- Pain management: Ibuprofen and Tylenol.
- Rinse the grafts with baby shampoo
- Clean and wash the donor area

Day Number 6:

- Refer to charts 2 and 3 above.
- ACell out by now. You can cleanse the donor area thoroughly with your fingertips.
- Take Antibiotics.
- As needed Ibuprofen and Tylenol.
- Rinse the grafts with baby shampoo.
- Clean and wash the donor area.

FAQ: Can't get rid of ACell from hair? Ask Dr Sean for assistance.

Day Number 7:

- Refer to charts 2 and 3 above.
- Take Antibiotics as needed, Ibuprofen and Tylenol if necessary.
- Rinse the grafts with baby shampoo.
- Clean and wash the donor area.

Preventing Ingrown hair in Donor area:

Do a vertical and horizontal squeeze on the donor area for 20 seconds, twice a day. This will eliminate ingrown hair in that region. Typically applies to ethnic hair or those with history of ingrown hair. Some people are prone to ingrown hair and may need topical ointment like clindamycin solution Or oral antibiotics such as doxycycline. Please keep Dr. Sean informed.

Days Number 8, 9, 10, 11:

- Refer to charts 2 and 3 above.
- Take Antibiotics. May stop if the donor is healed.
- As needed Ibuprofen and Tylenol if necessary.
- Rinse grafts with baby shampoo.
- Clean and wash the donor area.

Vertical and Horizontal Squeeze in Donor area.

Staple removal for those with FUT (strip) procedure (day 10). Not those with FUE.

FUT: How to make the staple removal more comfortable? One day prior to the staple removal date, apply a thick layer of Vaseline or Aquaphor or Neosporin and rub it into the staple area. Apply it in the morning and at night. Also apply it in the morning of the staple removal and one hour prior to coming in.

Days Number 12, 13, 14:

- Refer to charts 2 and 3 above.
- As needed Ibuprofen and Tylenol if necessary.
- Rinse the grafts with baby shampoo
- Clean and wash the donor area

Shampooing and removing the Scabs and crusts from Day 12 onward

This process can begin from the 12th day if there is significant crust built up or the 14th day if there is mild crusting and scabbing.

Please note that by day 12, please apply baby oil at night and let it sit overnight. In the morning please mix baby shampoo and any major brand conditioner together and lather it pretty well. Let it sit on the grafts for 20 minutes and then massage the area in circular fashion for four minutes. Most of the scabs should come off. Gently rinse the shampoo/conditioner off by rinsing water. Please note some hairs will fall out and it is OK. By now the grafts are rooted. Not all scabs and crusts will come off the first day. Repeat the process for a few more days until all scabs are removed.

No Shower pressure on the grafts for the first 4 weeks. You can shower regularly from neck down. Make sure the water is not too hot and the bathroom not too steamy.

Carefully blot dry your grafts with a dry towel or air-dry. **NO rubbing.**

You may use this technique using baby oil, shampoo and conditioner for the next several days until all scabs and crusts are out. Once they are out, become conservative again until the end of the month to resume regular showering.

Note ALL transplanted hairs will fall out and begin to regrow again by the 5th month. Good growth by the 8th month and full by the 12th month.

Shock Loss: Shock Loss is when some of your own natural hairs fall out after hair transplant. You may experience this shedding of your own natural hair in areas of thinning hair after hair transplant. 96% of the time, this does not happen.

...of the time, this does not happen. If it does, it is temporary due to the impact of hair transplant. It will grow back. These will regrow again. Please be patient. Take your protein and vitamins. Contact Dr. Sean please.

Ingrown hair is very common as the hair is growing out. It feels like acne and it might be painful. This is normal. Finishing your antibiotics will decrease the rate of ingrown hair. If you do get it, Dr. Behnam can prescribe clindamycin solution 1% to get rid of it.

Growth: Chart 4

Month 1	Months 2, 3, 4	Months 5, 6, 7	Months 8, 9, 10	Months 11, 12
Hairs begin to fall out	- All transplants have fell out - No growth - Shock loss present	Hairs begin to grow	Adequate growth	Final result

Follow ups:

After the first month, there will be multiple follow ups by either FaceTime or in person. I would like you to send me photos every two months. Please schedule a formal follow up by 12th month.

Please remember that it takes 4 months after transplant before the hair starts to grow. Please **BE PATIENT!** Hair grows only one inch every three months after the procedure starts. Full growth by 15 months.

There are two kinds of laser in our office. You get 10 LED red light low level laser therapy sessions. There is also Prodeep. Please call the office (310-315-4989) to make appointments for these sessions. Please note that these are different than the Prodeep 1340nm Laser sessions.

Please make regular appointments to follow your progress at 3 months, 6 months, 8 months, and 12 months and 15 months after the procedure. Dr. Behnam would love to continuously follow up to make sure everything is going well. If you can't make it to the office, please simply text Dr. Sean with your progress at the beginning of every month.

How can one get better healing of the Donor Area, smaller scarring and even some regrowth of the extracted hair?

Yes it is possible.

Material needed include: Lovtravel 16"x7.0" LED Red light (Amazon), Dermastamp 0.25mm (Amazon), Topical Solution such as Minoxidil (Amazon) or Compounded Minoxidil+Finasteride or compounded Minoxidil+Finasteride+Dutasteride.

[Product Diagrams: Lovtravel 16"x7.0" LED Red Light Therapy Device & Dermastamp 0.25mm]

When to begin?

21 days after the procedure. If you had a two day procedure, it would be 21 days after the 2nd day.

What's the technique?

Apply the dermastamp to the donor area for about two minutes. Not too deep and it should not bleed. Then apply the topical solution. Later that day or the following day, may use the LED light to the donor area. Perform this 3 times a week.

May begin the LED light and Solution around 21 days and then as sensitivity decreases add the dermastamp. Please perform this 3 (three) times a week.

Please clean the dermastamp after every use with isopropyl alcohol. Place isopropyl alcohol in spray bottle and spray it.

If the donor area gets irritated please stop.

There is also a new laser called Prodeep and this can help accelerate the healing of the donor area.

Prodeep laser for faster healing of the Donor Area after FUE

Prodeep ® laser is a non-ablative fractional laser at 1340 nm wavelength. Its unique 1340 nm wavelength allows it to penetrate deep into the skin to stimulate better wound healing of the donor area and faster hair regrowth. Only few sessions are necessary.

Minimal Downtime: No numbing required. No special care needed after procedure.

✓ Benefits After Hair Restoration

- ✓ Stimulates Dormant Hair Follicles
- ✓ Improves Hair Thickness and Density
- ✓ Enhances Blood Flow and Nutrient Supply to the Scalp
- ✓ Reduces Scalp Inflammation
- ✓ Promotes Healthier Scalp Tissue

Is it Possible to make the recipient area grow back faster after the shedding phase?

Medications that will regrow your hair faster:

1. **Oral Minoxidil:** It is the oral version of Rogaine. Works great with minimal side effects. Strongly recommended. Ask Dr. Sean for it.
2. **Oral Finasteride:** The oral is not Dr. Sean's favorite due to side effects. But if you're taking it and don't have side effects then that's great.
3. **Topical Finasteride/Minoxidil/Dutasteride:** Topical Finasteride/dutasteride has less side effects than oral does a very good job.

Technique:

May Begin one month after the procedure. **Not SOONER.**

You may use a Dermastamp 0.25mm and gently apply pressure on the recipient area for one minute. Make sure you're not applying too hard and no bleeding. Dermastamp three times a week only.

After the Dermastamping, may then apply the topical solution either Minoxidil 5% (Amazon) or Compounded Minoxidil+Finasteride or compounded Minoxidil+Finasteride+Dutasteride. These medications can be applied daily.

May also do LED LaserCap 3 times a week for 10 minutes.

There is a new laser called Prodeep 1340 nm laser that can accelerate hair regrowth. It may prevent shedding of the transplanted hair. If you are interested, please inform Dr. Sean. It will involve one session of 10 minutes a week for 8 weeks.

FREQUENTLY ASKED QUESTIONS (FAQ)

Activity / Item	Guideline / Restriction
Haircuts recipient area	Use Scissors for Six months on transplanted areas.
Clippers recipient area	Not recommended for 6 months.
Clippers donor area	10 days
Shaving whole head with Clippers	6 months
Hats	You may start wearing a CLEAN baseball hat from a few days after the procedure. Please make sure it is washed. I recommend trucker hats with links in preop. Loose. Straps in back. No tight fitting hat.
Jogging	14 days. Take it easy at first
Treadmill	14 days. Take it easy at first
Walking	Next day.
Heavy Lifting	21 days. Take It easy
Sports (Basketball, Baseball, Soccer, Football)	One month
Tennis	One month
Motorcycle helmet	1 month
Hard Hat	3-4 weeks
Alcohol	5 days. Do not take with Tylenol
Cigarette	10 days
Sex	5-6 Days
Swimming Pool	3 weeks, up to neck till 4th week.
Ocean	4 weeks
Sit ups FUT procedure	1 month
Sit ups FUE procedure	2-3 weeks
Hair system	3 weeks
Extensions	1-2 weeks. Make sure the...
Sauna	4 weeks
Hot showers	2-4 weeks. It will make your hair shafts swell and grafts may pop out.
Regular yoga	2 weeks

Activity / Item	Guideline / Restriction
Hot Yoga	4 weeks
Headstands	2 months
Color transplanted area	6 weeks, 10 minutes, least chemicals. No bleach.
Color non transplanted area	2 weeks
Blow Dryer with cold air	2 weeks, at a cool setting. Keep the comb or brush off of your scalp as you style.
Blow Dryer with warm air	1 month
Sun light	Don't get sunburn. Hats for 3 months if more than 20 minutes exposure.

Things that may delay or prevent your hair from growing:

- **Smoking:** Smoking cigarettes or marijuana will prevent your hair from growing. Smoking constricts blood vessels and doesn't allow oxygen to get to the grafts. So please stop smoking, definitely for the first 2 weeks.
- **Trauma to the scalp:** Please be careful. Don't do anything that can traumatize your grafts, i.e. surfing, motorcycling, bungee jumping.
- **Picking at scabs:** Don't pick at the grafts. If you have scabs, please follow the instructions above.
- **Significant weight loss in a short amount of time.**
- **General stress, shock etc.:** Stress can play a huge role in preventing hairs from growing. For example, death in the family, bad car accidents may cause a delay in hair growth.

Hair Stylist & Barber Contact:
Ron Dibuccio: 818-416-3881