

Contents

- [PART A](#)
- [PART B](#)
- [PART C](#)
- [PART D](#)
- [PART E](#)

```
%Tyler Matthews
%System Simluation Midterm P3
close; clc; close all;

A = [-4.7, -1.55, -0.55; 0.3, -2.75, -0.35; 1.1, 1.85, -2.55]
B = [1; 0; -1]
C = [2, 1, 1]
D = [0]
```

A =

```
-4.7000    -1.5500    -0.5500
 0.3000    -2.7500    -0.3500
 1.1000     1.8500    -2.5500
```

B =

```
1
0
-1
```

C =

```
2     1     1
```

D =

```
0
```

PART A

```
lamda = eig(A)
```

lamda =

```
-4.0000 + 0.0000i
-3.0000 + 0.8000i
```

-3.0000 - 0.8000i

PART B

```
Nt=21;
Nr=12;

theta=linspace(0,2*pi,1001);
rho=linspace(0,0.5256,1001);
tvec=linspace(0,2*pi,Nt);
rvec=linspace(0,0.5256,Nr);

T = linspace(0,1,1001);


hold on
plot(real(lamda(1)*T), imag(lamda(1)*T))
plot(real(lamda(2)*T), imag(lamda(2)*T))
plot(real(lamda(3)*T), imag(lamda(3)*T))
hold off

for k=1:length(rvec)
    z=rvec(k)*exp(i*theta);
    w=(z.^2-z.*1.45 + 0.45)./(z.*1.27-0.73);

    hold on
    plot(real(w), imag(w))
    hold off
end

for k=1:length(tvec)-1
    z=rho*exp(i*tvec(k));
    w=(z.^2-z.*1.45 + 0.45)./(z.*1.27-0.73);
    hold on
    plot(real(w), imag(w))
    hold off
end

grid on
axis([-1.25 0.75 -0.6 0.6])
```



PART C

```
stable_acc = 0.146;
stable_inacc = 0.3133;
unstable = 1;

disp("Stable and accurate at T = " + stable_acc);
disp("Stable and inaccurate at T = " + stable_inacc);
disp("Unstable and inaccurate at T = " + unstable);
```

```
Stable and accurate at T = 0.146
Stable and inaccurate at T = 0.3133
Unstable and inaccurate at T = 1
```

PART D

```
num = [0 1.27 -0.73];
den = [1 -1.45 0.45];

l1 = lamda * stable_acc
l2 = lamda * stable_inacc
l3 = lamda * unstable

stable_accurate_poles = roots(den - num*l1)
stable_inaccurate_poles = roots(den - num*l2)
unstable_inaccurate_poles = roots(den - num*l3)
```

```
l1 =  
  
    -0.5840 + 0.0000i  
    -0.4380 + 0.1168i  
    -0.4380 - 0.1168i
```

```
l2 =  
  
    -1.2532 + 0.0000i  
    -0.9399 + 0.2506i  
    -0.9399 - 0.2506i
```

```
l3 =  
  
    -4.0000 + 0.0000i  
    -3.0000 + 0.8000i  
    -3.0000 - 0.8000i
```

```
stable_accurate_poles =  
  
    0.6442 + 0.8364i  
    0.2689 - 0.4750i
```

```
stable_inaccurate_poles =  
  
    0.3648 + 1.0768i  
    0.0985 - 0.5902i
```

```
unstable_inaccurate_poles =  
  
    -0.1449 + 1.1381i  
    -0.1740 - 0.7222i
```

PART E

```
N = 10000;  
t = linspace(0,10,N);  
u = ones(1,N);  
fx1 = zeros(1,N);  
fx2 = zeros(1,N);  
fx3 = zeros(1,N);  
x1 = zeros(1,N);  
x2 = zeros(1,N);  
x3 = zeros(1,N);  
y = zeros(1,N);  
  
for k = 1:N-1  
    fx1(k+1) = -4.7*x1(k+1)-1.55*x2(k+1)-0.55*x3(k+1)+u(k+1);
```

```

fx2(k+1) = 0.3*x1(k+1)-2.75*x2(k+1)-0.35*x3(k+1);
fx3(k+1) = 1.1*x1(k+1)+1.85*x2(k+1)-2.55*x3(k+1)-u(k+1);

x1(k+2) = 1.45*x1(k+1) - 0.45*x1(k) + stable_acc * (1.27*fx1(k+1) - 0.73*fx1(k));
x2(k+2) = 1.45*x2(k+1) - 0.45*x2(k) + stable_acc * (1.27*fx2(k+1) - 0.73*fx2(k));
x3(k+2) = 1.45*x3(k+1) - 0.45*x3(k) + stable_acc * (1.27*fx3(k+1) - 0.73*fx3(k));

y(k) = 2*x1(k)+x2(k)+x3(k);
end
figure
plot(t,y)
xlim([0 0.1])
title('Stable and Accurate T')

N = 10000;
t = linspace(0,10,N);
u = ones(1,N);
fx1 = zeros(1,N);
fx2 = zeros(1,N);
fx3 = zeros(1,N);
x1 = zeros(1,N);
x2 = zeros(1,N);
x3 = zeros(1,N);
y = zeros(1,N);

for k = 1:N-1
    fx1(k+1) = -4.7*x1(k+1)-1.55*x2(k+1)-0.55*x3(k+1)+u(k+1);
    fx2(k+1) = 0.3*x1(k+1)-2.75*x2(k+1)-0.35*x3(k+1);
    fx3(k+1) = 1.1*x1(k+1)+1.85*x2(k+1)-2.55*x3(k+1)-u(k+1);

    %    x1(k+1) = x1(k) + stable_inacc*fx1(k);
    %    x2(k+1) = x2(k) + stable_inacc*fx2(k);
    %    x3(k+1) = x3(k) + stable_inacc*fx3(k);

    x1(k+2) = 1.45*x1(k+1) - 0.45*x1(k) + stable_inacc * (1.27*fx1(k+1) - 0.73*fx1(k));
    x2(k+2) = 1.45*x2(k+1) - 0.45*x2(k) + stable_inacc * (1.27*fx2(k+1) - 0.73*fx2(k));
    x3(k+2) = 1.45*x3(k+1) - 0.45*x3(k) + stable_inacc * (1.27*fx3(k+1) - 0.73*fx3(k));

    y(k) = 2*x1(k)+x2(k)+x3(k);
end
figure
plot(t,y)
xlim([0 0.1])
title('Stable and Inaccurate T')

N = 10000;
t = linspace(0,10,N);
u = ones(1,N);
fx1 = zeros(1,N);
fx2 = zeros(1,N);
fx3 = zeros(1,N);
x1 = zeros(1,N);
x2 = zeros(1,N);
x3 = zeros(1,N);
y = zeros(1,N);

```


```

for k = 1:N-1
    fx1(k+1) = -4.7*x1(k+1)-1.55*x2(k+1)-0.55*x3(k+1)+u(k+1);
    fx2(k+1) = 0.3*x1(k+1)-2.75*x2(k+1)-0.35*x3(k+1);
    fx3(k+1) = 1.1*x1(k+1)+1.85*x2(k+1)-2.55*x3(k+1)-u(k+1);


    x1(k+2) = 1.45*x1(k+1) - 0.45*x1(k) + unstable * (1.27*fx1(k+1) - 0.73*fx1(k));
    x2(k+2) = 1.45*x2(k+1) - 0.45*x2(k) + unstable * (1.27*fx2(k+1) - 0.73*fx2(k));
    x3(k+2) = 1.45*x3(k+1) - 0.45*x3(k) + unstable * (1.27*fx3(k+1) - 0.73*fx3(k));

    y(k) = 2*x1(k)+x2(k)+x3(k);
end
figure
plot(t,y)
xlim([0 0.1])
title('Unstable and Inaccurate T --> NOT REQUIRED FOR EXAM')

```



Stable and Inaccurate T



Unstable and Inaccurate T --> NOT REQUIRED FOR EXAM

