

# Scarcity of finite orbits for rational functions over a number fields.

Sebastian Troncoso

Birmingham-Southern College

Spring Southeastern Sectional Meeting  
Vanderbilt University, Nashville, TN

April 15, 2018.

# Notation

Let  $\phi : \mathbb{P}^n \rightarrow \mathbb{P}^n$  be an endomorphism defined over a number field  $K$ .

# Notation

Let  $\phi : \mathbb{P}^n \rightarrow \mathbb{P}^n$  be an endomorphism defined over a number field  $K$ .

**Periodic point:**  $\phi^m(P) = P$  for some  $m \geq 1$ .

# Notation

Let  $\phi : \mathbb{P}^n \rightarrow \mathbb{P}^n$  be an endomorphism defined over a number field  $K$ .

**Periodic point:**  $\phi^m(P) = P$  for some  $m \geq 1$ .

**Preperiodic point:**  $\exists m \geq 0$  such that  $\phi^m(P)$  is periodic.

The set of  $K$ -rational preperiodic points is denoted by  $PrePer(\phi, K)$ .

# Notation

Let  $\phi : \mathbb{P}^n \rightarrow \mathbb{P}^n$  be an endomorphism defined over a number field  $K$ .

**Periodic point:**  $\phi^m(P) = P$  for some  $m \geq 1$ .


**Preperiodic point:**  $\exists m \geq 0$  such that  $\phi^m(P)$  is periodic.

The set of  $K$ -rational preperiodic points is denoted by  $PrePer(\phi, K)$ .

**Tail point:** A point that is preperiodic but not periodic.

## Examples:

We can view  $\mathbb{P}^1(K)$  as  $K \cup \infty$  and endomorphism of  $\mathbb{P}^1$  as rational functions.



$\mathbb{Q}$ -rational tail points (red) and  $\mathbb{Q}$ -rational periodic points (green) of  $\phi_c(z) = z^2 + c$ .

## Question:

- Is the set  $\text{PrePer}(\phi, K)$  finite?

## Question:

- Is the set  $\text{PrePer}(\phi, K)$  finite?

**Yes.**



## Question:

- Is the set  $\text{PrePer}(\phi, K)$  finite?

**Yes.**

### Theorem (Northcott 1950)

Let  $\phi : \mathbb{P}^n \rightarrow \mathbb{P}^n$  be an endomorphism of degree  $\geq 2$  defined over a number field  $K$ . Then  $\phi$  has only finitely many preperiodic points in  $\mathbb{P}^n(K)$ .

## Question:

- Is the set  $\text{PrePer}(\phi, K)$  finite?

**Yes.**

### Theorem (Northcott 1950)

Let  $\phi : \mathbb{P}^n \rightarrow \mathbb{P}^n$  be an endomorphism of degree  $\geq 2$  defined over a number field  $K$ . Then  $\phi$  has only finitely many preperiodic points in  $\mathbb{P}^n(K)$ .

- How large is the set  $\text{PrePer}(\phi, K)$ ?
- Can we give an explicit bound?
- How is the bound depending on  $\phi$ ?

We can deduce from the original proof of Northcott's theorem a bound for  $|\text{PrePer}(\phi, K)|$  depending on

- $D = [K : \mathbb{Q}]$
- The dimension  $n$  of the projective space
- The degree  $d$  of  $\phi$ .
- height of the coefficients of  $\phi$

## Example

Consider  $f(x) = (x - 1)(x - 2)(x - 3) \dots (x - d) + x$ .

# Example

Consider  $f(x) = (x-1)(x-2)(x-3)\dots(x-d) + x$ .  
Notice that  $f$  has at least  $d$  periodic points.

1 ↻

2 ↻

⋮

$d$  ↻

# Example

Consider  $f(x) = (x-1)(x-2)(x-3)\dots(x-d) + x$ .  
Notice that  $f$  has at least  $d$  periodic points.

1 ↻

2 ↻

⋮

$d$  ↻

# The Dream:

Give explicit bounds for  $|\text{PrePer}(\phi, K)|$  in terms of:

# The Dream:

Give explicit bounds for  $|\text{PrePer}(\phi, K)|$  in terms of:

- $D = [K : \mathbb{Q}]$
- The dimension  $n$  of the projective space
- The degree  $d$  of  $\phi$ .



# The Dream:

Give explicit bounds for  $|\text{PrePer}(\phi, K)|$  in terms of:

- $D = [K : \mathbb{Q}]$
- The dimension  $n$  of the projective space
- The degree  $d$  of  $\phi$ .

Conjecture (Uniform Boundedness Conjecture - Morton–Silverman 1994)

There exists a bound  $B = B(D, n, d)$  such that if  $K/\mathbb{Q}$  is a number field of degree  $D$ , and  $\phi : \mathbb{P}^n \rightarrow \mathbb{P}^n$  is an endomorphism of degree  $d \geq 2$  defined over  $K$ , then

$$|\text{PrePer}(\phi, K)| \leq B.$$

# Goal:

- Give an explicit bound for  $|\text{PrePer}(\phi, K)|$ .
- To do so we need an extra parameter.
- Instead of the height of  $\phi$  we use a weaker and more natural parameter.
- This parameter is the number of places of bad reduction of  $\phi$

# Good reduction

- For simplicity in the notation we will defined good reduction only for rational maps  $\phi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ .

# Good reduction

- For simplicity in the notation we will defined good reduction only for rational maps  $\phi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ .
- Let  $K$  be a number field,  $\mathcal{O}_K$  its ring of algebraic integers,  $\mathfrak{p}$  a non zero prime ideal of  $\mathcal{O}_K$  and  $\mathcal{O}_{\mathfrak{p}}$  the local ring at  $\mathfrak{p}$ .

# Good reduction

- For simplicity in the notation we will defined good reduction only for rational maps  $\phi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ .
- Let  $K$  be a number field,  $\mathcal{O}_K$  its ring of algebraic integers,  $\mathfrak{p}$  a non zero prime ideal of  $\mathcal{O}_K$  and  $\mathcal{O}_{\mathfrak{p}}$  the local ring at  $\mathfrak{p}$ .
- Write  $\phi$  in normal form:

$$\phi([x : y]) = [F(x, y) : G(x, y)],$$

where  $F(x, y)$  and  $G(x, y)$  are coprime homogeneous polynomials of the same degree, with coefficients in  $\mathcal{O}_{\mathfrak{p}}$  and at least one a  $\mathfrak{p}$ -unit.

# Good reduction

- For simplicity in the notation we will defined good reduction only for rational maps  $\phi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ .
- Let  $K$  be a number field,  $\mathcal{O}_K$  its ring of algebraic integers,  $\mathfrak{p}$  a non zero prime ideal of  $\mathcal{O}_K$  and  $\mathcal{O}_{\mathfrak{p}}$  the local ring at  $\mathfrak{p}$ .
- Write  $\phi$  in normal form:

$$\phi([x : y]) = [F(x, y) : G(x, y)],$$

where  $F(x, y)$  and  $G(x, y)$  are coprime homogeneous polynomials of the same degree, with coefficients in  $\mathcal{O}_{\mathfrak{p}}$  and at least one a  $\mathfrak{p}$ -unit.

- We say  $\phi$  has **good reduction** at  $\mathfrak{p}$  if  $F$  and  $G$  do not have a common zero module  $\mathfrak{p}$  in  $\mathbb{P}^1$ .

# Bound on the set of preperiodic point for a rational map

## Theorem

Let  $\phi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$  be a rational map of degree  $d \geq 2$  defined over a number field  $K$  and  $[K : \mathbb{Q}] = D$ . Suppose  $\phi$  has good reduction outside a finite set of places  $S$ , including all archimedean ones. Let  $s = |S|$ . Then

# Bound on the set of preperiodic point for a rational map

## Theorem

Let  $\phi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$  be a rational map of degree  $d \geq 2$  defined over a number field  $K$  and  $[K : \mathbb{Q}] = D$ . Suppose  $\phi$  has good reduction outside a finite set of places  $S$ , including all archimedean ones. Let  $s = |S|$ . Then

- $|PrePer(K, \phi)| \leq d^{2^{16s}(s \log(s))^D}$   
J.K. Canci and L. Paladino (2015).



# Bound on the set of preperiodic point for a rational map

## Theorem

Let  $\phi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$  be a rational map of degree  $d \geq 2$  defined over a number field  $K$  and  $[K : \mathbb{Q}] = D$ . Suppose  $\phi$  has good reduction outside a finite set of places  $S$ , including all archimedean ones. Let  $s = |S|$ . Then

- $|PrePer(K, \phi)| \leq d^{2^{16s}(s \log(s))^D}$   
J.K. Canci and L. Paladino (2015).

- $|PrePer(K, \phi)| \leq 5 \left( 2^{16sd^3} \right) + 3$   
S. Troncoso (2017).

# Bound on the set of preperiodic point for a rational map

## Theorem

Let  $\phi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$  be a rational map of degree  $d \geq 2$  defined over a number field  $K$  and  $[K : \mathbb{Q}] = D$ . Suppose  $\phi$  has good reduction outside a finite set of places  $S$ , including all archimedean ones. Let  $s = |S|$ . Then

- $|PrePer(K, \phi)| \leq d^{2^{16s}(s \log(s))^D}$   
J.K. Canci and L. Paladino (2015).
- $|PrePer(K, \phi)| \leq 5 \left( 2^{16sd^3} \right) + 3$   
S. Troncoso (2017).
- $|PrePer(K, \phi)| \leq \alpha d^2 + \beta d + \gamma$   
where  $\alpha, \beta$  and  $\gamma$  are roughly  $2^{78s}$ .  
J.K. Canci, S. Troncoso and S. Vishkautsan (submitted).

# Tools

- Logarithmic  $v$ -adic distance between points in  $\mathbb{P}^1(K)$ .



# Tools

- Logarithmic  $v$ -adic distance between points in  $\mathbb{P}^1(K)$ .



- Study the distance between tail point and periodic.



# Tools

- Logarithmic  $v$ -adic distance between points in  $\mathbb{P}^1(K)$ .



- Study the distance between tail point and periodic.



- The set of tail points and the set of periodic points bound each other.



# Tools

- Logarithmic  $v$ -adic distance between points in  $\mathbb{P}^1(K)$ .



- Study the distance between tail point and periodic.



- The set of tail points and the set of periodic points bound each other.



- Number of solution of the  $S$ -unit equation.



# Tools

- Logarithmic  $v$ -adic distance between points in  $\mathbb{P}^1(K)$ .



- Study the distance between tail point and periodic.



- The set of tail points and the set of periodic points bound each other.



- Number of solution of the  $S$ -unit equation.



- Get a bound for the set of preperiodic point under a mild hypothesis



# Tools

- Logarithmic  $v$ -adic distance between points in  $\mathbb{P}^1(K)$ .



- Study the distance between tail point and periodic.



- The set of tail points and the set of periodic points bound each other.



- Number of solution of the  $S$ -unit equation.



- Get a bound for the set of preperiodic point under a mild hypothesis



- We use big theorems to lift the mild hypothesis and get the theorem.  
(Riemann-Hurwitz, Baker's Theorem, Kisaka's analysis on Baker's Theorem)



# THANK YOU