

K -rational preperiodic hypersurfaces on \mathbb{P}^n

Sebastian Troncoso

Michigan State University



Birmingham-Southern College

July 4, 2017.

Notation

Let K be a number field and \mathcal{O}_K its ring of algebraic integers,

Notation

Let K be a number field and \mathcal{O}_K its ring of algebraic integers,

$\phi : \mathbb{P}^n \rightarrow \mathbb{P}^n$ be an endomorphism defined over K
 ϕ^m the m th iterate of ϕ .

Notation

Let K be a number field and \mathcal{O}_K its ring of algebraic integers,

$\phi : \mathbb{P}^n \rightarrow \mathbb{P}^n$ be an endomorphism defined over K
 ϕ^m the m th iterate of ϕ .

The **orbit** of a point $P \in \mathbb{P}^n$ is the set

$$O_\phi(P) = \{P, \phi(P), \phi^2(P), \phi^3(P), \dots\}.$$

Notation

Periodic point: $\phi^m(P) = P$ for some $m \geq 1$.

Notation

Periodic point: $\phi^m(P) = P$ for some $m \geq 1$.
Minimal m is called the **period** of P .

Notation

Periodic point: $\phi^m(P) = P$ for some $m \geq 1$.

Minimal m is called the **period** of P .

The set of K -rational periodic points for ϕ is denoted by $\text{Per}(\phi, K)$.

Notation

Periodic point: $\phi^m(P) = P$ for some $m \geq 1$.

Minimal m is called the **period** of P .

The set of K -rational periodic points for ϕ is denoted by $\text{Per}(\phi, K)$.

Preperiodic point: $\exists m \geq 0$ such that $\phi^m(P)$ is periodic

Notation

Periodic point: $\phi^m(P) = P$ for some $m \geq 1$.

Minimal m is called the **period** of P .

The set of K -rational periodic points for ϕ is denoted by $\text{Per}(\phi, K)$.

Preperiodic point: $\exists m \geq 0$ such that $\phi^m(P)$ is periodic
i.e. P has finite orbit.

Notation

Periodic point: $\phi^m(P) = P$ for some $m \geq 1$.

Minimal m is called the **period** of P .

The set of K -rational periodic points for ϕ is denoted by $\text{Per}(\phi, K)$.

Preperiodic point: $\exists m \geq 0$ such that $\phi^m(P)$ is periodic
i.e. P has finite orbit.

The set of K -rational preperiodic points for ϕ is denoted by $\text{PrePer}(\phi, K)$.

Notation

Periodic point: $\phi^m(P) = P$ for some $m \geq 1$.

Minimal m is called the **period** of P .

The set of K -rational periodic points for ϕ is denoted by $\text{Per}(\phi, K)$.

Preperiodic point: $\exists m \geq 0$ such that $\phi^m(P)$ is periodic
i.e. P has finite orbit.

The set of K -rational preperiodic points for ϕ is denoted by $\text{PrePer}(\phi, K)$.

Tail point: A point that is preperiodic but not periodic.

Notation

Periodic point: $\phi^m(P) = P$ for some $m \geq 1$.

Minimal m is called the **period** of P .

The set of K -rational periodic points for ϕ is denoted by $\text{Per}(\phi, K)$.

Preperiodic point: $\exists m \geq 0$ such that $\phi^m(P)$ is periodic
i.e. P has finite orbit.


The set of K -rational preperiodic points for ϕ is denoted by $\text{PrePer}(\phi, K)$.

Tail point: A point that is preperiodic but not periodic.

The set of K -rational tail points for ϕ is denoted by $\text{Tail}(\phi, K)$.

Examples:

We can view $\mathbb{P}^1(K)$ as $K \cup \{\infty\}$ and endomorphism of \mathbb{P}^1 as rational functions.



\mathbb{Q} -rational tail points (red) and \mathbb{Q} -rational periodic points (green) of $\phi_c(z) = z^2 + c$.

Question:

- Are the sets $\text{Tail}(\phi, K)$, $\text{Per}(\phi, K)$ and $\text{PrePer}(\phi, K)$ finite?

Question:

- Are the sets $\text{Tail}(\phi, K)$, $\text{Per}(\phi, K)$ and $\text{PrePer}(\phi, K)$ finite?
Yes.

Question:

- Are the sets $\text{Tail}(\phi, K)$, $\text{Per}(\phi, K)$ and $\text{PrePer}(\phi, K)$ finite?
Yes.

Theorem (Northcott 1950)

Let $\phi : \mathbb{P}^n \rightarrow \mathbb{P}^n$ be an endomorphism of degree ≥ 2 defined over a number field K . Then ϕ has only finitely many preperiodic points in $\mathbb{P}^n(K)$.

Question:

- Are the sets $\text{Tail}(\phi, K)$, $\text{Per}(\phi, K)$ and $\text{PrePer}(\phi, K)$ finite?
Yes.

Theorem (Northcott 1950)

Let $\phi : \mathbb{P}^n \rightarrow \mathbb{P}^n$ be an endomorphism of degree ≥ 2 defined over a number field K . Then ϕ has only finitely many preperiodic points in $\mathbb{P}^n(K)$.

We can deduce from the original proof of Northcott's theorem a bound for $|\text{PrePer}(\phi, K)|$ depending on

- n
- $[K : \mathbb{Q}]$
- the degree of ϕ
- height of the coefficients of ϕ

Goals:

Give explicit bounds for $|\text{Tail}(\phi, K)|$, $|\text{Per}(\phi, K)|$ and $|\text{PrePer}(\phi, K)|$ in terms of:

Goals:

Give explicit bounds for $|\text{Tail}(\phi, K)|$, $|\text{Per}(\phi, K)|$ and $|\text{PrePer}(\phi, K)|$ in terms of:

- $D = [K : \mathbb{Q}]$

Goals:

Give explicit bounds for $|\text{Tail}(\phi, K)|$, $|\text{Per}(\phi, K)|$ and $|\text{PrePer}(\phi, K)|$ in terms of:

- $D = [K : \mathbb{Q}]$
- The dimension n of the projective space

Goals:

Give explicit bounds for $|\text{Tail}(\phi, K)|$, $|\text{Per}(\phi, K)|$ and $|\text{PrePer}(\phi, K)|$ in terms of:

- $D = [K : \mathbb{Q}]$
- The dimension n of the projective space
- The degree d of ϕ .

Goals:

Give explicit bounds for $|\text{Tail}(\phi, K)|$, $|\text{Per}(\phi, K)|$ and $|\text{PrePer}(\phi, K)|$ in terms of:

- $D = [K : \mathbb{Q}]$
- The dimension n of the projective space
- The degree d of ϕ .

Conjecture (Uniform Boundedness Conjecture - Morton–Silverman 1994)

There exists a bound $B = B(D, n, d)$ such that if K/\mathbb{Q} is a number field of degree D , and $\phi : \mathbb{P}^n \rightarrow \mathbb{P}^n$ is an endomorphism of degree $d \geq 2$ defined over K , then

$$|\text{PrePer}(\phi, K)| \leq B.$$

Goals:

In order to get explicit bounds for the cardinality of the set $\text{PrePer}(\phi, K)$ we need an extra parameter.

Instead of the height of ϕ we can use a weaker and more natural parameter to get bound on $|\text{PrePer}(\phi, K)|$.

Goals:

In order to get explicit bounds for the cardinality of the set $\text{PrePer}(\phi, K)$ we need an extra parameter.

Instead of the height of ϕ we can use a weaker and more natural parameter to get bound on $|\text{PrePer}(\phi, K)|$.

This parameter is the number of places of bad reduction of ϕ .

Goals:

In order to get explicit bounds for the cardinality of the set $\text{PrePer}(\phi, K)$ we need an extra parameter.

Instead of the height of ϕ we can use a weaker and more natural parameter to get bound on $|\text{PrePer}(\phi, K)|$.

This parameter is the number of places of bad reduction of ϕ .

Give explicit bounds for $|\text{Tail}(\phi, K)|$, $|\text{Per}(\phi, K)|$ and $|\text{PrePer}(\phi, K)|$ in terms of:

- $D = [K : \mathbb{Q}]$
- The dimension n of the projective space
- The degree d of ϕ .
- The number of places of bad reduction of ϕ .

Good Reduction

Let S be a finite set of places K , including all archimedean ones.

Good Reduction

Let S be a finite set of places K , including all archimedean ones.

- We say that ϕ has **good reduction outside S** if ϕ has good reduction for every $\mathfrak{p} \notin S$.

Good Reduction

Let S be a finite set of places K , including all archimedean ones.

- We say that ϕ has **good reduction outside S** if ϕ has good reduction for every $\mathfrak{p} \notin S$.

If we allow the number of primes of bad reduction as a parameter, much more is known for the cardinality of the set of K -rational preperiodic points in the case of \mathbb{P}^1 .

Bounds independent of the degree

Theorem (S. Troncoso 2016)

Let K be a number field and S a finite set of places of K containing all the archimedean ones. Let ϕ be an endomorphism of \mathbb{P}^1 , defined over K , and $d \geq 2$ the degree of ϕ . Assume ϕ has good reduction outside S .

(a) If there are at least three K -rational tail points of ϕ then

$$|\text{Per}(\phi, K)| \leq 2^{16|S|} + 3.$$

(b) If there are at least four K -rational periodic points of ϕ then

$$|\text{Tail}(\phi, K)| \leq 4(2^{16|S|}).$$

Bounds independent of the degree

Theorem (S. Troncoso 2016)

Let K be a number field and S a finite set of places of K containing all the archimedean ones. Let ϕ be an endomorphism of \mathbb{P}^1 , defined over K , and $d \geq 2$ the degree of ϕ . Assume ϕ has good reduction outside S .

(a) If there are at least three K -rational tail points of ϕ then

$$|\text{Per}(\phi, K)| \leq 2^{16|S|} + 3.$$

(b) If there are at least four K -rational periodic points of ϕ then

$$|\text{Tail}(\phi, K)| \leq 4(2^{16|S|}).$$

Notice that under these hypotheses the bounds are independent of the degree of ϕ . Those hypotheses are sharp, *i.e.* if there are two (three) K -rational tail (periodic) points then $|\text{Per}(\phi, K)|$ ($|\text{Tail}(\phi, K)|$) must depend on d .

Distance between periodic and tail points

Theorem (S. Troncoso 2016)

Let ϕ be an endomorphism of \mathbb{P}^1 , defined over K . Suppose ϕ has good reduction outside S . Let $R \in \mathbb{P}^1(K)$ be a tail point and let n be the period of the periodic part of the orbit of R . Let $P \in \mathbb{P}^1(K)$ be any periodic point that is not $\phi^{mn}(R)$ for some m . Then $\delta_{\mathfrak{p}}(P, R) = 0$ for every $\mathfrak{p} \notin S$.

For simplicity suppose \mathcal{O}_S is a PID and write $P = [x : y]$ and $Q = [w : t]$ in coprime S -integer coordinates.

Distance between periodic and tail points

Theorem (S. Troncoso 2016)

Let ϕ be an endomorphism of \mathbb{P}^1 , defined over K . Suppose ϕ has good reduction outside S . Let $R \in \mathbb{P}^1(K)$ be a tail point and let n be the period of the periodic part of the orbit of R . Let $P \in \mathbb{P}^1(K)$ be any periodic point that is not $\phi^{mn}(R)$ for some m . Then $\delta_{\mathfrak{p}}(P, R) = 0$ for every $\mathfrak{p} \notin S$.

For simplicity suppose \mathcal{O}_S is a PID and write $P = [x : y]$ and $Q = [w : t]$ in coprime S -integer coordinates.

Using the theorem we get that there is a S -unit element u such that

$$xt - yw = u$$

Theorem (S. Troncoso 2016)

Let K be a number field and S a finite set of places of K containing all the archimedean ones. Let ϕ be an endomorphism of \mathbb{P}^1 , defined over K , and $d \geq 2$ the degree of ϕ . Assume ϕ has good reduction outside S . Then

- (a) $|\text{Per}(\phi, K)| \leq 2^{16|S|d^3} + 3.$
- (b) $|\text{Tail}(\phi, K)| \leq 4(2^{16|S|d^3}).$
- (c) $|\text{PrePer}(\phi, K)| \leq 5(2^{16|S|d^3}) + 3.$

Notice that the bounds obtained in the theorem are a significant improvement from the previous bound given by Canci and Paladino which was of the order $d^{2^{16s}(s \log(s))^D}$ for the set $|\text{PrePer}(\phi, K)|$.

Another technique

Using another technique we can get a better result in terms of d .

Another technique

Using another technique we can get a better result in terms of d .

Theorem (S. Troncoso 2016)

Let K be a number field and S a finite set of places of K containing all the archimedean ones. Let ϕ be an endomorphism of \mathbb{P}^1 , defined over K , and $d \geq 2$ the degree of ϕ . Assume ϕ has good reduction outside S . Then

$$(a) \quad |\text{Tail}(\phi, K)| \leq d \max \left\{ (5 \cdot 10^6 (d^3 + 1))^{|S|+4}, 4(2^{64(|S|+3)}) \right\}.$$

Another technique

Using another technique we can get a better result in terms of d .

Theorem (S. Troncoso 2016)

Let K be a number field and S a finite set of places of K containing all the archimedean ones. Let ϕ be an endomorphism of \mathbb{P}^1 , defined over K , and $d \geq 2$ the degree of ϕ . Assume ϕ has good reduction outside S . Then

(a) $|\text{Tail}(\phi, K)| \leq d \max \left\{ (5 \cdot 10^6 (d^3 + 1))^{|S|+4}, 4(2^{64(|S|+3)}) \right\}.$

(b) In addition, if ϕ has at least one K -rational tail point then then

$$|\text{Per}(\phi, K)| \leq \max \left\{ (5 \cdot 10^6 (d - 1))^{|S|+3}, 4(2^{128(|S|+2)}) \right\} + 1.$$

Current project: Arithmetic dynamics in \mathbb{P}^n .

We are generalizing our results and techniques in \mathbb{P}^1 into results and techniques in \mathbb{P}^n .

Notation of preperiodic hypersurfaces

Notation of preperiodic hypersurfaces

Let $\phi : \mathbb{P}^n \rightarrow \mathbb{P}^n$ be an endomorphism defined over K and H an irreducible hypersurface defined over K of degree e .

Notation of preperiodic hypersurfaces

Let $\phi : \mathbb{P}^n \rightarrow \mathbb{P}^n$ be an endomorphism defined over K and H an irreducible hypersurface defined over K of degree e .

The **orbit** of H is the set

$$O_\phi(H) = \{H, \phi(H), \phi^2(H), \phi^3(H), \dots\}.$$

Notation of preperiodic hypersurfaces

Periodic hypersurface: $\phi^m(H) = H$ for some $m \geq 1$.

Minimal m is called the **period** of H .

The set of K -rational periodic hypersurface (of degree e) is denoted by $\text{HPer}(\phi, K)$ ($\text{HPer}(\phi, K, e)$).

Preperiodic hypersurface: $\exists m \geq 0$ such that $\phi^m(H)$ is periodic
i.e. H has finite orbit.

The set of K -rational preperiodic hypersurface (of degree e) is denoted by $\text{HPrePer}(\phi, K)$ ($\text{HPrePer}(\phi, K, e)$).

Tail hypersurface: A hypersurface that is preperiodic but not periodic.

The set of K -rational tail hypersurface (of degree e) is denoted by $\text{HTail}(\phi, K)$ ($\text{HTail}(\phi, K, e)$).

Questions:

- Are the sets $\text{HTail}(\phi, K)$, $\text{HPer}(\phi, K)$ and $\text{HPrePer}(\phi, K)$ finite?

Questions:

- Are the sets $\text{HTail}(\phi, K)$, $\text{HPer}(\phi, K)$ and $\text{HPrePer}(\phi, K)$ finite?

No, an example by J. Bell, D. Ghioca, and T. Tucker have shown that in general these sets are not finite.

Questions:

- Are the sets $\text{HTail}(\phi, K)$, $\text{HPer}(\phi, K)$ and $\text{HPrePer}(\phi, K)$ finite?

No, an example by J. Bell, D. Ghioca, and T. Tucker have shown that in general these sets are not finite.

- Are the sets $\text{HTail}(\phi, K, e)$, $\text{HPer}(\phi, K, e)$ and $\text{HPrePer}(\phi, K, e)$ finite?

Questions:

- Are the sets $\text{HTail}(\phi, K)$, $\text{HPer}(\phi, K)$ and $\text{HPrePer}(\phi, K)$ finite?

No, an example by J. Bell, D. Ghioca, and T. Tucker have shown that in general these sets are not finite.

- Are the sets $\text{HTail}(\phi, K, e)$, $\text{HPer}(\phi, K, e)$ and $\text{HPrePer}(\phi, K, e)$ finite?

Yes.

Theorem (B. Hutz 2016)

Let $\phi : \mathbb{P}^n \rightarrow \mathbb{P}^n$ be an endomorphism of degree ≥ 2 defined over a number field K . Then there are only finitely many preperiodic K -rational subvarieties of degree at most e .

Theorem (B. Hutz 2016)

Let $\phi : \mathbb{P}^n \rightarrow \mathbb{P}^n$ be an endomorphism of degree ≥ 2 defined over a number field K . Then there are only finitely many preperiodic K -rational subvarieties of degree at most e .

His result is based on theory of canonical heights for subvarieties of \mathbb{P}^N . From his proof we can give a bound for the cardinality of the set $\text{HPrePer}(\phi, K, e)$ depending on

- n
- $[K : \mathbb{Q}]$
- the degree of ϕ
- e
- height of the coefficients of ϕ

Goals

Just like the one dimensional case.

We would like to give explicit bounds for the cardinality of the sets $\text{HTail}(\phi, K, e)$, $\text{HPer}(\phi, K, e)$ and $\text{HPrePer}(\phi, K, e)$ in terms of:

Goals

Just like the one dimensional case.

We would like to give explicit bounds for the cardinality of the sets $\text{HTail}(\phi, K, e)$, $\text{HPer}(\phi, K, e)$ and $\text{HPrePer}(\phi, K, e)$ in terms of:

- $D = [K : \mathbb{Q}]$.
- The dimension n of the projective space.
- The degree d of ϕ .
- The degree e of the hypersurfaces.

Goals

Just like the one dimensional case.

We would like to give explicit bounds for the cardinality of the sets $\text{HTail}(\phi, K, e)$, $\text{HPer}(\phi, K, e)$ and $\text{HPrePer}(\phi, K, e)$ in terms of:

- $D = [K : \mathbb{Q}]$.
- The dimension n of the projective space.
- The degree d of ϕ .
- The degree e of the hypersurfaces.
- The number of places of bad reduction of ϕ .

- For now, we have just proven the following result for \mathbb{P}^2 .

- For now, we have just proven the following result for \mathbb{P}^2 .

If $T \in \text{HTail}(\phi, K, e)$ then n_T is the period of the periodic part of T . Consider N the number of monomials of degree e in three variables.

- For now, we have just proven the following result for \mathbb{P}^2 .

If $T \in \text{HTail}(\phi, K, e)$ then n_T is the period of the periodic part of T . Consider N the number of monomials of degree e in three variables.

Theorem (S. Troncoso 2017)

Let ϕ be an endomorphism of \mathbb{P}^2 , defined over K and suppose ϕ has good reduction outside S . Let $\{P_i\}_{i=1}^{2N+1}$ be a set of K -rational periodic points of \mathbb{P}^2 . Assume that no $N+1$ of them lie in a curve of degree e . Consider $\mathcal{B} = \{H' \in \text{HPer}(\phi, K) : \forall 1 \leq i \leq 2N+1 \quad P_i \notin \text{supp } H'\}$ and $\mathcal{A} = \{T \in \text{HTail}(\phi, K, e) : \text{there is } l \geq 0 \quad \phi^{ln_T}(T) \in \mathcal{B}\}$. Then

$$|\mathcal{A}| \leq (2^{33} \cdot (2N+1)^2)^{(N+1)^3(s+2N+1)}$$

- There is strong result from dynamical system that states that the set of periodic points is Zariski dense.

- There is strong result from dynamical system that states that the set of periodic points is Zariski dense.
- On \mathbb{P}^2 . We can give an alternative proof than the one given by B. Hutz for the finiteness of the set $\text{HPrePer}(\phi, K, e)$, **the idea** is to use the previous theorem and the Zariski density of periodic points.

Tools

The theorem is based on the following three tools:

Tools

The theorem is based on the following three tools:

- Logarithmic v -adic distance between a point and a hypersurface.

Tools

The theorem is based on the following three tools:

- Logarithmic v -adic distance between a point and a hypersurface.
- Study the distance between tail hypersurfaces and periodic points.

Tools

The theorem is based on the following three tools:

- Logarithmic v -adic distance between a point and a hypersurface.
- Study the distance between tail hypersurfaces and periodic points.
- Finiteness of integral points of \mathbb{P}^n - $\{2n + 1$ hyperplanes in general position $\}$ due to Ru and Wong.

Tools

The theorem is based on the following three tools:

- Logarithmic v -adic distance between a point and a hypersurface.
- Study the distance between tail hypersurfaces and periodic points.
- Finiteness of integral points of \mathbb{P}^n - $\{2n + 1$ hyperplanes in general position $\}$ due to Ru and Wong.
- Bounds for the number of solutions to decomposable form equations (Evertse).

logarithmic v -adic distance

Let H be a hypersurface of \mathbb{P}^n defined over K of degree e .

logarithmic v -adic distance

Let H be a hypersurface of \mathbb{P}^n defined over K of degree e .
Further, suppose H is defined by

$$f = \sum_{|\mathbf{i}|=d} a_{\mathbf{i}} \mathbf{x}^{\mathbf{i}} \in K[\mathbf{X}]$$

an homogeneous polynomial of degree e .

logarithmic v -adic distance

Let H be a hypersurface of \mathbb{P}^n defined over K of degree e .
Further, suppose H is defined by

$$f = \sum_{|\mathbf{i}|=d} a_{\mathbf{i}} \mathbf{x}^{\mathbf{i}} \in K[\mathbf{X}]$$

an homogeneous polynomial of degree e .
Let v be a nonarchimedean place of K

logarithmic v -adic distance

Let H be a hypersurface of \mathbb{P}^n defined over K of degree e .
Further, suppose H is defined by

$$f = \sum_{|\mathbf{i}|=d} a_{\mathbf{i}} \mathbf{x}^{\mathbf{i}} \in K[\mathbf{X}]$$

an homogeneous polynomial of degree e .

Let v be a nonarchimedean place of K and $P = [x_0 : \cdots : x_n]$ a point in $\mathbb{P}^n(K)$

logarithmic v -adic distance

Let H be a hypersurface of \mathbb{P}^n defined over K of degree e .
Further, suppose H is defined by

$$f = \sum_{|\mathbf{i}|=d} a_{\mathbf{i}} \mathbf{x}^{\mathbf{i}} \in K[\mathbf{X}]$$

an homogeneous polynomial of degree e .

Let v be a nonarchimedean place of K and $P = [x_0 : \cdots : x_n]$ a point in $\mathbb{P}^n(K)$ such that $P \notin \text{supp}(H)$.

logarithmic v -adic distance

Let H be a hypersurface of \mathbb{P}^n defined over K of degree e .
Further, suppose H is defined by

$$f = \sum_{|\mathbf{i}|=d} a_{\mathbf{i}} \mathbf{x}^{\mathbf{i}} \in K[\mathbf{X}]$$

an homogeneous polynomial of degree e .

Let v be a nonarchimedean place of K and $P = [x_0 : \cdots : x_n]$ a point in $\mathbb{P}^n(K)$ such that $P \notin \text{supp}(H)$.

logarithmic v -adic distance between P and H with respect to v is given by

$$\delta_v(P; H) = v(f(x_0, \dots, x_n)) - e \min_{0 \leq i \leq n} \{v(x_i)\} - \min_{|\mathbf{i}|=e} \{v(a_{\mathbf{i}})\}$$

Distance between tail hypersurfaces and periodic points

Theorem (S. Troncoso 2017)

Let ϕ be an endomorphism of \mathbb{P}^n , defined over K . Suppose ϕ has good reduction outside S . Let H be a K -rational tail hypersurface, m the period of the periodic part of the orbit of H and H' the periodic hypersurface such that $H' = \phi^{m_0 m}(H)$ for some $m_0 > 0$. Let $P \in \mathbb{P}^n(K)$ be any periodic point such that $P \notin \text{supp}\{H'\}$. Then $\delta_v(P; H) = 0$ for every $v \notin S$.

For simplicity suppose \mathcal{O}_S is a PID. Assume H is defined by

$$f = \sum_{|\mathbf{i}|=d} a_{\mathbf{i}} \mathbf{x}^{\mathbf{i}} \in \mathcal{O}_S[\mathbf{x}]$$

an homogeneous polynomial of degree e with at least one coefficient in \mathcal{O}_S^* and $P = [x_0 : \cdots : x_n]$ in coprime S -integer coordinates.

Distance between tail hypersurfaces and periodic points

Theorem (S. Troncoso 2017)

Let ϕ be an endomorphism of \mathbb{P}^n , defined over K . Suppose ϕ has good reduction outside S . Let H be a K -rational tail hypersurface, m the period of the periodic part of the orbit of H and H' the periodic hypersurface such that $H' = \phi^{m_0 m}(H)$ for some $m_0 > 0$. Let $P \in \mathbb{P}^n(K)$ be any periodic point such that $P \notin \text{supp}\{H'\}$. Then $\delta_v(P; H) = 0$ for every $v \notin S$.

For simplicity suppose \mathcal{O}_S is a PID. Assume H is defined by

$$f = \sum_{|\mathbf{i}|=d} a_{\mathbf{i}} \mathbf{x}^{\mathbf{i}} \in \mathcal{O}_S[\mathbf{x}]$$

an homogeneous polynomial of degree e with at least one coefficient in \mathcal{O}_S^* and $P = [x_0 : \cdots : x_n]$ in coprime S -integer coordinates.

Using the theorem we get that there is a S -unit element u such that

$$f(x_0, \dots, x_n) = u$$

If $T \in \text{HTail}(\phi, K, e)$ then n_T is the period of the periodic part of T . Consider N the number of monomials of degree e in three variables.

Theorem (S. Troncoso 2017)

Let ϕ be an endomorphism of \mathbb{P}^2 , defined over K and suppose ϕ has good reduction outside S . Let $\{P_i\}_{i=1}^{2N+1}$ be a set of K -rational periodic points of \mathbb{P}^2 . Assume that no $N+1$ of them lie in a curve of degree e . Consider $\mathcal{B} = \{H' \in \text{HPer}(\phi, K) : \forall 1 \leq i \leq 2N+1 \quad P_i \notin \text{supp } H'\}$ and $\mathcal{A} = \{T \in \text{HTail}(\phi, K, e) : \text{there is } l \geq 0 \quad \phi^{ln_T}(T) \in \mathcal{B}\}$. Then

$$|\mathcal{A}| \leq (2^{33} \cdot (2N+1)^2)^{(N+1)^3(s+2N+1)}$$

Idea of the proof:

- 1 Use the d -veronese map.

Idea of the proof:

- 1 Use the d -veronese map.

$$\{P_i\}_{i=1}^{2N+1} \rightarrow \{H_i\}_{i=1}^{2N+1}$$

$$T \rightarrow P_T$$

$$T(P_i) = H_i(P_T)$$

Idea of the proof:

- 1 Use the d -veronese map.

$$\{P_i\}_{i=1}^{2N+1} \rightarrow \{H_i\}_{i=1}^{2N+1}$$

$$T \rightarrow P_T$$

$$T(P_i) = H_i(P_T)$$

- 2 Use the arithmetic relation given by the v -adic distance between a periodic point and a tail curve.

Idea of the proof:

- 1 Use the d -veronese map.

$$\{P_i\}_{i=1}^{2N+1} \rightarrow \{H_i\}_{i=1}^{2N+1}$$

$$T \rightarrow P_T$$

$$T(P_i) = H_i(P_T)$$

- 2 Use the arithmetic relation given by the v -adic distance between a periodic point and a tail curve.

$T(P_i)$ is a S -unit $\Rightarrow H_i(P_T)$ is a S -unit.

Idea of the proof:

- 1 Use the d -veronese map.

$$\{P_i\}_{i=1}^{2N+1} \rightarrow \{H_i\}_{i=1}^{2N+1}$$

$$T \rightarrow P_T$$

$$T(P_i) = H_i(P_T)$$

- 2 Use the arithmetic relation given by the v -adic distance between a periodic point and a tail curve.

$T(P_i)$ is a S -unit $\Rightarrow H_i(P_T)$ is a S -unit.

- 3 Use the Finiteness of integral points of \mathbb{P}^n - $\{2n + 1$ hyperplanes in general position $\}$ to prove that our system of equation has finitely many solutions.

Idea of the proof:

- 1 Use the d -veronese map.

$$\{P_i\}_{i=1}^{2N+1} \rightarrow \{H_i\}_{i=1}^{2N+1}$$

$$T \rightarrow P_T$$

$$T(P_i) = H_i(P_T)$$

- 2 Use the arithmetic relation given by the v -adic distance between a periodic point and a tail curve.

$$T(P_i) \text{ is a } S\text{-unit} \Rightarrow H_i(P_T) \text{ is a } S\text{-unit.}$$

- 3 Use the Finiteness of integral points of \mathbb{P}^n - $\{2n+1 \text{ hyperplanes in general position}\}$ to prove that our system of equation has finitely many solutions.

Take $F = H_1 \cdot \dots \cdot H_{2N+1} \in O_S^*$ has finitely many solutions.

Idea of the proof:

- 1 Use the d -veronese map.

$$\{P_i\}_{i=1}^{2N+1} \rightarrow \{H_i\}_{i=1}^{2N+1}$$

$$T \rightarrow P_T$$

$$T(P_i) = H_i(P_T)$$

- 2 Use the arithmetic relation given by the v -adic distance between a periodic point and a tail curve.

$$T(P_i) \text{ is a } S\text{-unit} \Rightarrow H_i(P_T) \text{ is a } S\text{-unit.}$$

- 3 Use the Finiteness of integral points of \mathbb{P}^n - $\{2n+1$ hyperplanes in general position $\}$ to prove that our system of equation has finitely many solutions.

Take $F = H_1 \cdot \dots \cdot H_{2N+1} \in O_S^*$ has finitely many solutions.

- 4 Use the known bound for the number of solutions to the decomposable form equation (Evertse).

Idea of the proof:

- 1 Use the d -veronese map.

$$\{P_i\}_{i=1}^{2N+1} \rightarrow \{H_i\}_{i=1}^{2N+1}$$

$$T \rightarrow P_T$$

$$T(P_i) = H_i(P_T)$$

- 2 Use the arithmetic relation given by the v -adic distance between a periodic point and a tail curve.

$$T(P_i) \text{ is a } S\text{-unit} \Rightarrow H_i(P_T) \text{ is a } S\text{-unit.}$$

- 3 Use the Finiteness of integral points of \mathbb{P}^n - $\{2n + 1$ hyperplanes in general position $\}$ to prove that our system of equation has finitely many solutions.

Take $F = H_1 \cdot \dots \cdot H_{2N+1} \in O_S^*$ has finitely many solutions.

- 4 Use the known bound for the number of solutions to the decomposable form equation (Evertse).

We get an explicit bound previous decomposable equation.

Future work:

- 1 Generalize the explicit result that we have in \mathbb{P}^2 to a result in \mathbb{P}^n .

Future work:

- 1 Generalize the explicit result that we have in \mathbb{P}^2 to a result in \mathbb{P}^n .
- 2 Assume that we have enough K -rational periodic points $(2N + 1 + e^2)$ to get a bound for the cardinality of $\text{HTail}(\phi, K, e)$ in \mathbb{P}^2 in terms of the number of places of bad reduction, $[K : \mathbb{Q}]$ and e (independent on the degree of ϕ).

Future work:

- 1 Generalize the explicit result that we have in \mathbb{P}^2 to a result in \mathbb{P}^n .
- 2 Assume that we have enough K -rational periodic points $(2N + 1 + e^2)$ to get a bound for the cardinality of $\text{HTail}(\phi, K, e)$ in \mathbb{P}^2 in terms of the number of places of bad reduction, $[K : \mathbb{Q}]$ and e (independent on the degree of ϕ).
- 3 Get a bound for $\text{HPrePer}(\phi, K, e)$ in \mathbb{P}^2 in terms of the number of places of bad reduction, $[K : \mathbb{Q}]$, e and the degree of ϕ .

Future work:

- 1 Generalize the explicit result that we have in \mathbb{P}^2 to a result in \mathbb{P}^n .
- 2 Assume that we have enough K -rational periodic points $(2N + 1 + e^2)$ to get a bound for the cardinality of $\text{HTail}(\phi, K, e)$ in \mathbb{P}^2 in terms of the number of places of bad reduction, $[K : \mathbb{Q}]$ and e (independent on the degree of ϕ).
- 3 Get a bound for $\text{HPrePer}(\phi, K, e)$ in \mathbb{P}^2 in terms of the number of places of bad reduction, $[K : \mathbb{Q}]$, e and the degree of ϕ .
- 4 Do the previous results in \mathbb{P}^n .

THANK YOU