



Weakly-affine monads

Paolo Perrone  

Department of Computer Science, University of Oxford

Fabio Gadducci  

Department of Computer Science, University of Pisa, Pisa, IT

Davide Trotta  

Department of Computer Science, University of Pisa, Pisa, IT

Abstract

To be written.

2012 ACM Subject Classification


Keywords and phrases string diagrams, gs-monoidal categories

Digital Object Identifier 10.4230/LIPIcs...

1 Weakly Markov categories and weakly affine monads


Let \mathcal{C} be a GS-category. For every object X , the set $\mathcal{C}(X, I)$ has a canonical commutative monoid structure as follows: the monoidal unit is the discard map $X \rightarrow I$, and given $a, b : X \rightarrow I$, their product ab is given by copying, as follows.

How to call them? effects? co-states?



If a morphism $f : X \rightarrow Y$ is copyable and discardable, precomposition with f induces a morphism of monoids $\mathcal{C}(Y, I) \rightarrow \mathcal{C}(X, I)$.

The monoid $\mathcal{C}(X, I)$ acts on the set $\mathcal{C}(X, Y)$: given $a : X \rightarrow I$ and $f : X \rightarrow Y$, $a \cdot f$ is given as follows,



and the product $(f, g) \mapsto f \cdot g := (f \otimes g) \circ \text{copy}_X$ is equivariant for this action in both variables.

Definition 1.1. A GS-category \mathcal{C} is called *weakly Markov* if for every object X , the monoid $\mathcal{C}(X, I)$ is a group.

Every Markov category is weakly Markov: for each X , the monoid $\mathcal{C}(X, I)$ is the trivial group.

Definition 1.2. Given two parallel morphisms $f, g : X \rightarrow Y$ in a weakly Markov GS-category \mathcal{C} , we say that f and g are *equivalent*, and write $f \sim g$, if they lie in the same orbit for the action of $\mathcal{C}(X, I)$, i.e. if there is $a \in \mathcal{C}(X, I)$ such that $a \cdot f = g$. We say they are *uniquely equivalent* if there is a unique $a \in \mathcal{C}(X, I)$ such that $a \cdot f = g$.

XX:2 Weakly-affine monads

Let's now consider the case where the GS structure comes from a commutative monad on a cartesian monoidal category \mathcal{D} . In this case, the monoid structure of Kleisli morphisms $X \rightarrow 1$ comes from the following canonical internal monoid structure of $T1$ in \mathcal{D} , given by

$$1 \xrightarrow{\eta} T1, \quad T1 \times T1 \xrightarrow{c} T(1 \times 1) \xrightarrow{\cong} T1.$$

The monoid structure of Kleisli morphisms $X \rightarrow 1$ is now given as follows. The unit is given by

$$X \xrightarrow{!} 1 \xrightarrow{\eta} T1,$$

and the multiplication of the morphisms $f^\#, g^\# : X \rightarrow T1$ is

$$X \xrightarrow{\text{copy}} X \times X \xrightarrow{f^\# \times g^\#} T1 \times T1 \xrightarrow{c} T(1 \times 1) \xrightarrow{\cong} T1.$$

► **Definition 1.3.** A commutative monad T on a cartesian monoidal category is called *weakly affine* if $T1$ with its canonical internal monoid structure is a group.

► **Proposition 1.4.** Let \mathcal{D} be a cartesian monoidal category, and let T be a commutative monad on \mathcal{D} . The Kleisli category of T is weakly Markov if and only if T is weakly affine.

Proof. First, suppose that $T1$ is an internal group, and denote by $\iota : T1 \rightarrow T1$ its inversion map. The inverse of the morphism $f^\# : X \rightarrow T1$ in $\text{Kl}_T(X, 1)$ is given by $\iota \circ f$: indeed, the following diagram commutes,

$$\begin{array}{ccccc} X & \xrightarrow{\text{copy}} & X \times X & & \\ f^\# \downarrow & & f^\# \times f^\# \downarrow & \searrow f \times (\iota \circ f) & \\ T1 & \xrightarrow{\text{copy}} & T1 \times T1 & \xrightarrow{\text{id} \times \iota} & T1 \times T1 \xrightarrow{c} T(1 \times 1) \\ \downarrow ! & & & & \downarrow \cong \\ 1 & \xrightarrow{\eta} & & & T1 \end{array}$$

where the bottom rectangle commutes since ι is the inversion map for $T1$. The analogous diagram with $\iota \times \text{id}$ in place of $\text{id} \times \iota$ commutes analogously.

Conversely, suppose that for every X , the monoid structure on $\text{Kl}_T(X, 1)$ has inverses. Then in particular we can take $X = T1$, and the inverse of the Kleisli morphism $\text{id} : T1 \rightarrow T1$ is an inversion map for $T1$. ◀

1.1 In terms of the Yoneda embedding

For context:

► **Proposition 1.5.** A monoid (M, m, e) is a group if and only if the associativity square

$$\begin{array}{ccc} M \times M \times M & \xrightarrow{m \times \text{id}} & M \times M \\ \downarrow \text{id} \times m & & \downarrow m \\ M \times M & \xrightarrow{m} & M \end{array} \quad (1)$$

is a pullback.

The same can be said more generally for a monoid object in a cartesian monoidal category.

Proof. The square (1) is a pullback, both of sets and of groups, if and only if given $a, g, h, c \in M$ such that $ag = hc$, there exists a unique $b \in M$ such that $g = bc$ and $h = ab$. First, suppose that g is a group. The only possible choice of b is

$$b = a^{-1}h = gc^{-1},$$

which is unique by uniqueness of inverses.

Conversely, suppose that (1) is a pullback. We can set $g, h = e$ and $c = a$ so that $ae = ea = a$. Instantiating the pullback property, there is a unique b such that $ab = e$ and $ba = e$, that is, $b = a^{-1}$. ◀

Let \mathcal{D} be a cartesian monoidal category. Consider the presheaf category $[\mathcal{D}^{\text{op}}, \mathbf{Set}]$, equipped with the Day convolution product,

$$F \boxtimes G \cong \int^{A, B \in \mathcal{D}} \mathcal{D}(-, A \times B) \times F(A) \times G(B).$$

The Yoneda embedding $\mathcal{D} \rightarrow [\mathcal{D}^{\text{op}}, \mathbf{Set}]$ is strong monoidal: indeed, for each X ,

$$1 \cong \mathcal{D}(X, 1),$$

since 1 is terminal, and for each X and Y , by Yoneda reduction,

$$\begin{aligned} \mathcal{D}(-, X) \boxtimes \mathcal{D}(-, Y) &\cong \int^{A, B \in \mathcal{D}} \mathcal{D}(-, A \times B) \times \mathcal{D}(-, X) \times \mathcal{D}(-, Y) \\ &\cong \mathcal{D}(-, X \times Y). \end{aligned}$$

Therefore, and by the universal property of products, at the level of individual hom-sets the Day convolution product of representable presheaves just takes the cartesian products of sets:

$$(\mathcal{D}(-, X) \boxtimes \mathcal{D}(-, Y))(A) \cong \mathcal{D}(A, X \times Y) \cong \mathcal{D}(A, X) \times \mathcal{D}(A, Y).$$

Take now an object M of \mathcal{D} . Since the Yoneda embedding is fully faithful and strong monoidal, a monoid structure (M, m, e) on M is equivalently a monoid structure on the representable presheaf $\mathcal{D}(-, M)$. This makes the individual hom-sets monoids, with unit and multiplication as follows for each object X :

$$\begin{aligned} 1 &\xrightarrow{\cong} \mathcal{D}(X, 1) \xrightarrow{e_*} \mathcal{D}(X, M) \\ \mathcal{D}(X, M) \times \mathcal{D}(X, M) &\xrightarrow{\cong} \mathcal{D}(X, M \times M) \xrightarrow{m_*} \mathcal{D}(X, M) \end{aligned}$$

This is precisely the monoid structure that we have defined in the previous section for $M = T1$.

► **Proposition 1.6.** *M is an internal group if and only if all the monoids $\mathcal{D}(X, M)$ are groups.*

Proof. By Proposition 1.5, M is a group object if and only if its associativity square (1) is a pullback. Since the hom-functor preserves and reflects all limits in its second argument, we have that (1) is a pullback if and only if for each object X , the following diagram (or

XX:4 Weakly-affine monads

94 equivalently, its bottom right square) is a pullback,

$$\begin{array}{ccccc}
 \mathcal{D}(X, M) \times \mathcal{D}(X, M) \times \mathcal{D}(X, M) & \xrightarrow{\quad\quad\quad} & \mathcal{D}(X, M) \times \mathcal{D}(X, M) & & \\
 \downarrow & \searrow \cong & \downarrow \cong & & \\
 95 \quad \mathcal{D}(X, M \times M \times M) & \xrightarrow{(m \times \text{id})_*} & \mathcal{D}(X, M \times M) & & \\
 \downarrow (\text{id} \times m)_* & & \downarrow m_* & & \\
 \mathcal{D}(X, M) \times \mathcal{D}(X, M) & \xrightarrow{\cong} & \mathcal{D}(X, M \times M) & \xrightarrow{m_*} & \mathcal{D}(X, M)
 \end{array}$$

96 where the unlabelled arrows are the unique ones that make the diagram commute. Again by
 97 Proposition 1.5, the diagram above is a pullback if and only if $\mathcal{D}(X, M)$ is a group. ◀

98 1.2 Examples of weakly affine monads

99 ► **Example 1.7.** We present a family of examples of commutative monads that are weakly
 100 affine but not affine. Let A be a commutative monoid (written multiplicatively). Then the
 101 functor $T_A := A \times -$ on **Set** has a canonical structure of commutative monad, where the
 102 lax structure components $c_{X,Y}$ are given by multiplying elements in A while carrying the
 103 elements of X and Y along.

104 Since $T_A(1) \cong A$, the monad T_A is weakly affine if and only if A is a group, and affine if
 105 and only if $A \cong 1$.

106 ► **Example 1.8.** Let $M^* : \mathbf{Set} \rightarrow \mathbf{Set}$ be the monad assigning to every set the set of finitely
 107 supported discrete *nonzero* measures on M^* , or equivalently let $M^*(X)$ for any set X be the
 108 set of nonzero finitely supported functions $X \rightarrow [0, \infty)$. The monad structure is defined in
 109 terms of the same formulas as for the distribution monad on **Set** and the components $c_{X,Y}$
 110 are also given by the formation of product measures, or equivalently point-wise products of
 111 functions $X \rightarrow [0, \infty)$.


112 Since $M^*1 \cong (0, \infty) \not\cong 1$, this monad is not affine. However the monoid structure of
 113 $(0, \infty)$ induced by M^* is the usual multiplication of positive real numbers, which form a
 114 group. Therefore M^* is weakly affine, and its Kleisli category is weakly Markov.

115 On the other hand, if the zero measure is included, then we obtain a commutative monad
 116 M which can be seen as the monad of semimodules for the semiring of nonnegative reals.
 117 Since $M1 \cong [0, \infty)$ is not a group under multiplication, M is not weakly affine.

118 ► **Example 1.9.** Here is a negative example. Consider the free abelian group monad F on
 119 **Set**. Its functor takes a set X and forms the set FX of finite multisets (with repetition,
 120 where order does not matter) of elements of X and their formal inverses. We have that
 121 $F1 \cong \mathbb{Z}$, which is an abelian group under addition. However, the monoid structure on $F1$
 122 induced by the monoidal structure of the monad corresponds to the *multiplication* in \mathbb{Z} ,
 123 which does not have inverses. Therefore F is not weakly affine.

124 1.3 Conditional independence in weakly Markov categories

125 ► **Definition 1.10.** A morphism $f : A \rightarrow X_1 \otimes \cdots \otimes X_n$ in a GS-category \mathcal{C} is said to
 126 exhibit *conditional independence of the X_i given A* if and only if it can be expressed as a
 127 product of the following form.




Note that this is slightly different from [?, Definition 6.6], although it is equivalent for the case of Markov categories.

► **Proposition 1.11.** *Let $f : A \rightarrow X_1 \otimes \cdots \otimes X_n$ be a morphism in a GS-category \mathcal{C} . Then f exhibits conditional independence of the X_i given A if and only if it is equivalent to the product of all its marginals. Moreover, in that case f is uniquely equivalent to the product of its marginals.*


This generalizes the fact that, in Markov categories, a distribution exhibiting conditional independence is the product of its marginals [?, Section 12].

Proof. Denote the marginals of f by f_1, \dots, f_n . Suppose that f is a product as in Definition 1.10. For each $i = 1, \dots, n$, by marginalizing, we get that f_i is equal to the following.




Therefore for each i we have that $f_i \sim g_i$.

Conversely, suppose that f is equivalent to the product of its marginals, i.e. that there exists $a : X \rightarrow I$ such that f is equal to the following.




One can then choose $g_i = f_i$ for all $i < n$, and $g_n = a \cdot f_n$, so that f is in the form of Definition 1.10. Moreover, by marginalizing over all the X_i at once, we see that



so that a is uniquely determined. ◀

► **Remark 1.12.** For $n = 2$, a morphism $f : A \rightarrow X \otimes Y$ in a weakly Markov GS-category \mathcal{C} exhibits conditional independence of X and Y given A if and only if the following equation holds.



151

152 ► **Lemma 1.13.** *Let \mathcal{D} be a cartesian monoidal category, and let T be a commutative monad*
 153 *on \mathcal{D} . A Kleisli morphism $f^\sharp : A \rightarrow T(X_1 \times \cdots \times X_n)$ exhibits conditional independence of*
 154 *the X_i given A if and only if it factors as follows*

$$\begin{array}{ccc}
 & A & \\
 (g_1^\sharp, \dots, g_n^\sharp) \downarrow & \searrow f^\sharp & \\
 TX_1 \times \cdots \times TX_n & \xrightarrow{c} & T(X_1 \times \cdots \times X_n),
 \end{array}$$

156 *for some Kleisli maps $g_i^\sharp : A \rightarrow TX_i$, where the map c above is the one obtained by iterating*
 157 *the multiplication of the monoidal structure (such a map is unique by associativity).*

158 **Proof.** In terms of the base category \mathcal{D} , a Kleisli morphism in the form of Definition 1.10
 159 reads as follows.

$$A \xrightarrow{\text{copy}} A \times \cdots \times A \xrightarrow{g_1^\sharp \times \cdots \times g_n^\sharp} TX_1 \times \cdots \times TX_n \xrightarrow{c} T(X_1 \times \cdots \times X_n).$$

161 Therefore $f^\sharp : A \rightarrow T(X_1 \times \cdots \times X_n)$ is exhibiting conditional independence if and only if it
 162 is in the form above. ◀

163 ► **Definition 1.14.** Let \mathcal{C} be a GS-category. We say that \mathcal{C} satisfies the *localized independence*
 164 *property* if whenever a morphism $f : A \rightarrow X \otimes Y \otimes Z$ exhibits conditional independence of
 165 $X \otimes Y$ (jointly) and Z given A , as well as conditional independence of X and $Y \otimes Z$ given
 166 A , then it exhibits conditional independence of X , Y and Z given A .

167 ► **Theorem 1.15.** *Let \mathcal{D} be a cartesian monoidal category, and let T be a commutative*
 168 *monad on \mathcal{D} . The following conditions are equivalent.*

- 169 1. T is weakly affine;
- 170 2. $\text{Kl}(T)$ is weakly Markov;
- 171 3. $\text{Kl}(T)$ satisfies the localized independence property;
- 172 4. For all objects X , Y , and Z , the following associativity diagram is a pullback.

$$\begin{array}{ccc}
 T(X) \times T(Y) \times T(Z) & \xrightarrow{\text{id} \times c_{Y,Z}} & T(X) \times T(Y \times Z) \\
 c_{X,Y} \times \text{id} \downarrow & & \downarrow c_{X,Y \times Z} \\
 T(X \times Y) \times T(Z) & \xrightarrow{c_{X \times Y,Z}} & T(X \times Y \times Z)
 \end{array} \tag{2}$$

174 **Proof.** 1 \Leftrightarrow 2: see Proposition 1.4.

175 2 \Rightarrow 3: Suppose $f : A \rightarrow X \otimes Y \otimes Z$ exhibits conditional independence of $X \otimes Y$ (jointly)
 176 and Z given A , as well as conditional independence of X and $Y \otimes Z$ given A . By marginalizing
 177 out X , we have that f_{YZ} exhibits conditional independence of Y and Z given A . Since by
 178 hypothesis f exhibits conditional independence of X and $Y \otimes Z$ given A , by Proposition 1.11
 179 we have that f is equivalent to the product of f_X and f_{YZ} . But, again by Proposition 1.11,
 180 f_{YZ} is equivalent to the product of f_Y and f_Z , so we have that f is equivalent to the product

of all its marginals. Using Proposition 1.11 in the other direction, this means that f exhibits conditional independence of X , Y and Z given A .

3 \Rightarrow 4: By the universal property of products, a cone over the cospan in (2) consists of maps $g_1^\sharp : A \rightarrow TX$, $g_{23}^\sharp : A \rightarrow T(Y \times Z)$, $g_{12}^\sharp : A \rightarrow T(X \times Y)$ and $g_3^\sharp : A \rightarrow TZ$ such that the following diagram commutes.

$$\begin{array}{ccccc}
 A & & \xrightarrow{(g_1^\sharp, g_{23}^\sharp)} & & T(X) \times T(Y \times Z) \\
 & \searrow^{(g_{12}^\sharp, g_3^\sharp)} & & \searrow & \downarrow c_{X, Y \times Z} \\
 & & T(X) \times T(Y) \times T(Z) & \xrightarrow{\text{id} \times c_{Y, Z}} & T(X) \times T(Y \times Z) \\
 & & \downarrow c_{X, Y} \times \text{id} & & \downarrow c_{X, Y \times Z} \\
 & & T(X \times Y) \times T(Z) & \xrightarrow{c_{X \times Y, Z}} & T(X \times Y \times Z)
 \end{array}$$

By Lemma 1.13, this amounts to a Kleisli map $f^\sharp : A \rightarrow T(X \times Y \times Z)$ exhibiting conditional independence of X and $Y \otimes Z$ given A , as well as of $X \otimes Y$ and Z given A . By the localized independence property, we then have that f exhibits conditional independence of all X , Y and Z given A , and so, again by Lemma 1.13, f^\sharp factors through the product $TX \times TY \times TZ$. More specifically, by marginalizing over Z , we have that g_{12}^\sharp factors through $TX \times TY$, i.e. the following diagram on the left commutes for some $h_1^\sharp : A \rightarrow TX$ and $h_2^\sharp : A \rightarrow TY$, and similarly, by marginalizing over X , the diagram on the right commutes for some $\ell_2^\sharp : A \rightarrow TY$ and $\ell_3^\sharp : A \rightarrow TZ$.

$$\begin{array}{ccc}
 A & \xrightarrow{g_{12}^\sharp} & T(X \times Y) \\
 (h_1^\sharp, h_2^\sharp) \downarrow & & \downarrow c \\
 TX \times TY & \xrightarrow{c} & T(X \times Y)
 \end{array}
 \quad
 \begin{array}{ccc}
 A & \xrightarrow{g_{23}^\sharp} & T(Y \times Z) \\
 (\ell_2^\sharp, \ell_3^\sharp) \downarrow & & \downarrow c \\
 TY \times TZ & \xrightarrow{c} & T(Y \times Z)
 \end{array}$$

In other words, the upper and the left curved triangles in the following diagram commute.

$$\begin{array}{ccccc}
 A & & \xrightarrow{(g_1^\sharp, g_{23}^\sharp)} & & T(X) \times T(Y \times Z) \\
 & \searrow^{(g_{12}^\sharp, g_3^\sharp)} & & \searrow & \downarrow c_{X, Y \times Z} \\
 & & T(X) \times T(Y) \times T(Z) & \xrightarrow{\text{id} \times c_{Y, Z}} & T(X) \times T(Y \times Z) \\
 & & \downarrow c_{X, Y} \times \text{id} & & \downarrow c_{X, Y \times Z} \\
 & & T(X \times Y) \times T(Z) & \xrightarrow{c_{X \times Y, Z}} & T(X \times Y \times Z)
 \end{array}$$

By marginalizing over Y and Z , there exists a unique $a^\sharp : A \rightarrow T1$ such that $h_1 = a \cdot g_1$. Therefore

$$g_{12} = h_1 \cdot h_2 = (a \cdot g_1) \cdot h_2 = g_1 \cdot (a \cdot h_2),$$

and so in the diagram above we can equivalently replace h_1 and h_2 with g_1 and $a \cdot h_2$. Similarly by marginalizing over X and Y , there exists a unique $c^\sharp : A \rightarrow T1$ such that $\ell_3 = c \cdot g_3$, so that

$$g_{23} = \ell_2 \cdot \ell_3 = \ell_2 \cdot (c \cdot g_3) = (c \cdot \ell_2) \cdot g_3$$

XX:8 Weakly-affine monads

and in the diagram above we can replace ℓ_2 and ℓ_3 with $c \cdot \ell_2$ and g_3 , as follows.

$$\begin{array}{c}
 \begin{array}{ccc}
 A & \xrightarrow{(g_1^\sharp, g_{23}^\sharp)} & T(X) \times T(Y) \times T(Z) \\
 \searrow (g_1^\sharp, (c \cdot \ell_2)^\sharp, g_3^\sharp) & \searrow (g_1^\sharp, (a \cdot h_2)^\sharp, g_3^\sharp) & \downarrow c_{X,Y} \times \text{id} \\
 & T(X) \times T(Y \times Z) & \downarrow c_{X,Y \times Z} \\
 & \downarrow c_{X,Y} \times \text{id} & \\
 & T(X \times Y) \times T(Z) & \downarrow c_{X \times Y, Z} \\
 & T(X \times Y \times Z) &
 \end{array}
 \end{array}$$

Now, marginalizing over Z and Z , we see that necessarily $a \cdot h_2 = c \cdot \ell_2$. Therefore there is a unique map $A \rightarrow TX \times TY \times TZ$ making the whole diagram commute, which means that (2) is a pullback.

4 \Rightarrow 1: If T is weakly affine, then taking $X = Y = Z = 1$ in (2) shows that this monoid must be an abelian group: we obtain a unique arrow $\iota: T(1) \rightarrow T(1)$ making the following diagram commute,

$$\begin{array}{ccccc}
 T1 & \xrightarrow{(id, \eta_1!)} & T1 \times T1 \times T1 & \xrightarrow{id \times c_{1,1}} & T1 \times T(1 \times 1) \xrightarrow{\cong} T1 \times T1 \\
 \searrow (id, \iota, id) & & \downarrow c_{1,1} \times id & & \downarrow c_{1,1 \times 1} \quad \downarrow c_{1,1} \\
 & & T(1 \times 1) \times T1 & \xrightarrow{c_{1 \times 1, 1}} & T(1 \times 1 \times 1) \xrightarrow{\cong} T(1 \times 1) \\
 & & \cong \downarrow & & \downarrow \cong \quad \downarrow \cong \\
 & & T1 \times T1 & \xrightarrow{c_{1,1}} & T(1 \times 1) \xrightarrow{\cong} T1
 \end{array}$$

and the commutativity shows that ι satisfies the equations making it the inversion map for a group structure. \blacktriangleleft

2 Additional material (to be added to section)

► **Proposition 2.1.** Let $(G, \cdot, 1)$ be a group and let X be a set. A function $\alpha: M \times X \rightarrow X$ determines a left action if and only if the square

$$\begin{array}{ccc}
 G \times G \times X & \xrightarrow{\cdot \times id} & M \times X \\
 \downarrow id \times \alpha & & \downarrow \alpha \\
 G \times X & \xrightarrow{\alpha} & X
 \end{array} \tag{3}$$

commutes and it is a pullback.

Proof. By definition, the square (3) commutes if and only if α and \cdot are compatible. Now we show that the commutative square (3) is a pullback if and only if α satisfies the identity axiom, i.e. $\alpha(e, x) = x$ for every x in X . Now, if (3) is a pullback, then there exists a

function $\beta : X \rightarrow G$ such that the diagram

$$\begin{array}{ccccc}
 & & \langle e!, \text{id} \rangle & & \\
 & \searrow & & \searrow & \\
 X & & & & G \times X \\
 & \swarrow \langle e!, \beta, \text{id} \rangle & & \swarrow \text{id} \times \alpha & \\
 & G \times G \times X & \xrightarrow{\text{id} \times \alpha} & G \times X & \\
 & \downarrow \cdot \times \text{id} & & \downarrow \alpha & \\
 & G \times X & \xrightarrow{\alpha} & X & \\
 & \swarrow \langle e!, \text{id} \rangle & & \swarrow & \\
 & & & &
 \end{array}$$

commutes, where $e! : X \rightarrow G$ is the function assigning the identity element e to every element x of X . Now, since the left triangle commutes, then we have that $e = e \cdot \beta(x)$ for every x of X , i.e. $\beta(x) = e$ for every x of X . Now, since the right triangle commutes, we can conclude that $\alpha(\beta(x), x) = \alpha(e, x) = x$ for every x in X .

Now we show that $\alpha(e, x) = x$ implies that the commutative square (3) is a pullback. Let us consider a set Y and the functions $\langle f_1, f_2 \rangle : Y \rightarrow G \times X$ and $\langle g_1, g_2 \rangle : Y \rightarrow G \times X$ such that $\alpha(f_1(y), f_2(y)) = \alpha(g_1(y), g_2(y))$. By applying $\alpha(f_1(y)^{-1}, -)$ to both sides, and then combining the compatibility of α with the assumption that $\alpha(e, x) = x$, we can conclude that $f_2(y) = \alpha(f_1(y)^{-1} \cdot g_1(y), g_2(y))$. Therefore, we can conclude that the diagram

$$\begin{array}{ccccc}
 & & \langle f_1, f_2 \rangle & & \\
 & \searrow & & \searrow & \\
 Y & & & & M \times X \\
 & \swarrow \langle f_1, \gamma, g_2 \rangle & & \swarrow \text{id} \times \alpha & \\
 & M \times M \times X & \xrightarrow{\text{id} \times \alpha} & M \times X & \\
 & \downarrow \cdot \times \text{id} & & \downarrow \alpha & \\
 & M \times X & \xrightarrow{\alpha} & X & \\
 & \swarrow \langle g_1, g_2 \rangle & & \swarrow & \\
 & & & &
 \end{array}$$

commutes, where the function $\gamma : Y \rightarrow M$ is defined by $\gamma(y) := f_1^{-1}(y) \cdot g_1(y)$. By the unicity of the inverse in a group, this function is also unique, and hence we can conclude that the commutative square (3) is a pullback. ◀

► **Proposition 2.2.** *If T is weakly affine, then for every object X , the morphism $c_{1,X} : T(1) \times T(X) \rightarrow T(X)$ determines a (left) group action.*

Proof. The compatibility axiom follows from the fact that the diagram

$$\begin{array}{ccc}
 T(1) \times T(1) \times T(X) & \xrightarrow{\text{id} \times c_{1,X}} & T(1) \times T(X) \\
 c_{1,1} \times \text{id} \downarrow & & \downarrow c_{1,X} \\
 T(1) \times T(X) & \xrightarrow{c_{1,X}} & T(X)
 \end{array}$$

commutes for every strong and commutative monad. Moreover, following the same proof used for Proposition 2.1, we can conclude that the identity axiom is satisfied since T is weakly affine. In particular, because $T(1)$ is a group by ??, and the previous square is a pullback (by definition of weakly affine monad). ◀

247 ► **Proposition 2.3.** *Let T be a weakly affine monad. If the diagram*

$$\begin{array}{ccc} T(1) & \xrightarrow{\text{id}} & T(1) \\ \downarrow \iota & & \downarrow \eta_{T1} \\ T(1) & \xrightarrow{T(\eta_1)} & T^2(1) \end{array}$$

249 *commutes, then $T^2(1) \cong T(1)$ in \mathcal{A} .*

250 **Proof.** To prove the result it is enough to show that $T(1) \cong 1$ in the Kleisli category \mathcal{A}_T .
 251 We know from Lemma that $T(1)$ is a group in \mathcal{A} , where the arrow $\eta_1: 1 \rightarrow T(1)$ is the
 252 unit of the group, and $\iota: T(1) \rightarrow T(1)$ is the inversion map. Therefore, we have that the
 253 composition $\iota\eta_1: 1 \rightarrow T(1)$ has to be equal to η_1 . Therefore, we can consider the arrows
 254 $1 \rightarrow T(1)$ and $T(1) \rightarrow 1$ in the Kleisli category \mathcal{A}_T given by $T(\eta_1)\eta_1$ and ι respectively. The
 255 composition $T(\eta_1)\eta_1$ with ι in \mathcal{A}_T is given by $\mu_{1,1}T(\iota)T(\eta_1)\eta_1$. Employing the naturality
 256 of η_1 and the fact that $\iota\eta_1 = \eta_1$, it is direct to check that $\mu T(\iota)T(\eta_1)\eta_1 = \eta_1$, that is the
 257 identity $1 \rightarrow 1$ in \mathcal{A}_T . Now to show that the other composition gives the identity on $T(1)$ in
 258 \mathcal{A}_T , it is enough to show that $T(\eta_1)\iota = \eta_{T(1)}$, but this follows by hypothesis. ◀

(Paolo) Credo che $T(\eta_1)\iota \neq \eta_{T(1)}$ nell'esempio delle misure non zero. Per ogni x in $(0, \infty) = T1$ abbiamo che $\eta_{T(1)}(x) = \delta_x$ (delta di Dirac), mentre $T\eta_1(\iota(x)) = T\eta_1(1/x) = 1/x \delta_1$.

260 ► **Corollary 2.4.** *Let T be a weakly affine monad. If the diagram*

$$\begin{array}{ccc} T(1) & \xrightarrow{\text{id}} & T(1) \\ \downarrow \iota & & \downarrow \eta_{T1} \\ T(1) & \xrightarrow{T(\eta_1)} & T^2(1) \end{array}$$

262 *commutes, then $T(1)$ is an idempotent group, namely $\iota = \text{id}_{T1}$.*

263 **Proof.** By ?? we have that $T(1)$ is a group. If $\eta_{T1} = T(\eta_1)\iota$, then we can apply the
 264 multiplication of the monad to both sides, obtaining $\iota = \text{id}_{T1}$. ◀

265 The following result shows that weak affinity occurs frequently. Recall that a strong monad
 266 $T: \mathcal{A} \rightarrow \mathcal{A}$ on a category \mathcal{A} with finite products is **affine** if $T(1) \cong 1$ (see also Remark ??).
 267 Three relevant examples of affine monads are the distribution monad on **Set** (for discrete
 268 probability), the Giry monad on the category of measurable spaces (for measure-theoretic
 269 probability, see Examples ?? and ??), and the expectation monad, see [?].

270 ► **Remark 2.5.** We are not aware of any relation between weakly affine monads in our sense
 271 and Jacobs' *strongly affine* monads [?], other than the fact that strongly affine implies affine
 272 implies weakly affine.