Feedback - Assignment 1

Help

You submitted this quiz on **Thu 10 Apr 2014 12:01 PM PDT**. You got a score of **8.00** out of **8.00**.


Question 1

These exercises are meant to give you practice with the MATLAB programming environment and are designed for you to demonstrate your competence performing basic computations using this software.

Part 1: Array computations for simple data analysis

For this question you will read in data that were obtained experimentally using a laser scanning confocal microscope operating in line-scan mode. The manipulations required in the assignment include: 1) accessing only a particular portion of the data (a defined spatial position), 2) spatial averaging to reduce noise, 3) normalization, and 4) converting the data into different units. First, for those of you who are interested, a few words about what the data represent. In this experiment a rat ventricular myocyte was loaded with the Ca2+ sensitive indicator fluo-3, and a confocal microscope was used to record changes in intracellular [Ca2+]. The microscope was operated in "line scan" mode such that one dimension of the resulting "pseudo-image" is space and the other is time. This cell was also loaded with the Ca2+ buffer NP-EGTA. This buffer has unique properties such that when it is exposed to high-intensity ultraviolet light, it loses its ability to bind Ca2+. Thus, delivering a pulse of UV light to a cell loaded with NP-EGTA provides a means to rapidly increase [Ca2+] by "uncaging" it from the NP-EGTA. The initial, local increase in [Ca2+] in this image results from a spatially localized flash of UV light. The second increase results from electrical stimulation of the cell. The goal is to determine how the increase in [Ca²⁺] due to the flash affects the later increase due to the electrical stimulus. The regions of the cell not exposed to the UV light serve as internal controls.

The data appear as follows:



To perform this analysis, you have to take the following steps:

a. Dowbload flash4.jpg from here.

Read in the file: data = imread('flash4.jpg','jpg');

- b. Take a look at it using the imagesc function. Orient the data so that time runs from left to right. Transpose the matrix if necessary.
- c. Average over the region of the UV flash. Store this in the variable flash. This should have dimensions 1 x 634.
- d. Average over a control region that does not contain the flash. Store this in the variable noflash
- e. The fluorescence units are arbitrary, since the number depends on laser intensity, dye concentration, microscope detector gain, etc. Thus, we are interested in relative changes in fluorescence (\mathbf{F}) over the baseline value (\mathbf{F}_0) in a resting cell. Convert from raw fluorescence to units of \mathbf{F}/\mathbf{F}_0 by normalizing flash and noflash to the average fluorescence in a region with no activity. (Hint: between lines 70 and 100 is a good region).
- f. To a first approximation, one can assume that the dye used in this experiment, fluo-3, only emits fluorescence when it is bound to Ca²⁺. Thus, where KD is the dissociation constant of the dye, and [Fluo3]TOTAL is the dye concentration, fluorescence is proportional to:

$$F \sim \frac{[Ca^{2+}][Fluo3]_{TOTAL}}{[Ca^{2+}] + K_D}$$

If one makes a reasonable assumption for baseline [$^{Ca^{2+}}$] (e.g. 100 nm), the following equation can be used to convert from a ratio R (units of $^{F/F_0}$) to [$^{Ca^{2+}}$] in units of concentration:


$$[Ca^{2+}] = \frac{RK_D}{\frac{K_D}{[Ca^{2+}]_{hawbw}} - R + 1}$$


Implement this equation to convert flash and noflash from units of F/F_0 to units of $[Ca^{2+}]$, in nM. You can assume $[Ca^{2+}]_{baseline} = 100$ nM, and $K_D = 700$ nM. Keep in mind that your variables flash and noflash are not just scalars (numbers) but are defined at many time points.


g. Plot both versus time, on the same scale, in different colors. The spacing between lines in this image was 1.53 ms. Thus your time vector should begin at zero and consist of points spaced 1.53 ms apart.


Which of the following plots can represent the output plot if you assume $[Ca^{2+}]_{baseline} = 150 \text{ nM}$ and $K_D = 1000 \text{ nM}$?

Your Answer Score Explanation









Question 2

Part 2: Visualization of data for statistical classification

For this question you will perform a rudimentary analysis to discriminate between two categories of data.

a. Read in the results contained in the file sampledata2.mat. (Download it from here.) Using the following MATLAB command:

>> load sampledata2

The simulated data in this file is structured as follows:

Column 1 patients' ages

Column 2 self-reported number of drinks per week

Column 3 clinical status: 1 = cancer, 0 = no cancer

Which of the following statements is correct:

Your Answer	Score	Explanation
Number of patients who have cancer is less than those without cancer.		
There are more patients who drink more than 6 times per week compared to those who drink less than 5 times per week.		
More than half of the patients drink more than 5 times per week.		
There are more patients who are older than 50 compared to those who are younger than 20.	✓ 1.00	
Total	1.00 / 1.00	


Question 3


Next, generate two histograms illustrating the following: (i) the ages of the patients who have cancer, (ii) the ages of the patients who do not have cancer. Plot them on the same graph so that it's easy to visualize whether the age distributions in the two groups are different.


Which of the following graphs correctly illustrates the histogram of the ages of the patients with


cancer?

Your Answer		Score	Explanation
•	~	1.00	











Total 1.00 / 1.00


Question 4

Which of the following graphs correctly illustrates the histogram of the ages of the patients who do not have cancer?

Your Answer Score Explanation







Question 5

Which of the following statements is correct?

Your Answer	Score	Explanation
Among patients without cancer, there are more people younger than 35 compared to those older than 35.		
Among patients without cancer, there are more people older than 55 compared to those younger than 25.		
• Among patients with cancer, there are more people younger than 30 compared to those older than 60.	1.00	
Among patients with cancer, there are more people older than 45 compared to those younger than 45.		
Total	1.00 /	
	1.00	

Question 6

What percent of patients with cancer are older than 35?

Your Answer		Score	Explanation
80.2	~	1.00	
90.2			
<u></u>			
60.2			
Total		1.00 / 1.00	

Question 7

What percent of all patients are younger than 25?

Your Answer		Score	Explanation
6	~	1.00	
<u>12</u>			
O 9			
_ 4			
Total		1.00 / 1.00	

Question 8

What percent of all patients who drink more than 3 times per week have cancer?

Your Answer		Score	Explanation
<u></u>			
44.5			
74.5			
6 4.5	~	1.00	
Total		1.00 / 1.00	