

# The Effects of Asymmetric Dark Matter on Stellar Evolution I: Spin-Dependent Scattering

Troy J. Raen,<sup>1\*</sup>, Héctor Martínez-Rodríguez<sup>1</sup>, Travis J. Hurst<sup>2</sup>, Andrew R. Zentner<sup>1</sup>, Carles Badenes<sup>1</sup>, and Rachel Tao<sup>3</sup>

<sup>1</sup>Department of Physics and Astronomy & Pittsburgh Particle Physics, Astrophysics, and Cosmology Center (Pitt PACC), University of Pittsburgh, Pittsburgh, PA 15260, USA

<sup>2</sup>Department of Mathematics and Physics, Colorado State University - Pueblo, Pueblo, CO, 81001

<sup>3</sup>Department of Physics, Emory University, Atlanta, GA 30322

10 September 2020

## ABSTRACT

Most of the dark matter (DM) search over last few decades has focused on WIMPs but the viable parameter space is quickly shrinking. Asymmetric Dark Matter (ADM) is a WIMP-like DM candidate with slightly smaller masses and no present day annihilation, meaning that stars can capture and build up large quantities of it. The captured ADM can transport energy through a significant volume of the star. We investigate the effects of spin-dependent ADM energy transport on stellar structure and evolution in stars with  $0.9 \leq M_*/M_\odot \leq 5.0$  in varying DM environments. We wrote a MESA module† that calculates the capture of DM and the subsequent energy transport within the star. We fix the DM mass to 5 GeV and the cross section to  $10^{-37} \text{ cm}^2$ , and we study varying environments by scaling the DM capture rate. For stars with radiative cores ( $M_* \lesssim 1.3 M_\odot$ ), the presence of ADM flattens the temperature and burning profiles in the core and increases MS lifetimes ( $X_c > 10^{-3}$ ) by up to  $\sim 20\%$ . In higher mass stars ( $1.3 M_\odot \leq M_* \leq 4.0 M_\odot$ ), ADM energy transport shuts off the convection in the core, limiting the fuel available and therefore shortening MS lifetimes by as much as  $\sim 40\%$ . This may translate to changes in the luminosity and effective temperature of the MS turnoff in stellar population isochrones. The with isochrones of  $\sim 0.1 \text{ Gyr}$  having slightly hotter MS turnoffs, and those with  $\sim 1 \text{ Gyr}$  having slightly cooler turnoffs. The tip of the red giant branch may occur at lower luminosities. The effects are largest in DM environments with high densities and/or low velocity dispersions, making dwarf and early forming galaxies most likely to display the effects.

**Key words:** keyword1 – keyword2 – keyword3

## 1 INTRODUCTION

A preponderance of the evidence suggests that approximately 84% of the matter budget of the universe consists of a form of non-baryonic dark matter that has yet to be identified (e.g., ???). In the standard picture of cosmological structure formation, galaxies form within the potential wells of large, nearly virialized halos of dark matter (??). If the dark matter interacts with standard model particles, it can be captured by stars moving through dark matter halos (????). Once captured, continued scattering within the stellar interior contributes to energy transport within the star, potentially altering its evolution (?????????) (?????????). The significance of this energy transport depends on the following properties (in addition to the properties of the star): (1) the DM mass,  $m_{\text{DM}}$ ; (2) the DM-nucleon scattering cross section,  $\sigma_n$ ; and (3) the total number of DM particles captured by a star,  $N_{\text{DM}}$ , which itself depends on  $m_{\text{DM}}$  and  $\sigma_n$  as well as the local DM environment from which the particles are captured (see § 2). We study the effects of energy transport by asymmetric dark matter (ADM, see below) in stars of mass

$0.9 \leq M_*/M_\odot \leq 5.0$  living within a variety of dark matter environments using the publicly-available code Modules for Experiments in Stellar Astrophysics (MESA, ????)

Evidence supporting the claim that  $\sim 84\%$  of the matter in the universe is in an unknown form of dark matter is abundant and varied, ranging from the anisotropy of the microwave background radiation to formation and structures of galaxies (e.g., ???). For several decades, the leading candidate has been the so called Weakly-interacting massive particle (WIMP). The classic WIMP is a heavy ( $m_{\text{DM}} \sim 10^2 - 10^3 \text{ GeV}$ ) thermal relic whose contemporary abundance is set by its annihilation rate in the early universe (e.g., ?). Therefore, WIMPs are thought to have a fairly well established “standard” annihilation cross section (e.g., ?), which is comparable to typical weak-scale cross sections,  $\langle \sigma v \rangle \sim 10^{-26} \text{ cm}^3/\text{s}$ . This annihilation of WIMPs, which is so critical to guaranteeing that the correct abundance of dark matter in the contemporary Universe, in turn, limits the number of particles that can accumulate within a star. The rate of capture of new dark matter particles comes

to equilibrium with dark matter particle annihilation in the stellar interior (?). Despite numerous ongoing terrestrial direct detection experiments (see ?, for a review) and efforts to detect dark matter indirectly through its annihilation products (reviewed in ?), dark matter has not been observed non-gravitationally. The available parameter space for relatively light ( $m_{\text{DM}} \lesssim 10^2 \text{ GeV}$ ) DM is rapidly shrinking, which has triggered a surge in research into alternatives to the long-favored WIMP.

Asymmetric dark matter (ADM) is an alternative to the classic WIMP in which the relic abundance of the dark matter particle is set by a primordial asymmetry rather than via annihilation (for a review, see ?, and references therein). If the baryon and dark matter asymmetries are related, then such models have the appealing property that they explain the fact that the contemporary dark matter and baryon abundances are of the same order of magnitude, which is otherwise surprising because these relic abundances are determined by unrelated physics in the WIMP scenario. Indeed, this was one of the early motivations for ADM-like models (e.g., ???). The variety of specific incarnations of ADM is broad, but ADM models often predict particle masses smaller than the classic WIMP ( $m_{\text{DM}} \sim 1 - 10 \text{ GeV}$ ) and little or no contemporary dark matter annihilation for lack of relic dark matter anti-particles.

These predictions motivate studies to constrain ADM indirectly through stellar astrophysics. The lack of annihilation means that ADM may build up to very large quantities within stars because the capture of ADM is never countered by annihilation. Meanwhile, the relatively low masses compared to the classic WIMP mean captured ADM particles orbit within a significant volume of the star, out to  $r_{\text{DM}} \sim 0.1R_{\odot}$  for a Sun-like star, which means that they experience large differences in ambient temperature throughout their orbits and can thus transport energy outward from the stellar core extremely efficiently (?). These features of ADM have already motivated research into the possibility that ADM may alter stellar evolution (e.g. ????????) (e.g. ?????????). In this paper we undertake a study of the properties and evolution of stellar populations within halos of ADM. In this first paper on the topic, we further narrow our study to spin-dependent ADM-nucleus scattering. Spin-independent ADM-nucleus scattering leads to behaviors that are qualitatively distinct from spin-dependent scattering; therefore we will present results for the former case in a forthcoming manuscript.

We generally find that ADM captured by stars can cool stellar cores to a degree that can have potentially observable effects on stellar populations. In general, the extra cooling due to ADM reduces the main sequence (MS) lifetimes of stars with masses between  $\sim 1.2M_{\odot}$  and  $\sim 5M_{\odot}$ , and alters their subsequent evolution. Stars with masses between  $0.9M_{\odot}$  and  $1.2M_{\odot}$  have their lifetimes altered in complicated manner that is not a monotonic function of the model parameters. We summarize our results on stellar lifetimes in Figure 3 and the net effect on stellar populations in the form of stellar isochrones in § 4. The effects of stellar cooling are particularly large in environments in which the ambient dark matter density is high and velocity dispersion is low, such that the capture of dark matter is extremely efficient. Thus, these effects will be largest in dwarf satellite galaxies and high-redshift galaxies.

In the following section, we summarize the dependence of

the capture rate of dark matter within stars on both dark matter and stellar properties. In § 3, we describe our simulations of stellar evolution including cooling due to ADM. We present our results in § 4. We discuss our results and draw conclusions in § 5.

## 2 DARK MATTER PROPERTIES AND CAPTURE IN STARS

Probing the parameter space of ADM with simulations of stellar evolution is computationally expensive. Consequently, we show results for a representative set of ADM parameters that we initially chose in order to: (1) make the effects of ADM on stellar evolution significant; and (2) remain consistent with contemporary constraints on dark matter properties (but see the discussion in the next paragraph). For our models we choose  $m_{\text{DM}} = 5 \text{ GeV}$  and a spin-dependent dark matter-nucleon scattering cross section of  $\sigma_p = 10^{-37} \text{ cm}^2$ . Hereafter we will discuss ADM-proton scattering since protons are the only nuclei in MS stars with both a significant abundance and a net spin. We assume that ADM self-interactions are negligible throughout; however, it is likely that self-interactions, if they existed, would lead to enhanced cooling (e.g., ?) and exploring such models would constitute a potentially interesting follow-up to this work.

While we were performing the numerical simulations described in this work, the PICO collaboration was able to reduce its thresholds, unambiguously excluding dark matter with the specific parameter values listed above (?). Unfortunately, the simulations that we have carried out are computationally intensive and it is impractical to repeat each of the  $\sim 600$  stellar evolution simulations. Consequently, we choose to present these results as a qualitative indication of the effects that ADM can have upon stellar evolution. As we will see below, there is significant uncertainty involved in associating a particular stellar effect with a particular dark matter-nucleon scattering cross section due to a variety of model uncertainties. Consequently, while the effects of dark matter within stars will likely be milder than those that we describe here, it is possible that the same, qualitative effects could be realized in nature.

The amount of energy transported by dark matter (see § 3.1) is proportional to the amount of ADM within the star. In ADM models, in which annihilation of dark matter is negligible, the number of dark matter particles within the star at any given time,  $t$ , is determined by  $dN_{\text{DM}}/dt = C_{\text{DM}}$ , where  $C_{\text{DM}}$  is the instantaneous ADM capture rate. We use the capture rate from ?, which is a simplified form valid for dark matter particle masses  $m_{\text{DM}} \lesssim 20 \text{ GeV}$  (see ?, for more complete capture rates) (see ?, for more complete capture rates):

$$C_{\text{DM}} = C_{\odot} \left( \frac{\rho_{\text{DM}}}{0.4 \text{ GeV cm}^{-3}} \right) \left( \frac{270 \text{ km s}^{-1}}{\bar{v}} \right) \times \left( \frac{\sigma_p}{10^{-43} \text{ cm}^2} \right) \left( \frac{5 \text{ GeV}}{m_{\text{DM}}} \right) \times \left( \frac{v_{\text{esc}}}{618 \text{ km s}^{-1}} \right)^2 \left( \frac{M_{\star}}{M_{\odot}} \right) \quad (1)$$

where  $C_{\odot} = 5 \times 10^{21} \text{ s}^{-1}$ ,  $\rho_{\text{DM}}$  is the DM density in the stellar environment,  $\bar{v}$  is the velocity dispersion of dark matter

particles in the stellar neighborhood, and  $v_{\text{esc}}$  is the escape speed from the surface of the star.

The first line of Eq. (1) gives the dependence of the capture rate on the stellar environment. Both  $\rho_{\text{DM}}$  and  $\bar{v}$  are properties of the local stellar environment and are degenerate with one another in Eq. (1). A higher ambient density of dark matter leads to a higher rate of dark matter capture, while a lower relative velocity between the star and the infalling dark matter leads to a higher probability for capture. Because of this degeneracy, coupled with the fact that these parameters carry considerable uncertainty themselves, it is convenient to parameterize a star's local dark matter environment by an overall factor (??),

$$\Gamma_B = \left( \frac{\rho_{\text{DM}}}{0.4 \text{ GeV cm}^{-3}} \right) \left( \frac{270 \text{ km s}^{-1}}{\bar{v}} \right). \quad (2)$$

Normalized in this way,  $\Gamma_B$  specifies the capture rate,  $C_{\text{DM}}$ , relative to the rate that would be realized in the solar neighborhood for the same star. From this point on, we will characterize a star's dark matter environment using  $\Gamma_B$ . In general, we will be most interested in values of  $\Gamma_B > 1$ . A value of  $\Gamma_B = 0$  describes a stellar environment with no dark matter (hereafter referred to as 'standard models' and labeled 'NoDM'), and  $\Gamma_B = 1$  describes the solar neighborhood. A value of  $\Gamma_B = 10^2$  may specify an environment in which the dark matter density is 100 times that in the solar neighborhood at the same velocity dispersion, an environment in which the velocity dispersion is 1/100 that of the solar neighborhood at the same density, or any of an infinite number of other possible combinations.

It is interesting to consider the range of  $\Gamma_B$  values that would be considered reasonable. If the distribution of dark matter within galaxies, such as the Milky Way, follows a profile that diverges as the ?, NFW density profile, then one might expect to find dark matter environments near the centers of galaxies with densities significantly higher than the local value and velocity dispersions significantly lower than the local value, giving  $\Gamma_B \gg 1$ . Such scenarios were explored in ?, ?, and ?. While such large values of  $\Gamma_B$  may well lead to large effects on stellar structure, stellar populations near the Galactic Center are difficult to observe and any assumption about the dark matter density profile in the inner regions of any galaxy must be considered speculative. Interestingly, Local Group dwarf galaxies are extremely dark matter-dominated and have well-constrained dark matter profiles and velocity dispersions. In some cases, the Local Group dwarfs have densities  $\sim 3$  orders of magnitude higher than the dark matter density in the Solar neighborhood and velocity dispersions that are at least  $\sim 2$  orders of magnitude smaller than the local value. This suggests that values of  $\Gamma_B \sim 10^{5-6}$  could be realized within Local Group dwarf galaxies and has the further merit that  $\Gamma_B$  within Local Group dwarfs can be measured more precisely in the future. A third possibility for large values of the environmental factor are early-forming, very high-redshift galaxies. These galaxies begin forming in small, dense halos where the environmental boost factor can reach  $\Gamma_B \gtrsim 10^6$  at redshifts  $z \gtrsim 10$  (?). Of course, these stars will not be directly observable, but it is interesting to speculate that such stars could be detected as remnants of early mergers with the proto-Milky Way and/or that changes to the structure and evolution of these stars could be detected indi-

rectly in the chemical evolution of the larger, lower-redshift galaxies in which they will be found today.

Finally, while we have focused on the environmental parameter,  $\Gamma_B$ , as a proxy for the dark matter environment in which a star is embedded, we note that values of  $\Gamma_B \neq 1$  can also be mimicked through dark matter physics. In particular, dark matter self-interactions can greatly enhance the capture rates of dark matter within stars (?). This effect of dark matter self-capture itself grows with increasing ambient density and decreasing ambient velocity dispersion, so the two effects reinforce one another. For example, a value of  $\Gamma_B \sim 10^4$  may be realized by increasing the ambient dark matter density by a factor of  $\sim 10^3$ , while simultaneously introducing a dark matter self-interaction that boosts the number of captured dark matter particles by a factor of  $\sim 10$ . We relegate the separation of these effects to future work and encapsulate all of this uncertainty into the single parameter  $\Gamma_B$ .

### 3 METHODS

We study the impact of dark matter on the evolution of  $0.9 \leq M_*/M_\odot \leq 5.0$  stars ~~through the thermal pulse (or equivalent) phase (with a mass step of  $0.05 M_\odot$ ) through core helium depletion ( $Y_e = 10^{-3}$ ) or a maximum age of 10 Gyr, whichever comes first~~, using the publicly-available code Modules for Experiments in Stellar Astrophysics (**MESA**, ?????), release 12115. We used the **MESA** SDK version 20190830<sup>1</sup> to compile **MESA**. We base our stellar parameter inlist on the **MESA** test suite `1M_pre_ms_to_wd` inlists and use a metallicity of  $Z = 0.0142$ . ~~12 of the models we ran did not complete (e.g., due to requiring unreasonably small timesteps), and we have excluded them from our final data set. Of these, 2 were  $\Gamma_B = 10^4$  models, and none of them were either NoDM or  $\Gamma_B = 10^6$  models (these are the three  $\Gamma_B$  values we highlight below).~~ We wrote a module that calculates dark matter capture and energy transport (see § 3.1) and connects to **MESA** simulations via the provided `extra_energy_implicit` hook. Our code is available on GitHub<sup>2</sup> and through the **MESA** Marketplace<sup>3</sup>.

To generate isochrones from **MESA**'s stellar models we must interpolate the data because time steps are adaptive and therefore data is not generated at a consistent set of ages. We attempted to use the **MIST** **MIST** code (**MESA** Isochrones and Stellar Tracks, ??), which uses a multi-step process to interpolate both the stellar tracks and the mass grid and therefore produces isochrones that are well-sampled, even through dynamic but short lived phases such as the sub-giant branch. However, this method failed to produce isochrones in much of our parameter space of interest, ~~including producing no isochrones older than 1 Gyr~~. This is likely due to non-monotonicity in the relation between initial stellar mass and age of a given evolutionary phase, which is present in our data (see § 4.3) but violates assumptions of the code. Instead we simply ~~interpolate~~ perform a linear interpolation of our stellar tracks to a uniform set of ages and choose isochrones that are as well-sampled as possible in regions of interest. **MESA**'s

<sup>1</sup> <https://zenodo.org/record/3560834>

<sup>2</sup> <https://github.com/troyraen/DM-in-Stars>

<sup>3</sup> [http://cococubed.asu.edu/mesa\\_market/add-ons.html](http://cococubed.asu.edu/mesa_market/add-ons.html)

adaptive time steps resolve dynamic phases of evolution quite well, and this interpolation is not problematic. We do not interpolate the mass grid.

### 3.1 Energy Transport by Dark Matter

The energy transported by captured ADM can, in principle, be computed by solving the Boltzmann equation; however, this strategy is too computationally intensive to combine with a full-scale simulation of the evolution of stellar structure. To reduce the computational costs of our simulations, we estimate ADM energy transport using the approximations of ?. In particular, we assume a Maxwellian phase-space distribution for the ADM and calculate an orbit-averaged temperature,  $T_{\text{DM}}$ , by requiring that the distribution satisfy the first moment of the Boltzmann equation. This amounts to a requirement on energy conservation: ADM should neither inject nor remove a net energy from the star. The rate of energy transfer (per unit mass) from dark matter to protons is then

$$\epsilon_{\text{DM}}(r) = 8\sqrt{\frac{2}{\pi}} \frac{n_{\text{DM}}(r)n_{\text{p}}(r)}{\rho(r)} \frac{m_{\text{DM}}m_{\text{p}}}{(m_{\text{DM}} + m_{\text{p}})^2} \sigma_{\text{p}} \\ \times \left( \frac{m_{\text{p}}kT_{\text{DM}} + m_{\text{DM}}kT(r)}{m_{\text{DM}}m_{\text{p}}} \right)^{1/2} k[T_{\text{DM}} - T(r)], \quad (3)$$

where  $n(r)$  is a number density,  $\rho(r)$  is the mass density,  $k$  is Boltzmann's constant, and the subscript p refers to protons. (See ?, for a detailed derivation).

Generally,  $n_{\text{p}}$ ,  $n_{\text{DM}}$ , and  $T$  all peak at the center (exceptions noted below), so the energy transport is most efficient here. The number density of dark matter particles,  $n_{\text{DM}}$  increases in proportion with  $N_{\text{DM}}$ , so we can expect the effects to increase with both  $\Gamma_{\text{B}}$  and stellar age through the MS, while hydrogen is abundant. As a star leaves the MS,  $n_{\text{p}}$  drops in the core and spin-dependent ADM energy transport is greatly diminished because there are relatively few protons left with which dark matter may scatter<sup>4</sup>.

The sign of  $\epsilon_{\text{DM}}(r)$  is given by the final term in (3),  $T_{\text{DM}} - T(r)$ , which is used to define an ADM characteristic radius,  $r_{\text{DM}}$ , implicitly as

$$T(r = r_{\text{DM}}) = T_{\text{DM}}. \quad (4)$$

Then dark matter takes energy from  $r < r_{\text{DM}}$  and deposits it at  $r > r_{\text{DM}}$  for a standard MS temperature profile (monotonically decreasing from the center outward). With our chosen ADM parameters we see typical values:

$$r_{\text{DM}} \sim \mathcal{O}(0.1R_{\star}) \quad (5)$$

$$l_{\text{DM}} = (\sigma_{\text{p}}n_{\text{p}})^{-1} \sim \mathcal{O}(1R_{\star}) \quad (6)$$

where  $l_{\text{DM}}$  is the ADM mean free path (implying that it completes several orbits between scattering events). These values allow dark matter to travel much larger distances than photons or ions within the star (which have  $l \lesssim 10^{-10}R_{\star}$ ) and to

<sup>4</sup> This is one of the primary reasons that spin-dependent and spin-independent scattering give qualitatively distinct results. As the star burns H on the MS, the number of protons is reduced, reducing the importance of spin-dependent scattering processes. In the case of spin-independent scattering, the effect gets more important as helium is produced from H burning during the MS.

traverse qualitatively distinct regions of the star. This large mean free path is what enables dark matter to serve as such an effective coolant despite being far less numerous than either photons or ions (?).

## 4 RESULTS

In standard stellar evolution, with no influence from dark matter, stars with  $0.9 \leq M_{\star}/M_{\odot} \leq 5.0$  naturally split into two groups with qualitatively different structures, based on the dominant channel through which they burn hydrogen. Spin-dependent asymmetric dark matter (ADM) affects core hydrogen burning, mainly by flattening the temperature gradient. In this section, we will review standard stellar astrophysics (?) and then describe the effects of asymmetric dark matter seen in our MESA simulations.

The dominant burning channel is determined by the core temperature, with the transition happening at  $T_c \sim 2 \times 10^7 \text{ K}$ , which corresponds to a stellar mass of  $M_{\star} \sim 1.3 M_{\odot}$ . Stars with  $M_{\star} \lesssim 1.3 M_{\odot}$ , which we call low-mass stars, burn hydrogen primarily through the proton-proton (pp) chain for which the burning rate scales with temperature very roughly as  $\epsilon_{\text{pp}} \propto T^4$ . For stars in the mass range  $0.4 \lesssim M_{\star}/M_{\odot} \lesssim 1.3$ , the transport of energy away from the core burning region is dominated by photon diffusion. Energy transport in the cores of such stars is said to be radiative.


Stars with  $M_{\star} \gtrsim 1.3 M_{\odot}$ , which we call high-mass stars, are dominated by the carbon-nitrogen-oxygen (CNO) cycle, for which the burning rate scales much more strongly with temperature,  $\epsilon_{\text{CNO}} \propto T^{16-20}$ . In CNO-dominated stars, radiative energy transport is insufficient to carry away the energy produced by hydrogen burning; consequently, they have convective cores.

In § 4.1 and § 4.2, we will consider results for low-mass stars and high-mass stars separately and we will demonstrate that ADM has distinct effects on the evolution of the two groups. § 4.3 details changes in MS lifetimes due to ADM energy transport. § 4.5 discusses the effects on surface properties of individual stars and in § 4.4, and on the isochrones of stellar populations in § 4.5. Note that all logarithms in this paper are base 10.

### 4.1 Low-Mass Stars

Standard model stars in the mass range  $0.9 \leq M_{\star}/M_{\odot} \lesssim 1.3$  have relatively low central temperatures and so are powered primarily by the pp chain, which is much less sensitive to the temperature than burning via the CNO cycle. This means the burning does not peak as strongly at the center and radiative transport is sufficient to carry the energy flux, so the core is radiative. Without the mixing provided by convection, hydrogen depletes first at the very center and the burning shifts gradually outward into a shell.

As seen in Figure 1, energy transport by large amounts of ADM causes flatter temperature profiles gradients in the center than those seen in the NoDM model. This reduces the burning rate in the center (where ADM is removing energy) and increases it in a shell (where ADM deposits energy). Indeed, the increased burning at larger radii ( $m(< r) \sim 0.1 M_{\odot}$ ) causes a small net increase in the total luminosity of the star. Because ADM probes temperature differences



**Figure 1.**  $1.0 M_{\odot}$  profiles for NoDM (grey) and  $\Gamma_B = 10^6$  (dark blue) models. Each set of 3 panels shows stellar profiles of the stars at different evolutionary phases indicated by the fraction of hydrogen in the center,  $X_c$ , which decreases as the star evolves. (ZAMS is “zero-age main sequence”. The profiles in each panel are: 1)  $\epsilon_{\text{nuc}}$ , the nuclear burning rate in [erg/g/s]; 2)  $\epsilon_{\text{DM}}$ , the rate at which DM transports energy (negative values indicate that energy is being removed), also in [erg/g/s]; 3)  $\log(T/K)$ ,  $\log_{10}$  of the temperature in [K]. ADM energy transport decreases the temperature and burning rate in the center and increases them in a shell at  $m(< r) \sim 0.1 M_{\odot}$ . The  $\Gamma_B = 10^6$  model reaches the  $X_c$  evolutionary markers at older ages, relative to the NoDM model, because the changes to the burning rates cause the central hydrogen fraction to decline more slowly.


over large portions of the star, energy transport by ADM remains efficient despite the shallower temperature gradient. The increased temperature at  $m(< r) \gtrsim 0.1 M_{\odot}$  means that more hydrogen burns during the MS in low-mass ADM models than in their NoDM counterparts. These two competing effects dictate the gross evolution of the star. For low values of  $\Gamma_B$  (weaker ADM influence), the increase in the total amount of fuel wins and the stellar MS lifetime is an increasing function of  $\Gamma_B$ . At values of  $\Gamma_B \gtrsim 10^3$ , the burning rate continues to increase, but little additional fuel is burned because of the precipitous drop in temperature at larger radii, which cannot be overcome by the energy transport. The result is that the MS lifetime of the star decreases with increasing environmental factor for  $\Gamma_B \gtrsim 10^3$ . (We will discuss stellar lifetimes further in § 4.3). Stars in this regime have a higher surface luminosity than their NoDM counterparts, at fixed central hydrogen fraction, and move along their. However, at fixed luminosity we should expect their effective temperatures to be roughly the same, because these are equilibrium points dictated by the equations of stellar structure. Indeed, this is what we see in our models. The result is that stars of a given mass but different ADM content move along roughly the same tracks in an HR diagram at a faster pace, but they do so at different rates. (We will discuss stellar tracks in § 4.4 and population isochrones in § 4.5.)

Previous work by ? found that solar mass stars in dense ADM environments may experience dramatic oscillations in both their core structure and surface properties, due to the large amounts of energy that ADM removes from the center, but noted that they were unable to determine whether it was a physical effect or a numerical artifact. Our initial simulations, using MESA release 10398, resulted in similar oscillations. In those models, ADM energy transport caused the temperature profile to become inverted, with the central temperature falling below the ADM temperature. Once this occurred, ADM began transporting energy back toward the center of the star. This raised the central temperature until it surpassed the ADM temperature again, causing ADM energy transport to again reverse direction and move energy away from the center. This cycle was self-reinforcing, causing large oscillations in the core temperature, density, etc., which propagated outward and resulted in large oscillations in surface properties as well.

Upon further investigation, we found that these simulations had poor energy conservation due to numerical artifacts. A new MESA version had been released since we had begun this work that included improved energy conservation schemes. When we updated to MESA release 12115 and ran these models again, the energy conservation was much improved, and the central temperature was reduced such that it was very close to the ADM temperature but never dropped below it. Since  $T_c > T_{\text{DM}}$  throughout the star’s lifetime, ADM energy transport never reverses direction and the oscillations seen previously are absent. We conclude that the oscillations in our initial simulations were a numerical artifact.

#### 4.2 High-Mass Stars

In standard models, MS stars with  $M_{\star} \gtrsim 1.3 M_{\odot}$  are powered primarily by the CNO cycle. This has several important consequences: (1) the burning rate is much higher than in pp-dominated stars; (2) the burning rate is extremely sensitive



**Figure 2.** Same as Figure 1 for  $3.5 M_{\odot}$  models, except that the 3rd panel in each set shows  $\log(D)$ , where  $D$  is the diffusion coefficient for convective mixing in  $[cm^2/s]$ . In the NoDM model the convective core retreats toward the center over time, and the burning rate peaks at the center until the end of the main sequence when the burning rate drops dramatically and a shell of strong burning appears suddenly. In the  $\Gamma_B = 10^6$  model, convection at the very center shuts off relatively early in the MS and a convective shell retreats away from the center over time. The peak burning rate shifts gradually outward, following the inner edge of the convective shell. The  $\Gamma_B = 10^6$  model reaches the  $X_c$  evolutionary markers at younger ages, relative to the NoDM model, since convection cannot replenish the fuel at the center.

to core temperature; (3) temperature gradients in the stellar core are relatively steep; and (4) stellar cores must be convective in order to carry away the energy produced by core hydrogen burning. Convective energy transport in the star also replenishes the core with unburnt hydrogen as the star evolves. Once hydrogen throughout the convective zone is depleted, the burning rate rapidly decreases and the star loses more energy at its surface than is being generated by burning. Gravity temporarily overcomes pressure support and the star contracts until the internal temperature increase is sufficient to ignite hydrogen in a shell outside the depleted core. This restructuring produces the so-called "convective hook" in an HR diagram as the star leaves the main sequence (?).


If a star captures enough ADM, the combination of dark matter + radiative energy transport becomes sufficient to carry the flux from nuclear burning at a temperature gradient that is insufficient to support convection. In other words, the additional energy transport by ADM can turn off convection within the stellar core. This can be seen in Figure 2 for a  $3.5 M_{\odot}$ ,  $\Gamma_B = 10^6$  star. Convection disappears from the center first (where ADM energy transport is most efficient) and retreats away from the core, into a narrowing shell. Without convective mixing, the hydrogen fuel supply depletes first at the very center (instead of simultaneously throughout the core) and the burning concomitantly shifts gradually into a shell, following the lower boundary of the convective zone. This can be seen in the time progression (down the page) of the  $\Gamma_B = 10^6$  (dark blue) model in Figure 2. The shift to shell burning is similar to the behavior of standard *low mass* stars.

#### 4.3 Main Sequence Lifetimes

#### 4.3 Main Sequence Lifetimes

In Figure 3 we summarize the effects of ADM on main sequence (MS) lifetimes relative to a standard NoDM star of the same mass. For the purposes of this [plotpaper](#), we have defined the MS to end when the fractional abundance of hydrogen in the center,  $X_c$ , falls below  $10^{-3}$ .  $X_c$  declines rapidly. Once  $X_c < 10^{-3}$  the hydrogen burning rate is greatly reduced and the star transitions out of the MS and onto the sub-giant branch. This transition period is marked by relatively sudden and dramatic changes to the star's structure. Stars that capture large amounts of ADM can have significantly different core structures at the end of the MS, so our results are not strongly affected by the exact choice—than their standard model counterparts, and these differences affect the stars' transition out of the MS, including the duration, in ways that are qualitatively different than ADM's effect on the MS itself. Therefore different choices in the definition of when a star leaves the MS can affect the results. Our relatively moderate choice of  $10^{-3}$  highlights changes in the core of the star during the bulk of the MS, rather than changes in the transition period between the MS and the sub-giant branch. We discuss ADM's affects on this transition period in § 4.4.

Lifetimes of low-mass stars, which have radiative (not convective) cores even in models. The MS lifetimes of relatively low-mass stars (near  $\sim 1 M_{\odot}$ ) can be altered by up to 20%. The, however the sense and degree of the shift in MS lifetime for relatively low-mass stars near  $M_{\star} \sim M_{\odot}$  is not a



**Figure 3.** Changes in MS lifetimes, relative to a star of the same mass with no dark matter, seen in our simulations. Diamonds mark the transition from radiative to convective cores (left to right). For the purposes of this figure this is defined as the lowest  $M_\star$  for which the average (over the MS) mass of the convective core is greater than 0.01 Mstar. Stars to the right of the NoDM marker (grey diamond) have decreased lifetimes due to a reduction in the size of the convective core, which reduces the amount of hydrogen available for burning. The effect abruptly disappears as stellar lifetimes become shorter than the time required to build up a sufficient amount of ADM. Stars to the left of this marker show mixed behavior due to the competing effects of decreased central burning rates and higher temperatures around  $m(< r) \gtrsim 0.1$  M $_\odot$  which give the star access to more fuel. In addition to these trends, there are several abrupt dips (e.g., at  $2.4$  M $_\odot$ ) and spikes (e.g., at  $2.55$  M $_\odot$ ). This is due to rotational mixing that turns on ~~in a shell~~ part-way through the MS and funnels fresh hydrogen fuel to the center, which increases the lifetimes (spikes). The dips result when the NoDM model exhibits this feature, but the ADM model of the same mass does not.

monotonic function of the strength of the dark matter effect, parameterized by  $\Gamma_B$ . This complicated dependence on the amount of captured ~~dark matter~~ADM is due to the competition between ~~increases burning rate~~increased burning rates and increased availability of burnable hydrogen fuel as discussed in § 4.1.

At higher masses, the influence of ADM on stellar lifetimes is clearer. ADM shortens the lifetimes of high mass stars ( $M_\star \gtrsim 1.3$  M $_\odot$ ). In NoDM models, the central convection zone extends beyond the burning region, giving the star a source of fresh nuclear fuel as hydrogen from outside of the core is mixed into the center. Since ADM shuts off convection in the center, the star no longer gets this influx of fresh hydrogen. Consequently, the star has less fuel available to burn, and so it leaves the MS faster than the NoDM models. Note that the appearance of a convective core (diamonds, Fig 3) shifts to higher masses with increasing  $\Gamma_B$  (diamonds in Figure 3) due to larger amounts of ADM which can carry larger energy fluxes. The effect disappears abruptly as  $M_\star$  increases because stellar lifetimes (which scale as  $M_\star^{-2.5}$ ) and quickly become too short for a sufficient amount of ADM to build up ~~in the star~~ (recall that the ADM capture rate scales roughly linearly with  $M_\star$ ), while the luminosity of the star increases rapidly with mass (roughly,  $L \propto M_\star^{3.5}$ ), meaning that more energy must be transported in order to alter the stellar structure.


In addition to the trends we have discussed, Figure 3 has

several abrupt dips (e.g., at  $2.4$  M $_\odot$ ) and spikes (e.g., at  $2.55$  M $_\odot$ ). This is due to rotational mixing that turns on ~~in a shell~~ part-way through the MS and funnels fresh hydrogen fuel to the center, which increases the lifetimes (spikes). The dips result when the NoDM model exhibits this feature, but the ADM model of the same mass does not. This rotational mixing occurs sporadically (i.e. at isolated masses, not in a continuous range of masses) in our models and may or may not be physical. However, it cannot be a bug introduced by our ADM energy transport module since ~~it happens in both ADM and some NoDM models. This type of behavior (stars of display the feature, but our module is not called in this case. This phenomenon where a star of a given mass experiencing increases in their central hydrogen fractions receives an influx of hydrogen to the center due to the onset of mixing, while stars of bracketing masses experience no such mixing~~ has also been reported previously in the MESA mailing lists<sup>5</sup>.

#### 4.4 Stellar Evolutionary Tracks and Isochrones

One of the goals of this work is to determine whether or not ADM can cause any gross changes to the properties of stars. We begin to answer this question with Figure 4, which shows

<sup>5</sup> <https://lists.mesastar.org/pipermail/mesa-users/>



**Figure 4.** Stellar evolution tracks, from ZAMS to core helium depletion ( $Y_c < 10^{-3}$ ,  $Y_c = 10^{-3}$ ), of select masses with  $\Gamma_B = 10^4$  (left) and  $\Gamma_B = 10^6$  (right). Tracks of NoDM models of the same mass are overplotted with a higher transparency. The chosen masses highlight some of the most dramatic changes seen in Figure 3. ~~x's~~ ~~Triangles~~ mark the location where stars leave the MS, which we define here as core hydrogen depletion below  $X_c = 10^{-3}$ . The location of the ZAMS and core hydrogen depletion for NoDM models are plotted as dotted and solid black lines respectively. The spike in the  $X_c, \text{NoDM} < 10^{-3}$  line near the  $2.5 M_\odot$  track is due to rotational mixing in the  $2.4 M_\odot$  model, which is discussed in § 4.3. The main effect of ADM on a star’s surface properties is to move the star through roughly the same sequence of events at a faster or slower pace, causing the offset of the  $X_c < 10^{-3}$ ,  $X_c = 10^{-3}$  milestone relative to NoDM. ~~High mass stars with sufficient ADM skip the convective hook because ADM shuts off convection in the core. These stars transition into shell burning, and therefore the sub-giant branch, more smoothly, similar to low mass stars.~~

evolutionary tracks on the HR diagram for many of our models. The tracks begin on the zero-age main sequence (ZAMS), delineated by the dotted black lines at the lower left of each panel. Stars evolve off of the ZAMS in a mass-dependent manner that is familiar from well-known aspects of standard stellar evolution. The tracks that we show leave the MS, defined as  $X_c < 10^{-3}$ , at the points marked by ~~crosses~~ ~~triangles~~. The points at which stars exit the MS in a standard model with no dark matter are indicated by the solid black lines in each panel. ~~(The spike near the  $2.5 M_\odot$  track is due to rotational mixing in the  $2.4 M_\odot$  model, which is discussed in § 4.3.)~~ Stars spend the majority of their lives on the MS and move more rapidly through the subsequent phases of stellar evolution. Our evolutionary tracks terminate when the core helium fraction falls below  $10^{-3}$ .

As is evident in Fig. 4, the effects of ADM on the evolutionary track of any individual star are generally subtle. Roughly speaking, this is not surprising. ~~The cooling induced by ADM effects only stellar cores. The chemical compositions, At a fixed central hydrogen fraction, stars containing ADM have different surface luminosities than their standard model counterparts, which is a result of ADM altering the structure of the star. This can be seen in the difference in location between the  $X_c = 10^{-3}$  markers in Fig. 4 (x's for ADM models, solid black line for NoDM models). However, if we instead consider stars at fixed luminosity, the temperature profiles, chemical compositions, opacities, and other proper-~~

ties of the overlying zones are approximately unaltered by ADM cooling. Consequently, the gross properties of the stellar photosphere, which are determined via the equations of stellar structure, are approximately fixed ~~at fixed luminosity. The result is that both ADM and standard model stars of a given mass follow roughly the same tracks in the HR diagram, but they do so at different rates.~~

Nonetheless, there are some small differences between standard evolutionary tracks and the tracks of stars with ADM. For example, consider the track of the  $1.75 M_\odot$  star in the left panel, corresponding to an environmental factor of  $\Gamma_B = 10^4$ . The standard model of stellar evolution shows a kink in the evolutionary track as the star exits the MS. This kink is known as the *convective hook*. The convective hook is caused by an overall contraction of the convective cores of the stars after hydrogen depletion. During this phase,  $T_{\text{eff}}$  increases. Eventually, at the hottest point on the hook, contraction of the former convective core is sufficient to ignite burning in a shell. After this point, shell burning ensues and the star continues to evolve along the sub-giant branch. What is clear from the evolutionary track of the  $1.75 M_\odot$  star in the left panel of Fig. 4 is that this evolutionary track exhibits *no convective hook*. This is because convection within the stellar core has been shut off by the ADM in these models. Instead of going through a phase of core collapse followed by shell burning, such stars make a smooth transition to shell burning and, thus, a smooth transition to the sub-giant branch.




Figure 5. Same as Figure 4 except that here we plot effective temperature as a function of stellar age. The sub-giant branch phase in a star's evolution is seen here as a sharp drop in  $T_{\text{eff}}$ , and is a result of structural changes in the star that are triggered by large reductions in the core burning rate due to hydrogen depletion at the end of the MS. The duration of the transition period between the MS and the sub-giant branch is seen here as the temporal difference between the locations of  $X_c = 10^{-3}$  (triangles) and the drop in  $T_{\text{eff}}$ . ADM alters the duration of the transition period, tending to increase it in high mass stars and decrease it in low mass stars. This is opposite of ADM's effect on MS lifetimes. The net effect is that the feature in  $T_{\text{eff}}$  always occurs either concurrently or earlier in the ADM model than in its NoDM counterpart.

The absence of convective hooks is evident for a wider range of masses in the right hand panel of Fig. 4, which corresponds to a larger environmental factor of  $\Gamma_B = 10^6$ .

The convective hook feature has been clearly seen in many open clusters for which the main sequence turn off lies between  $\sim 1.3 M_{\odot}$  and  $\sim 2 M_{\odot}$ , corresponding to stellar ages of  $\sim 1 \text{ Gyr}$  to  $\sim 4 \text{ Gyr}$ . However, this does not yet provide any strong statement about the nature of dark matter because none of those environments are thought to contain significant amounts of dark matter. Exploiting the convective hook feature would require the serendipitous identification of a stellar population of the appropriate age, associated with a significant amount of dark matter, and containing a sufficiently large number of stars such that the ephemeral convective hook and sub-giant phases of evolution are well sampled.

In Figure 5 we plot the effective temperatures with respect to stellar age to better understand the transition from the MS to the sub-giant branch, seen here as the difference between  $X_c = 10^{-3}$  (triangles) and the sharp drop in  $T_{\text{eff}}$ . A star's exit off of the main sequence is triggered when the core hydrogen fuel supply is depleted and the burning rate decreases such that it no longer provides sufficient pressure support and the star begins to collapse. Densities and temperatures increase until the bottom layer of hydrogen, now in a shell surrounding the core, ignites. The outward pressure resulting from increased shell burning causes the star's outer envelope to expand and cool at roughly constant luminosity, seen in Figure 5 as a large, sudden drop in  $T_{\text{eff}}$ . In models with no ADM, this transition period is more abrupt in

high mass stars due to the mixing induced by their convective cores, which causes hydrogen to become depleted throughout the core simultaneously and shell burning to appear suddenly (see § 4.2). In standard model low mass stars the cores are not convective, and so hydrogen depletes first at the very center and the burning shifts into a shell more gradually (see § 4.1).

The temporal difference between ADM and standard models in the location of the sudden drop in  $T_{\text{eff}}$  (when increased shell burning causes the envelope to expand) is another indicator of the change in MS lifetime. Unlike our definition of the end of the main sequence ( $X_c = 10^{-3}$ ) this indicator is based on surface properties and occurs towards the end of the large structural changes that happen during the transition period. In some cases (e.g., in the  $1.75 M_{\odot}$ ,  $\Gamma_B = 10^4$  model) this temporal difference is much smaller than the change in MS lifetime given by our definition of leaving the MS (seen here as the difference between the triangle markers of the ADM model and its standard model counterpart), and in other cases (e.g., in the  $1.0 M_{\odot}$ ,  $\Gamma_B = 10^6$  model) it is larger.

ADM can affect both burning rates and stellar structure (e.g., convection), and therefore it is not surprising that ADM affects the timescale of a star's transition off of the main sequence. High mass stars that skip the convective hook due to ADM energy transport take longer to move through this transition period because the burning shifts gradually into a shell (see § 4.2 for details). This behavior is very similar to standard model low mass stars. Conversely, the

$1.0 M_{\odot}$  AMD models move through this period more quickly than their NoDM counterpart. This is likely due to the fact that ADM has caused higher burning rates at the outer edge of the core during the MS, so that mixing during this transition period brings more helium into the center than in the standard model star. These shifts are opposite the shifts in MS lifetimes, and the net result is that this feature in  $T_{\text{eff}}$  always occurs either concurrently or earlier in the ADM model than in its standard model counterpart.

#### 4.5 Stellar Population Isochrones

Though the evolutionary tracks are quite similar across all models, ADM changes the rate of evolution and, hence, the stellar ages at which stars reach particular evolutionary stages. To convey some of the information that is obscured in an evolutionary track, we present stellar isochrones in Figure ?? and Figure ???. Each isochrone is a line on this plot that represents the locus of points that would be occupied by a population of stars of fixed age, but a wide range of masses.

The change in MS lifetimes due to ADM is seen here. The changes caused by ADM to individual stars' MS evolution is seen in these populations as a shift in the location of the MS turn-off (where the isochrones take a hard right turn, e.g., near the yellow triangle in Figure ??). Because the general effect on stars that are altered by ADM is to shorten MS lifetimes, . We have chosen these particular ages to maximize the sampling around this period and the subsequent crossing of the sub-giant branch. We do not show the giant branches because we do not have enough data points there to be meaningful, but we discuss the red giant branch further below. Due to the fact that stars move through the MS turnoff, sub-giant, and giant branches rather quickly, our mass sampling limits our ability to resolve these phases.


At around 1 Gyr, stars of  $M_* \sim 1.75 M_{\odot}$  are leaving the MS. In Figure 6, we see that the MS turnoffs around this time occur at a higher effective temperature and skip the convective hook, reflecting ADM's affects on these stars, discussed in § 4.2. These isochrones also tend to cross the turn-off happens at a lower mass and the stars evolve through the sub-giant branch at a lower luminosity. This causes isochrones of clusters in high  $\Gamma_B$  environments to appear older than their standard model counterparts. This becomes noticeable in the highest  $\Gamma_B$  environments around 0.2 Gyr as stars with  $M \approx 4 M_{\odot}$  begin to leave the MS. Prior to this, the stars have not had enough time to capture a sufficient number of DM particles for ADM driven energy transport to be a significant contribution to the overall energy transfer within the stars reflects the fact that ADM speeds up the evolution of these stars, and so stars with smaller initial masses, which have lower effective temperatures, are crossing the sub-giant branch earlier than they otherwise would. This happens in a wider range of ages in populations that live in richer ADM environments (right panel). Unfortunately stars move through this phase very quickly, meaning isochrones of real stellar populations are very sparsely populated in this region, which is known as the Hertzsprung Gap.

Younger isochrones display a sharp feature called the convective hook near the MS turn-off, which is noticeably absent from the  $\Gamma_B = 10^6$  isochrones. Since the models have convective cores, hydrogen is depleted nearly simultaneously throughout the volume of the core at the end of the MS.

This causes The oldest isochrones contain only low mass stars, since higher mass stars have already evolved into the giant branches and beyond. Here, the burning rate to decrease dramatically, which reduces the pressure support and results in gravitational contraction. The contraction increases the temperature until the bottom layer of hydrogen (now in a shell surrounding the core) is hot enough to ignite, reestablishing the pressure support. Our high ADM isochrones look very similar to their standard model counterparts, except that populations in high  $\Gamma_B$  isochrones lack this convective hook because ADM shuts off core convection. This lack of mixing causes the hydrogen to deplete first at the very center, and the burning gradually shifts into a shell, avoiding the sudden instability that causes the convective hook. This gradual shift to shell burning is very similar to the low mass models (which also do not experience a convective hook), which contributes to environments appear slightly older due to their decreased luminosity and temperature. This indicates that ADM causes the stars' surface properties to evolve more quickly, likely due to the increase in shell burning.

To better resolve the isochrone's MS turnoffs, Figure 7 shows the effective temperature of the high MS turnoff star, which we define as the hottest MS star at a given age. At younger ages there is no difference between the ADM and standard models because the stars have not yet captured enough ADM to be significantly affected. Around 0.15 Gyr, isochrones of the  $\Gamma_B = 10^6$  model start to display higher temperatures, remaining high until  $\sim 3$  Gyr, after which their temperatures are cooler than their NoDM counterparts. The  $\Gamma_B = 10^4$  isochrones show similar trends, but they occur at later times, since it takes longer for stars in lower  $\Gamma_B$  appearing older environments to build up sufficient ADM. The waviness in the lines at older ages is a result of limited mass resolution.

The gaps in the data are due to difficulty interpolating masses at a fixed time. Stellar masses generally increase when following a single isochrone from bottom to top. To build the isochrones from MESA's stellar models, MIST identifies the ages at which each star reaches a series of evolutionary milestones, and interpolates the stellar mass to generate properties of a group of stars at a fixed age. More massive stars generally evolve more quickly, and so the mapping between age and mass at a fixed milestone is usually monotonic, which the interpolation relies on. Exceptions to the monotonicity occur in mass regions where slightly less massive stars evolve more quickly. For example, in trends seen in our isochrones are consistent with what we were able to see in the isochrones generated by the MIST code, in regions where that code's interpolation was successful. In addition, we noticed from those isochrones that the tip of the red giant branch tends to occur at a lower luminosity in ADM models. The tip of the red giant branch is commonly used as a distance indicator, so ADM may add a source of uncertainty to these studies. To give the reader a sense of this shift, and to show a more well resolved MS turnoff and sub-giant branch, we show one particularly successful MIST isochrone for NoDM models, the emergence of a convective core around  $1.3 M_{\odot}$  results in stars just above this threshold living slightly longer than stars just below it. With large amounts of ADM, the increase in stellar lifetimes through



**Figure 6.** Isochrones for  $\Gamma_B = 10^4$  isochrones (left) and  $\Gamma_B = 10^6$  (right) models, marked by circles, with NoDM models overplotted as thin lines at higher transparency and marked by crosses. Triangles mark  $3.5 M_\odot$ , circles mark  $1.0 M_\odot$ . The positions of the markers here data points are very similar to interpolated from the ease stellar tracks to a common set of times (not shown isochrone ages). Gaps in We connect the interpolated data points of a single isochrone with straight lines to guide the eye. The lowest data point on every isochrone is a  $0.9 M_\odot$  star. As stars leave the main sequence they evolve rapidly, and therefore subsequent phases are less well sampled due to the difficulty interpolating in regions where mass resolution. Isochrone ages have been chosen to maximize the evolution sampling around the MS turnoff and sub-giant branch. We do not show the giant branches because we do not have enough data points there to be meaningful. The MS turnoff of isochrones around 1 Gyr happens at a higher effective temperature and a lower luminosity and skips the convective hook. This happens in a wider range of ages in populations that live in richer ADM environments (right panel). The oldest isochrones contain only low mass stars is faster than, and the ADM and NoDM populations look very similar except that of stars with populations in high  $\Gamma_B$  environments appear slightly higher masses (see § 4.5) older due to their decreased luminosity and temperature.

this transition to a convective core is much larger than in models and occurs over a wider mass range. These regions are not interpolated and appear as "missing" data in the isochrones and  $\Gamma_B = 10^6$  models in Figure 8. The lowest mass star in both isochrones (lower left) has  $M_* = 2.25 M_\odot$  (the interpolation was not successful for lower masses), and we show data through core helium depletion ( $Y_c = 10^{-3}$ ) which corresponds to  $M_* = 3.2 M_\odot$  in both cases.

PLACEHOLDER for MS turnoff plots, Teff and/or L. Main sequence turnoff temperature (temperature of the hottest star that is still on the main sequence). Residuals are with respect to (...?)


Paragraph(s) on MS turnoff plots (Teff and/or L as fnc of isochrone age)

## 5 DISCUSSION AND CONCLUSIONS

We have studied the potential impact of asymmetric dark matter interacting with nucleons through a spin-dependent coupling on the gross evolution of stars. We accomplished this by incorporating a module that approximates heat transport

by dark matter into the MESA stellar evolution software. We have identified several interesting qualitative distinctions between the standard evolution of stars and the evolution of stars in environments with a very high dark matter content. These include, 1) changes to the core temperature gradient, which alters the burning rates and stellar structures of low and high mass stars in qualitatively different ways; 2) low mass stars' MS lifetimes (defined as  $X_c > 10^{-3}$ ) are increased by as much as 20%; 3) high mass stars' MS lifetimes are reduced by as much as 40%; 4) stars in both mass regimes cross the sub-giant branch at younger ages and may reach the tip of the red giant branch at lower luminosities.

It is interesting to speculate on ways in which these effects could be used to identify and/or constrain dark matter or ways in which these effects may, at least, serve as an element of uncertainty in the analysis of stellar populations. Any constraint on dark matter arising from these effects requires very high-quality observations of a stellar population residing in an environment with a large ambient dark matter density and thus there will be a significant element of serendipity involved. If such a population is observed, our models suggest that if it is  $\sim 0.1$  Gyr old, the hottest MS



**Figure 7.** Same Effective temperature of the main-sequence-turnoff star, defined here as Figure ?? but for the hottest MS star, at a given age. We show only the NoDM,  $\Gamma_B = 10^4$ , and  $\Gamma_B = 10^6$  models to allow the reader to see them clearly. The MS turnoff temperature of the  $\Gamma_B = 10^4$  models starts to become hotter than in the NoDM isochrones between (insert ages) happens at a lower luminosity and skips  $\sim 0.7$  Gyr when stars of  $\sim 2 M_\odot$  begin leaving the convective hookMS, causing them while those of the  $\Gamma_B = 10^6$  models become hotter at  $\sim 0.1$  Gyr when stars of  $\sim 3.5 M_\odot$  begin leaving the MS. As we move to appear older relative to isochrones the effect is reversed, and ADM models have lower effective temperatures at the turnoff. The lines terminate when there are no more stars on the main sequence (the lowest mass in our set of models is  $0.9 M_\odot$ ). For the purposes of this figure, we exclude stars in the convective hook because this feature is not well-resolved in our isochrones (see Fig. 6).


star should be slightly *hotter* than expected for a population without ADM, and it should be slightly *colder* at  $\sim 1$  Gyr. In addition, the tip of the red giant branch may occur at a lower luminosity. The tip of the red giant branch is commonly used as a distance indicator so ADM may add a source of uncertainty to these studies. Finally, the metallicity is known to affect many of the properties we have discussed (e.g., the locations of various phases in the HR diagram), so ADM may be a contaminant here as well.

Future work along these lines includes 1) chemical abundance studies exploring the effects of altered core burning; 2) asteroseismology of Sun-like MS and red giant branch stars, which could be seen in (e.g.,) the small frequency separation—a diagnostic that is sensitive to the core structure of the star. 3) spin-independent ADM-nucleus scattering, which should have a larger effect during later phases when stars are burning helium.

This paper has been typeset from a T<sub>E</sub>X/L<sub>A</sub>T<sub>E</sub>X file prepared by the author.

## ACKNOWLEDGEMENTS

$T_{\text{eff}}$  as a function of age for select  $\Gamma_B$ , with with models overplotted as thin lines. Stars undergo a sharp decrease in  $T_{\text{eff}}$  as they leave the MS. The change in MS lifetimes due to ADM can be seen in the time difference between these features.



**Figure 8.** Isochrones generated by the MISTcode for NoDM (grey) and  $\Gamma_B = 10^6$  (dark blue) populations. The lowest mass star in both isochrones has  $M_* = 2.25 M_{\odot}$  (the interpolation was not successful for lower masses), and we show data through core helium depletion ( $Y_c = 10^{-3}$ ) which corresponds to  $M_* = 3.2 M_{\odot}$  in both cases. The  $\Gamma_B = 10^6$  isochrone skips the convective hook and crosses the sub-giant branch at a lower luminosity, consistent with Figure 6. Additionally, the tip of its red giant branch occurs at a lower luminosity, a trend which appears in most of the  $\Gamma_B = 10^6$  MIST isochrones with  $0.1 \text{ Gyr} \lesssim \text{Isochrone age} \lesssim 1 \text{ Gyr}$ .