

Observations of Galactic and Extragalactic Novae

Massimo Della Valle and Luca Izzo

Received: date / Accepted: date

Abstract The recent GAIA DR2 measurements of distances to galactic novae have allowed to re-analyse some properties of nova populations in the Milky Way and in external galaxies on new and more solid empirical bases. In some cases we have been able to confirm results previously obtained, such as the concept of nova populations into two classes of objects, that is *disk* and *bulge* novae and their link with the Tololo spectroscopic classification in *Fe II* and *He/N* novae. The recent and robust estimates of nova rates in the Magellanic Clouds galaxies provided by the OGLE team have confirmed the dependence of the normalized nova rate (i.e. the nova rate per unit of luminosity of the host galaxy) with the colors and/or class of luminosity of the parent galaxies. The nova rates in the Milky Way and in external galaxies have been collected from literature and critically discussed. They are the necessary ingredient to assess the contribution of novae to the nucleosynthesis of the respective host galaxies, particularly to explain the origin of the overabundance of lithium observed in young stellar populations. A direct comparison between distances obtained via GAIA DR2 and Maximum Magnitude vs. Rate of Decline (MMRD) relationship points out that the MMRD can provide distances with an uncertainty better than 30%. Multiwavelength observations of novae along the whole electromagnetic spectrum, from radio to gamma-rays, have revealed that novae undergo a complex evolution characterized by several emission phases and a non-spherical geometry for the nova ejecta.

Keywords novae, cataclysmic variables - distance scale - Nuclear reactions, nucleosynthesis, abundances - supernovae: general - Galaxy: stellar content

M. Della Valle
INAF-Capodimonte, Naples, Italy & ESO, Garching, Germany
E-mail: massimo.dellavalle@inaf.it

L. Izzo
DARK/NBI, Copenhagen, Denmark & IAA/CSIC, Granada, Spain
E-mail: luca.izzo@nbi.ku.dk

Contents

1	Observations of Guest Stars	3
1.1	First Theories	6
1.2	Modern Times	6
1.3	Classical Novae as Cataclysmic Variables	10
2	Introduction	12
3	Nova Populations in the Milky Way	14
3.1	Nova lightcurves: the speed classes	16
3.2	Disk and Bulge Novae	19
3.3	Post-Novae	23
3.4	The interaction of Nova with the Interstellar medium	26
3.5	The Galactic Nova rate	27
4	The Spectral evolution of Novae	29
4.1	The Tololo Spectroscopic Taxonomy	33
4.2	The THEAs: Transient Heavy Elements Absorptions	34
5	The Maximum Magnitude vs. Rate of Decline Relationship in the Milky Way (MMRD)	37
5.1	The Shape of the MMRD Relationship	41
5.2	How good is the MMRD?	44
6	Nova populations in external galaxies	45
6.1	Novae in M31	45
6.2	Novae in M33	49
6.3	Novae in the Magellanic Clouds	50
6.4	Novae in M81	53
6.5	Novae in Virgo and Fornax Ellipticals	53
6.6	Nova Rates as function of the Hubble type of the hosts	54
6.7	Nova outbursts as a function of the Metallicity content and class of luminosity of the Hosts	57
7	Novae in Globular Clusters	57
8	The MMRD Relationship in Galaxies of different Hubble types	58
8.1	Outliers of the MMRD	59
9	Novae as Distance Indicators	65
9.1	The distance to M31, LMC and Virgo	65
9.2	Novae as Standard Candles	66
10	The role of Novae in the Galactic Nucleosynthesis	67
10.1	The mass of the ejecta	68
10.2	The abundances in the nova ejecta	69
10.3	Novae as Lithium Factories	70
10.4	"The Hunt for Red Lithium"	72
11	Novae as Supernova-Ia progenitors	73
12	Multi-wavelength Novae	77
12.1	Novae in the Infrared	77
12.2	X-ray Novae	78
12.3	Gamma-ray Novae	79
12.4	Novae at Radio frequencies	81
13	Conclusions	82
A	Tables	107

1 Observations of Guest Stars

In the ancient chronicles, particularly in the almanacs of the Chinese astronomers, all "temporary" or "transient" phenomena that suddenly appeared in the sky, to then fade and disappear after days or months, were generally indicated with the name of "guest stars" or "visitor stars". This definition was very broad and it was used to classify a large number of astronomical events such as comets, planets in opposition or conjunction, and new stars. In those days there were no clues about the different origin of these phenomena and the common idea was that they were somehow related to each others. In Europe these events were also reported as "*Stella Nova*", the Latin name for "New Star" (*Stellae Novae* in his plural form). Unlike the locution *Supernova* that is a recent term, coined in the 30's, *Stella Nova* in astronomy was used for the first time about 2,000 years ago in the Natural History of Pliny the Elder. In his second book¹ dedicated to Astronomy and Meteorology, Pliny reports:

"Hipparchus himself ... detected a "new star" that led him to wonder whether this was a frequent occurrence" and "he dared to prepare a list of stars for posterity to indicate their positions and magnitudes, in order that from that time onward it might be possible to discern not only whether stars perish and are born, transit and in motion, and also whether they increase and decrease in magnitude"

To preserve the Aristotle's dogma on the perfection and immutability of the heavens, in which *Stellae Novae* obviously did not fit, the ancient astronomers considered nova stars sub-lunar phenomena or even Earth's atmosphere occurrences. Therefore, in this view, these objects did not belong to the "octave sphere" the so called "sphere of fixed stars": the skies beyond Saturn were indeed immutable by definition. Despite these ideological prejudices, the observation of new stars was never abandoned, and therefore today we know, at least approximately, the positions in the sky of many remnants of those stellar explosions and some of them have been targeted with modern instrumentation, (see for example Shara *et al.*, 2017b). The reason for which the ancient observers keep alive the interest in observing these phenomena was simple. Until few centuries ago, very little or even nothing was known about the real nature of the stars and planets. According to the Italian poet, Giacomo Leopardi (1798-1837):

"the knowledge of the effects and the ignorance of the causes produced astrology",

therefore we should not be surprised to learn that in those times astrology and astronomy were indistinguishable. They were both part of a culture that considered the earthly events announced in the sky through the manifestation of strange and unusual phenomena, such the sudden appearance of comets or "new stars". We can only imagine what was the apprehension or excitement in the naive and unprepared observers of those strange phenomena. Due to the way in which nova stars appeared in the sky, suddenly and without any warning signs, these events were often considered omens of misfortune, so their observational follow-up was extremely important: it could help in predicting the "unknown" that a

¹ Out of 37 that constitute his entire work

little later would have happened on Earth. This climate is well represented by the chronicles of observations made in the Benedictine Abbey of St. Gall, today in Switzerland, about the "nova star" of 1006 that we know today to have been a Supernova explosion indeed. The monks reported an immense light that *blinded sight* and aroused *a certain fear*.

Novae and Supernovae were not considered two distinct families of objects until last century. The rapidly increasing number of nova discovery in Spiral Nebulae due to systematic searches (Ritchey, 1917a,b, 1918; Curtis, 1917b,c; Pease, 1917; Duncan, 1918; Sanford, 1918, 1919) started to cast the first doubts about a common origin of these phenomena. In those years, novae played a pivotal role in the debate about the galactic or extra-galactic nature of the nebulae. For example, in his paper on "Novae in Spiral Nebulae and the Island Universe Theory" Heber Curtis at the end of his short note (Curtis, 1917a) correctly inferred:

"If we assume equality of absolute magnitude for galactic and spiral novae, then the latter, being apparently 10 magnitudes the fainter, are of the order of 100 times as far away as the former."

However, this simple conclusion was questioned by the strange case of S Andromedae (And). This *nova* occurred in M31 in 1885 and today is classified as sub-luminous SN-Ia. The 8th magnitude achieved by S And at peak luminosity was much brighter than the apparent magnitudes of all other nova stars in Andromeda Galaxy and in other spiral Nebulae. As noted by Shapley (1917):

"... the average (magnitude) for the eleven (novae) given in the table is at least magnitude -7, and the maximum for S Andromedae is at least -15, a luminosity nearly a hundred million times that of our Sun, the equivalent of the light emission of a million stars of zero absolute magnitude, and probably, therefore, much greater than the total light of all the stars seen with naked eye ..."

an argument that led him to conclude:

"... hence stellar luminosities of these orders seem out of the question and accordingly the close comparability of spirals containing such Novae to our Galaxy appears inadmissible ...".

For many decades novae have been used as rulers of distance outside our Galaxy (Pritchett and van den Bergh, 1987; Della Valle and Livio, 1995; Shara *et al.*, 2018), so it appears somehow ironic that the erroneous interpretation of S And in terms of nova explosion has significantly delayed the correct measurements of the distance to M31 and more generally the correct interpretation of the spiral Nebulae as systems external to the Milky Way. However this conundrum was taken by Lundmark (1923) the other way around, he assumed that spiral Nebulae were extragalactic systems and in a series of paper published between 1920 and 1923 he realized first that *"... Spiral Nebulae were extragalactic objects ...some Nova got very bright..."*, and therefore *"...a division into two magnitude classes is not impossible."* (see Fig. 1).

The brightest class of explosive variables was labelled in the following decades in different ways. Lundmark (1927) called them upper-class Novae, while Hubble (1929) calls S Andromedae *"exceptional case"* and finally Baade and Zwicky (1934)




Fig. 1 Number of Novae brighter than magnitude m . Plot from Lundmark (1923).

proposed the definition currently adopted of *Super-novae*. In the following years the hyphen has disappeared. Lundmark (1935) pointed out the existence, at the faint tail of the luminosity distribution of novae, of two other groups of "temporary stars": the U Gem or Dwarf Novae and nova-like objects. He also provided the first classification in terms of luminosity at maximum for the whole family of explosive variables:

- A. The upper-class novae or super-novae, having an absolute magnitude at maximum (M_{max}) around -15 mag;
- B. The middle-class novae, or ordinary novae, with an absolute magnitude at maximum very close to -7 mag;
- C. The lower-class novae, or the dwarf novae, having an absolute magnitude at maximum around +3 mag or +4 mag.

Lundmark also provided the first estimate of their frequency of occurrence in the Milky Way: 1 in 50 years for super-novae and about 50 ordinary novae per year. "Thus the ordinary Novae are at least 2500 times more frequent than the super-Novae." We note that current estimates of this ratio for spirals galaxies range between $2 \times 10^3 - 2 \times 10^4$ (Della Valle and Livio, 1994).

1.1 First Theories

Since the times of Riccioli (1596-1671), Hevelius (1611-1687) and Fabricius (1564-1617) many theories were developed to explain the nature of the nova phenomenon. Gian Battista Riccioli, in the ninth book of its encyclopedic work "*Almagestum Novum*", dedicated to "Comets and New Stars", listed a dozen of theories to explain the nature of these transients phenomena. At that time the physical understanding of nova explosions was completely inaccessible, so it is not surprising that the theories presented in the Riccioli's work appear today as naive and extravagant. For example, it was assumed that these "temporary" stars were not "new" but in fact they were "old" and that they were hidden in the depths of the ether, invisible to us, because of their vast distance. These stars were supposed to be in motion and when they approached the Earth, they became visible and when they moved away they became invisible again. This idea was initially conceived to explain the behaviour of Tycho's new star, but it was also used to explain the appearance and disappearance of periodic stars as Mira Ceti. Many other theories were devised in the following centuries and in some cases they were characterized by more physical insight with respect to the oldest ones. For example de Maupertuis (1732) suggested that the appearance and disappearance of new stars could be originated by the unbalanced action of planets around stars that were flattened by their very high angular momentum. When the geometrical alignment between the flattened star and the observer turned from edge-on to face-on, the observer could measure a dramatic change in the brightness of the star. According to Newton (1729), novae were old stars that had exhausted their reserve of light and vapors and therefore were not anymore visible in the sky. They could have returned to "new splendor" after being fed with fresh supply of gas provided by comets that fell upon them. In this case the idea that a nova was an "old" star that could be rejuvenated or reinvigorated by gas coming from another external celestial body, appears remarkable and not distant, at least in its general principle, from modern ideas, in which the fuel source are not comets but a companion star. Finally, we note the very imaginative hypothesis elaborated by Monck (1885) to explain the nature of S Andromedae:

"Gaseous nebulae sometimes occupy degrees in the sky ... The rush of a star through a gaseous nebula is therefore no improbable event and it is certain that if occurred the star would be highly heated, and its light greatly intensified. An idea that was borrowed by shooting stars events that are known to be dark bodies rendered luminous for a short time by rushing through our atmosphere ... "

1.2 Modern Times

We had to wait for the 20th century, particularly the 50's and 60's to come to understand the Nova phenomenon in its general lines. The beginning of "modern times" might be set by the work of Schatzman (1951) who highlighted the role of ^3He in triggering "*une onde de detonation*" that might be at the origin of the Nova phenomenon. The following steps, i.e. the discovery that DQ Her was a binary system (Walker, 1954) and that the binary condition was common to

all cataclysmic variables including Novae (Kraft, 1964), laid the physical basis to conclude that Nova outbursts are related to mass accretion phenomena onto the surface of degenerate hot companion (Kraft, 1965; Paczyński, 1965). All previous items, complemented with the first hydrodynamic studies of Nova ejections (Sparks, 1969), provided the pillars on which the modern studies on Nova explosions have then progressed (Starrfield *et al.*, 1975; Prialnik *et al.*, 1978; Nariai *et al.*, 1980).

The current framework for Classical Novae establishes that they are explosive events caused by the ignition of the thermonuclear runaway (TNR) on the surface of a white dwarf that is accreting material, under degenerate conditions, from a low main sequence companion (e.g. Gallagher and Starrfield, 1978). Most novae are semi-detached binary star systems where a late-type main sequence star, which has filled its Roche-lobe, transfers mass through the inner L_1 Lagrangian point, to the white dwarf companion. When the gas falls onto the compact object, because of its angular momentum with respect to the accreting star, it will not fall directly onto the white dwarf but will start orbiting around it. Gas particles, having different orbits will collide with each other and, shortly after their bulk motion will be circularized through friction and viscosity processes, the gas will eventually form an accretion disk before falling onto the compact object. In passing we note that this process can be partially or totally inhibited by the presence of strong magnetic fields (Wickramasinghe *et al.*, 2014). If the mass accretion rate onto the white dwarf is sufficiently low the accreted gas becomes degenerate, and both pressure and temperature in the accreted envelope will dramatic increase. Finally, TNR will ignite at the base of the accreted envelope leading to a nova eruption. The explosion is not destructive for the white dwarf, therefore nova explosions are "recurrent" and the period of time between two explosions can be as short as a few months (Kato *et al.*, 2014) and as long as $\sim 10^5$ years. The length of the time scale marks the difference from the so called "Recurrent Novae", (RNe) i.e. Novae for which at least two outbursts have been recorded (typically RNe show more explosions over the lifetime of an astronomer) and Classical Novae (CNe) characterized by much longer interval of time between two explosions.

In a very simplified scheme a white dwarf composed of either carbon-oxygen (CO) or oxygen-neon-magnesium (ONeMg) accretes hydrogen rich material from a companion star at a rate of $10^{-9\pm1} M_\odot/yr$ and as the matter piles up onto its surface, it becomes dense and hot. It is heated at its bottom mainly by gravitational compression. When the pressure and the temperature at the bottom of the accreted layer reaches the critical values of

$$P_{crit} = \frac{GM_W D \Delta m}{4\pi R^4}, \quad (1)$$

of $\approx 10^{19} - 10^{20}$ dyne cm^{-2} (Fujimoto, 1982; Truran and Livio, 1986) and $T \sim 10^7 K$, the nuclear reactions ignite. In fact theoretical calculation for realistic nova outbursts show that P_{crit} also depends on other quantities such as \dot{M} and ΔM_{ign} (Kato *et al.*, 2014). These reactions produce short-lived (about 100s) β^+ radioactive nuclei that can be transported by convection in the outer envelope. This is efficiently heated up, leading to higher temperatures, which in turn causes more nuclear burning. Eventually, the temperature rises to the point where the layers above the nuclear burning are no longer degenerate and a TNR flash occurs

Mass WD (M_\odot)	Interval (10^5) years
0.60	12.9
0.70	7.3
0.80	4.2
0.90	2.4
1.00	1.2
1.10	0.64
1.20	0.28
1.30	0.09
1.35	0.04

Table 1 Nova recurrence time scale (table from Truran and Livio, 1986).

then expelling the ejecta. At the critical pressure, it corresponds a critical ignition mass, which is a function of the radius of WD and its mass:

$$\Delta M_{ign} \approx \frac{4\pi P_{ign}}{G} \frac{R_{WD}^4}{M_{WD}}. \quad (2)$$

Therefore for a given \dot{M} , the recurrence time:

$$\Delta T_{rec} \sim \Delta M_{ign} / \dot{M} \sim R^4 / M_{WD} \quad (3)$$

is proportional to R^4 / M_{WD} .

From the radius of the WD expressed as a function of its mass (Eggleton, 1983):

$$R_{WD} \approx 8.5 \times 10^8 \left[1.286 \left(\frac{M_{WD}}{M_\odot} \right)^{-2/3} - 0.777 \left(\frac{M_{WD}}{M_\odot} \right)^{2/3} \right]^{1/2} \text{ cm}, \quad (4)$$

we derive that the more massive WDs, e.g. $1.35 M_\odot$, are characterized by recurrence time shorter by a factor 282 with respect to light WDs with masses of $0.6 M_\odot$. This first order computation is in good agreement with the recurrence times scale obtained by several authors (Truran and Livio, 1986; Politano *et al.*, 1990; Ritter *et al.*, 1991), see also Table 1, for which $\Delta T_{0.6} / \Delta T_{1.35} = 322$.

Yaron *et al.* (2005) have computed an extended grid of multi-cycle nova evolution models and obtained recurrence times at varying the mass of and temperature of the WD and \dot{M} . For a typical value of $\dot{M} = 10^{-9} M_\odot / \text{yr}$ we consistently derive a ratio of 390. Accurate grids provided by Kato *et al.* (2014) computed for a range $10^{-6} < \dot{M} < 10^{-9} M_\odot / \text{yr}$ give consistent results within a factor about two for $\dot{M} \approx 10^{-9} M_\odot / \text{yr}$, see Fig. 2. Now we can estimate on the basis of simple arguments the duration of the nova duty cycle. A system formed by a low main sequence companion of $0.5 M_\odot$ can experience about $0.5 M_\odot \times \alpha / M_{env}$ outbursts. The parameter α is a coefficient that accounts for the efficiency of the accretion through L_1 the inner Lagrangian point, which is estimated to be of the order of 20% (see Sect. 12). For $M_{env} \sim 10^{-5} M_\odot$ and $\sim 10^{-4} M_\odot$ for fast and slow novae (see Sect. 10) we derive about $10^{4 \div 3}$ outbursts during the nova cycle, consistent with the "classical" estimates (Bath and Shaviv, 1978).




Fig. 2 The distribution of recurrence periods of classical novae compared with the accretion rate and the mass of the primary white dwarf, according to the simulations by Kato *et al.* (2014).

Thus the mean lifetime for nova systems turns out to be $10^{4 \div 3} \times \Delta T_{rec}$, therefore $\sim 10^7$ years and $\sim 10^9$ years for massive and light nova progenitors, respectively. The connection among M_{WD} , \dot{M} , and M_{env} was explained by several authors e.g. Nomoto (1982a) (Fig. 3) and revised in recent years by Kato *et al.* (2014) (Fig. 4). The more massive WDs expel the lightest envelopes with the highest ejection velocities, up to 4,000-5,000 km/s in the most extreme cases. In many post-novae a "quiet" H-burning continues in a kind of steady state then originating the so called SuperSoft X-ray phase (Oegelman *et al.*, 1984; Orio *et al.*, 2001; Henze *et al.*, 2010), see also Sec. 12.2.

The physical processes regulating the ejection of the gas are still matter of debate. The entire evolution of a nova outburst can be explained in terms of a continuum radiation wind (Kato and Hachisu, 1994) or through an homologous ballistic ejection of a discrete shell (Shore *et al.*, 1996). More recently, the detection of high-energy radiation from novae combined with radio and optical high-resolution observations led to the proposal of a "multi-phase" ejection mechanisms for novae. The current evidence is consistent with an early time ballistic ejection, with a toroidal geometry distribution and a late time bi-polar faster wind (Mukai and Sokoloski, 2019). Their interaction would give rise to shocks and then to the observed high-energy radiation (Metzger *et al.*, 2015).




Fig. 3 Masses of the accreted envelopes as a function of the mass of the WD and \dot{M} . Figure taken from Nomoto (1982a).

1.3 Classical Novae as Cataclysmic Variables

Classical Novae fall in the faint tail of stellar explosions that occur in galaxies. With an emitted radiant and kinetic energies of about 10^{45} erg (Gallagher and Starrfield, 1978). They are many orders of magnitude less energetic than gamma-ray bursts (Amati, 2006), Kilonovae (Abbott *et al.*, 2017) and Supernovae (Gal-Yam, 2012). In terms of magnitude at maximum these eruptions can reach an absolute magnitude as bright as Nova LMC 1991, $M_V \approx -10$ mag (Della Valle, 1991; Schwarz *et al.*, 2001), approximately one order of magnitude less luminous than Luminous Red Novae (Pastorello *et al.*, 2019).

Novae belong to the very heterogeneous family of semi-detached binaries and different phenomena are observed according to the nature of the receiving star. If it is a main sequence (MS) star, we have an Algol-type eclipse system characterized by periods of hours/days. If it is a collapsed object, as a white dwarf (WD), we enter in the heterogeneous realm of Cataclysmic Variables (CVs), normally characterized by periods of few hours (but see GK Per, Bianchini *et al.*, 1981). With the limits, that are inevitable when we proceed to a schematization, we can broadly divide the CVs family in different sub-classes according to masses of the companion stars and the amount of mass transfer rates. Decades of studies on CV properties effectively summarized by Warner (2003) converge toward the following "outline":




Fig. 4 The distribution of the ignition mass as a function of the accretion rate and the mass of the primary white dwarf, according to the simulations by Kato *et al.* (2014).

- if the donor star is a low mass MS star and the receiving star is a WD ($0.7M_{\odot}$) and $\dot{M} \sim 10^{-11} M_{\odot}/\text{year}$, the binary system is labeled as Dwarf Nova (DN);
- if the donor star is a low mass MS star and the receiving star is a high mass WD ($0.7M_{\odot} < M < 1.1M_{\odot}$) and $\dot{M} \sim 10^{-9} - 10^{-10} M_{\odot}/\text{year}$, the binary system is a Classical Nova (CN);
- if the donor star is a MS star (or a giant) and the receiving star is a high mass WD ($\sim 1.3M_{\odot}$) and $\dot{M} \sim 10^{-7} - 10^{-8} M_{\odot}/\text{year}$ the system is a Recurrent Nova (RN);
- if the donor star is a low mass MS star and the receiving star is a NS/BH, then, the system is a Low Mass X-ray Binary (LMXB);
- if the donor star is a high mass MS (e.g. blue giant) star and the receiving star is a NS/BH, the system is a High Mass X-ray Binary (HMXB);
- if the donor star is a low MS star and the receiving star is a very high mass WD ($> 1.3M_{\odot}$) the transfer can bring the WD across the Chandrasekhar limit and the WD explodes as SN Ia (Whelan and Iben, 1973; Nomoto, 1982a,b; Starrfield *et al.*, 2019);
- if the donor star is another WD, the coalescence can bring the binary system across the Chandrasekhar limit and the double degenerate (DD) system will explode as SN Ia (Iben and Tutukov, 1984, 1985). As an alternative the coalescence of two WDs may not result in a SN-Ia explosion, but eventually producing an “accretion-induced-collapse” to a neutron star (Nomoto and Kondo, 1991).




Fig. 5 The number of novae vs. year of discovery. The total number of novae plotted in this figure can be found at the *Project Pluto* website (<https://projectpluto.com/galnovae/galnovae.txt> maintained by Bill Gray).

- if the donor star transfer Helium onto the WD it can explode as SN Ia by igniting off-center at sub-Chandrasekhar masses (Yungelson and Livio, 1998), or feeding the WD in a Helium-rich supersoft X-Ray system ((Kato and Hachisu, 1989), Hachisu *et al.* (1999)) and typified by the Recurrent Nova U Sco, in which the WD grows in mass to the Chandrasekhar limit and may explode as SN Ia (Kato et al. 2008).

In the 80's it was proposed (Shara *et al.*, 1986) an unified scenario in which Classical Novae and Dwarf Novae were the same systems observed at different evolutionary stages (see sect. 3.3).

With "Nova-like" we label all poorly studied cataclysmic variable stars which resemble Novae due to their photometric behaviour at maximum or minimum or due to their spectral development. It is not infrequent that as soon as more observations are collected it is possible to reclassify some "Nova-like" objects as belonging to one of the above reported classes.

2 Introduction

About 400 novae have been observed in the Milky Way since CK Vul 1604 and WY Sge 1783, but only a fraction of them has been observed into some details sufficient to carry out interesting analysis (Fig. 5).

Collections of Nova data are periodically updated, reviewed and enriched, see for example (Cecchini and Gratton, 1942; Duerbeck, 1981, 1987; Özdönmez *et al.*, 2018; Hachisu and Kato, 2018), (Schaefer, 2018) - S18 hereafter - and Selvelli and Gilmozzi (2019) - SG19 hereafter - and some data from these papers have been used in this work. Reviews on novae have not been very popular since the

exhaustive paper by Gallagher and Starrfield (1978). More recently, we notice there has been an increased interest in this field: the sections 5-8 in the book "Cataclysmic Variable Stars" by Warner (2003), the book *Classical Novae* by Bode and Evans (2008), and the more recent reviews by Woodward and Starrfield (2011), Poggiani (2018a), Darnley *et al.* (2019) and (Shafter, 2019).

Spectroscopic and photometric data for historical Novae have been extracted from the classical books "The Galactic Novae" by Payne-Gaposchkin (1957) and "Classical Novae" by Bode and Evans (2008), while more recent data have been directly derived from literature, e.g. (Ribeiro *et al.*, 2013b; Munari *et al.*, 2017). However, both historical and new records have been revised according to the Tololo Nova Classification in the spectroscopic "types" Fe II and He/N Novae (Williams *et al.*, 1991; Williams, 1992, 1994; Williams *et al.*, 1994). The modern spectroscopic classification, usually based on low-intermediate resolution, was subsequently integrated by high resolution observations ($R \sim 50,000$) carried out at ESO-La Silla at the end of 90's for Milky Way and Magellanic Clouds Novae (Della Valle *et al.*, 2002; Mason *et al.*, 2005). These high resolution data set have provided the observational basis for the detections in the nova ejecta of the so called Transient Heavy Element Absorptions (THEAs, Williams *et al.* 2008) and ${}^7\text{Li}/{}^7\text{Be}$ (Izzo *et al.*, 2015; Tajitsu *et al.*, 2015; Molaro *et al.*, 2016; Izzo *et al.*, 2018; Molaro *et al.*, 2020). Lithium and Berillium were ones of the most important and puzzling missing pieces of chemical evolution of the Milky Way (Romano and Matteucci, 2003) and their detection have significantly contributed to assess the role of Novae in the Galactic Nucleosynthesis.

Systematic surveys to search for Novae in external galaxies have a long historical record. Pioneer surveys on photographic plates were mostly devoted to study the nova population in M31 (Hubble, 1929; Arp, 1956; Rosino, 1956, 1964, 1973; Rosino *et al.*, 1989), M33 (Rosino and Bianchini, 1973), and in the LMC (Graham, 1979). We note that the former surveys were often supplemented by more fragmentary observations (Sharov, 1979) or by "new concept" surveys, as the one carried out on the M31 bulge in $\text{H}\alpha$ narrow band filter with one-meter class telescopes equipped with CCD detectors (Ciardullo *et al.*, 1987). Photographic data obtained in the 50's Palomar campaign for the discovery of classical novae in M81 and in the early 70's at Asiago Observatory for studying novae in M33, were published only in the 90's (Shara *et al.*, 1999; Della Valle *et al.*, 1994). The modern search for Novae in extra-galactic systems has received a significant boost of interest after the seminal paper by Pritchett and van den Bergh (1987) dedicated to study Novae in the Virgo Cluster. The paper was a "classical" search for Novae in B filter, carried out with a four-meter class telescope (the Canada-France-Hawaii Telescope, CFHT) equipped with CCDs. A work that, despite of only 9 novae discovered in 15 half-nights run, has considerably influenced the subsequent studies in the extragalactic nova topics including the distance scale debate (Jacoby *et al.*, 1992; van den Bergh, 1996). In the following years surveys for Nova searches in extragalactic systems (corroborated by some spectroscopy) have been carried out in M31 (Tomaney and Shafter, 1992; Darnley *et al.*, 2004, 2006; Shafter and Irby, 2001; Shafter *et al.*, 2011; Kasliwal *et al.*, 2011; Cao *et al.*, 2012), M51 and M101 (Shafter *et al.*, 2000), in NGC 1316 (Della Valle and Gilmozzi, 2002), M49 (Ferrarese *et al.*, 2003; Curtin *et al.*, 2015), M33 (Shafter *et al.*, 2012), M87 (Shafter *et al.*, 2000; Madrid *et al.*, 2007; Curtin *et al.*, 2015; Shara *et al.*, 2016), M84 (Curtin *et al.*, 2015), NGC 2403 (Franck *et al.*, 2012). Finally we point out the

first homogeneous survey on novae in the Magellanic Clouds (Mróz *et al.*, 2016) since Graham (1979). Sporadic observations carried out on M32 and NGC 205 (Neill and Shara, 2005) have discovered one or possibly two nova outbursts.

The structure of this review is as follows: Sect. 3, 4 and 5 discuss the properties of the Galactic Nova populations, their photometric and spectroscopic behaviours, their locations inside the Milky Way and the Maximum Magnitude vs. Rate of decline Relationship (MMRD). Sect. 6 makes a summary of the main nova surveys in the Local Group galaxies and beyond, up to Virgo and Fornax clusters and discusses their frequency of occurrence. Sect. 7 is dedicated to estimate the nova rate in the Globular Cluster systems. Sect. 8 compares the MMRDs in different Hubble type galaxies and discusses the outliers of the MMRD. In Sect. 9 we calibrate the LMC, M31 and Virgo MMRD relations with GAIA Data Release 2 (DR2 - Gaia Collaboration *et al.*, 2018) data to determine the distances toward the respective hosts. Sect. 10 dwells on the contribute of Novae to the chemical evolution of the host galaxy. Sect. 11 discusses the role of CNe as possible progenitors of SNe-Ia. In Sect. 12 we briefly report on the observations of novae at different wavelengths (radio, X- and γ -rays) and, finally, in Sect. 13 we give our conclusions.

3 Nova Populations in the Milky Way

Baade (1944, 1957) introduced the concept into astronomy that different kinds of stellar populations have different spatial distribution within the galaxies, therefore the starting point to search for a link between speed classes and properties of the nova progenitors is to establish the spatial distribution of Novae inside the respective parent galaxies. Due to their intrinsic luminosity, novae can be easily identified in the Milky Way and in external galaxies, thus novae are ideal tracers of their parent stellar population. However, in spite of these favorable conditions, the nova parent population is still an open question. Historical data on galactic novae (Hubble, 1929; McLaughlin, 1942, 1945, 1946; Payne-Gaposchkin, 1957) have received discrepant interpretations. According to Hubble:

"In general, the distribution of novae follows the distribution of luminosity in the nebula. The concentration in the nuclear region is conspicuous."

Modern data confirm the strong concentration of novae in the direction of the galactic bulge (Fig. 6). Kukarkin (1949); Kopylov (1955) and Plaut (1965) pointed out the existence of a thickening of novae towards the galactic plane and the galactic center, and the latter author classified novae as belonging to the 'disk population'. Minkowski (1948) and Payne-Gaposchkin (1957) showed that the galactic longitudes of novae and planetary nebulae (PNe) have similar distributions and therefore novae, like PNe, should belong to Pop II stellar population. Baade (1958) assigned novae to Pop II stellar population because of the occurrence of a few ones, like T Sco 1860, in very old stellar population systems of the Globular Clusters M80. Iwanowska and Burnicki (1961) suggested that novae are a mixture of Pop I and Pop II objects. Patterson (1984) pointed out that novae belong to an 'old disk' population. Observations of novae in extra-galactic systems have helped to clarify the picture: Della Valle and Duerbeck (1993) compared the cumulative distributions of the rates of decline for M31, LMC and Milky Way nova populations and found that galactic and M31 distributions are indistinguishable, whereas M31 and




Fig. 6 Distribution of the galactic longitudes of Novae inside the Milky Way. Data from Özdönmez *et al.* (2018)




Fig. 7 Distributions of the rates of decline for Milky Way, M31 and LMC. MW data (Özdönmez *et al.*, 2018), M31 data (Della Valle and Livio, 1995), LMC data (Della Valle and Livio, 1995; Shafter, 2013).

LMC distributions do not come from the same population at $\gtrsim 99\%$ significance level, see Fig. 7.

Since the speed class of a nova is related to some of the most important properties characterizing a nova system, like the mass of the WD and the amount of the ejected matter (see Sec. 3.2), the lack of systematic differences in the distributions of the rates of decline between Milky Way and M31 novae would suggest that they come from similar parent stellar populations, which are different from the one in which LMC novae are produced. On the other hand LMC is a bulge-less galaxy and




Fig. 8 Lightcurve of V382 Vel (Nova Velorum 1999) characterized by a smooth decline. Data obtained from the AAVSO Data Archive.

therefore novae in the LMC originate from disk population by definition. Therefore it seems logical to mainly associate novae in M31 and in the Milky Way to a bulge stellar populations. This conclusion is strengthened by spectroscopic observations of "disk" novae in the M33 and LMC (Shafter *et al.*, 2012; Shafter, 2013), which show remarkable differences compared with "bulge" M31 novae (Shafter *et al.*, 2011) (see also Sec. 6).

3.1 Nova lightcurves: the speed classes

The systematic studies of novae aimed at measuring their distances and luminosity at maximum, date back to the 20's. Lundmark (1922) derived an average absolute magnitude at maximum of $M_V = -6.2$ mag, but characterized by a broad dispersion, typified by the cases of Nova Lac 1910 and T Sco 1860, with $M_V = -1.1$ mag and $M_V = -9.1$ mag, respectively. In a following paper, Lundmark (1923) revised the previous value by averaging the distances obtained for about two dozen of novae with four different methods and he obtained $\langle M_V \rangle = -7.2$ mag, not far from $\langle M_V \rangle = -7.5$ mag found in modern studies (SG19). Novae normally show a fast rise to maximum light of the order of few hours, but in some case may last up to several days (e.g. RR Pictoris). The timescale and the morphology of lightcurves during the decline exhibit a very high degree of individuality (see Figs. 8, 9 and 10). In recent years, the possibility of using multi-wavelength observations led to important developments in understanding the evolution of nova outbursts. Novae lightcurves observed at different wavelengths (optical, UV and X-ray) exhibit different phenomenology. The ionisation of the ejecta generally increases with time and the peak of the emission shifts from optical to UV and finally to X-rays. A clear example is v1974 Cyg in Fig. 11, which reports the evolution of the emission as observed in different wavelengths (Cassatella *et al.*, 2004).




Fig. 9 Lightcurve of RR Pic (Nova Pictoris 1925) characterized by a slow rising to maximum light and slow decline exhibiting secondary maxima. Data obtained from the AAVSO Data Archive.




Fig. 10 Lightcurve of DQ Her (Nova Herculis 1934) characterized by a slow decline with oscillations, followed by a deep trough due to dust formation. Data obtained from the AAVSO Data Archive.

In the attempt to understand the link between timescale and morphology, several authors have tried to classify the nova light curve on the basis of either the rates of decline, expressed in magnitude per day (McLaughlin, 1939, 1942; Payne-Gaposchkin, 1957) or by grouping Novae in several families (Duerbeck, 1981) according to common behaviors exhibited by lightcurves after maximum light. For example, class A of Duerbeck (1981) was defined as "smooth fast decline", class C was defined as "extended maximum, deep minimum" and class E, defined as




Fig. 11 Lightcurve of V1974 Cyg (Nova cygni 1992) as observed in the optical (upper panel), in the 1455 Å UV continuum emission (second panel from the top) and in several UV emission line fluxes (third to seventh panels from the top). The last panel shows the evolution in X-rays as observed from ROSAT (Cassatella *et al.*, 2004).

speed class	t_2	mag/day
very fast	10^d	> 0.2
fast	$11^d - 25^d$	0.18 - 0.08
moderately fast	$26^d - 80^d$	0.07 - 0.025
Slow	$81^d - 150^d$	0.024 - 0.013
very slow	$151^d - 250^d$	0.013 - 0.008

Table 2 The nova speed classification as reported in Payne-Gaposchkin (1957).

”extremely slow nova with irregular lightcurve”. The classification in speed class, reported here (see Tab. 2) is the one adopted by Payne-Gaposchkin (1957).

The rate of decline are expressed in terms of mag/day or t_2 and t_3 , i.e. the times that a nova takes to decay by 2 or 3 magnitudes from maximum light. Multiple-peaks are often observed in slow novae, thus, this ambiguity might hamper the correct use of the MMRD for this nova subclass. This situation is well represented by the light curves of many galactic novae (Duerbeck, 1981) such as NQ Vul 1976 or RR Pictoris 1925, for example. If we compute the rate of decline after the first maximum we would obtain t_3 of a few days rather than 65 and 130 days as actually measured after averaging and smoothing the secondary maxima.

This classification is very popular and still adopted (see e.g. Warner, 2003) and it is certainly useful for quickly providing the observers with some important pieces of information necessary to trigger and plan the photometric and spectroscopic follow-up of an outbursting nova. However, in the last years it emerged the need to move from a simple grouping of novae, according to their speedy class or morphology of the lightcurves, toward a physical classification of nova outbursts. The idea that we develop in this paper is to relate the morphological classification of Payne-Gaposchkin with: i) the ”disk” and ”bulge” Galaxy environments by measuring the galactic latitude and longitude of nova systems and their height above the galactic plane (Della Valle *et al.*, 1992; Della Valle and Livio, 1998); ii) the spectral classification of novae in Fe II and He/N classes (Williams *et al.*, 1991; Williams, 1992). Recently Darnley *et al.* (2012) have proposed a CN classification in three classes, based on the evolutionary stage of the respective secondary stars: main sequence Nova, sub-giant Nova, and Red Giant Nova.

3.2 Disk and Bulge Novae

The quantitative characterization of the concept of nova populations into two classes of objects, i.e. fast and bright ’*disk novae*’ and slow and faint ’*bulge novae*’, has been independently elaborated by Duerbeck (1990), and by Della Valle *et al.* (1992), Della Valle and Livio (1994), Della Valle and Livio (1998). The former demonstrated that nova counts in the Milky Way do not follow a unique distribution (see Fig. 12), the latter authors showed that the rates of decline, which traces at first order some of the parameters driving the strength of the outburst, correlates with the spatial distributions of novae inside the Milky Way. Theoretical calculations suggest that the strength of a nova outburst is a strong function of the mass of the underlying WD (Shara, 1981; Livio, 1992): the brighter is the nova at maximum, the larger is the amplitude of the outburst (see Fig. 13) the more massive is the underlying WD.




Fig. 12 Adapted and updated from Fig. 2 of Duerbeck (1990). Galactic nova counts vs. apparent magnitude at maximum. The distribution shows that up to mag ~ 2 we mostly observe bright "disk" novae. As we proceed toward fainter magnitudes the number of bright novae is vanishing and the distribution starts to increase due to the observations of faint bulge novae.




Fig. 13 Distributions of the outburst amplitude plotted against the rate of decline (expressed as $v_d = 3/t_3$). Data from S19 and SS19, see also Table 9.

Since the luminosity at maximum of a nova correlates with the rate of decline, the distribution of the rates of decline traces the distribution of the masses of the WDs in nova systems. On the other hand, there exists a strong relation between the final mass of the WD and the initial mass of the progenitor star on the main sequence (MS): the higher the initial mass, the higher will be the mass of the WD. For example, Cummings *et al.* (2018) found - admittedly with some degree




Fig. 14 Relation between the height z above the galactic plane and the rate of decline obtained from novae which distances could be determined with GAIA DR2. The gray dashed lines mark the 3σ error strips. Data from S18 and SG19, see also Table 9.

of uncertainty - that a star of $7.5 M_{\odot}$ produces a WD of $\sim 1.3 M_{\odot}$, while stars of initial mass $\sim 1 M_{\odot}$ produces a WD of $\sim 0.6 M_{\odot}$. Galactic novae then should be characterised by different scale heights above the galactic plane according to the initial mass of the progenitor star. Novae originating from the more massive WDs should be located in the thin disk of the Galaxy, while novae from less massive WDs should occur in the thick disk, in the bulge or in the Galactic halo. Therefore we should observe a trend between the rates of decline and their heights above the galactic plane. To test this hypothesis we use the GAIA DR2 distances reported in the golden sample of S18 and SG19, to derive the height above the galactic plane:

$$z = 10^{0.2 \times (m - M + 5 - A)} \times \sin b \quad (5)$$

where b is the galactic latitude of each nova. The S18 and SG19 samples include 15 novae that are in common to both samples. To test our prediction we have merged the two data sets into a "bona fide" sample (Tab. 9) complemented by the recent GAIA DR2 measurement of the distance to V1500 Cyg (Nova Cyg 1975)². Fig. 14 shows strong evidence for the existence of a trend between the rate of decline and the height above the galactic plane. The correlation:

$$\log z = -0.797(\pm 0.199) \times \log v_d + 2.692(\pm 0.206) \quad (6)$$

is characterized by a confidence level of $\gtrsim 3\sigma$, according to the Student's t -distribution.

The trend illustrated in Fig. 14 indicates that the fastest nova systems, which contain the most massive WDs, are concentrated close to the galactic plane. It is

² We derived the GAIA DR2 distance of V1500 Cyg using the parallax provided in the GAIA DR2 database and using the formula given in Bailer-Jones *et al.* (2018), see also Muraveva *et al.* (2018). The GAIA DR2 archive is available at <https://gea.esac.esa.int/archive/>




Fig. 15 The distribution of the absorptions A_V (mag) as given in Downes and Duerbeck (2000), S18 and SG19 (see also Table 10)

difficult to understand how such a distribution could be due to selection effects, since there is no obvious mechanism to prevent the discovery of fast and bright novae at high z . Obviously, we cannot exclude that some heavily absorbed slow and faint nova at small z can have been overlooked.

However we note that the observed distribution (Fig. 15) of nova absorptions as estimated by different authors peaks at $A_V \sim 1$ mag and extends up to ~ 3 mag. Therefore the faint tail of the luminosity function of novae at maximum light, characterized by an absolute magnitude at maximum of ~ -6 mag, is bright enough to be detected, with modern surveys, at several kpc distance toward the galactic center even under more severe absorption constraints.

We note that the attached error bars to each data point, due to the exquisite accuracy of the GAIA DR2 distances, are very small. However, the scatter of the relation is intrinsically large. This fact implies that other parameters in addition to M_{WD} , such as the \dot{M} (Prialnik *et al.*, 1982; Townsley and Bildsten, 2004; Kato *et al.*, 2014), the envelope mass at ignition (Hachisu and Kato, 2010), the temperature of the WD (Prialnik and Kovetz, 1995; Yaron *et al.*, 2005) or the chemical composition of the accreted matter (Kovetz and Prialnik, 1985; Starrfield, 1999) affect the luminosity at maximum, the rate of decline, the ejection velocity and the mass of the expelled layers.

In the following we label 'disk' novae the objects characterized by $z \lesssim 150$ pc. They are mostly 'fast' declining objects characterised by a bright peak at maximum $M_V \sim -9$ mag, $t_2 \lesssim 12^d$ and $t_3 \lesssim 20^d$ and mainly belongs to the He/N class. Slow novae occur at higher z , up to 1 kpc, and are characterized by $M_V \sim -7.5$ mag at maximum and t_2 and $t_3 \gtrsim 12^d$ and 20^d , respectively. Hereafter we label these high z slow declining objects 'bulge' novae (Della Valle and Livio, 1998).




Fig. 16 v841 Oph lightcurve. Inset: an example of DN-like outburst occurred during the post-outburst stage.

3.3 Post-Novae

We can estimate as $\Omega \sim 1.5 \times GM^2/R \sim 10^{51}$ erg the binding energy of a $1.4 M_{\odot}$ WD. On the other hand, the energy budget associated with a nova outburst is $\sim 10^{45}$ erg. This simple order of magnitude comparison provides a clear indication that the nova outburst is not a catastrophic event for the WD, unlike from SN-Ia explosions. As a consequence, some decades after the outbursts the systems go back to quiescence and the post-nova magnitudes become very similar to those observed during the pre-nova stage (Robinson, 1975). The return to the minimum light is not as quick as the luminosity increasing to maximum. This fact can be quantified by studying the lightcurves of the oldest galactic novae for which a significant record of observations since the maximum epoch do exist: Nova Oph 1848 (v841 Oph), Nova Cyg 1872 (Q Cyg) and Nova Cir 1906 (AR Cir). The light curves reported in Figs. 16, 17, 18 have been obtained from the Steavenson's observations published between 1920 and 1950. An inspection of the data shows that v841 Oph and Q Cyg have returned to the quiescence state, at $V \sim 12.5$ mag and $V \sim 15$ mag in about 80 and 50 years, respectively.

Post-nova stages observations are not very frequent. When they are available, they are characterized by variation of the magnitude at minimum as large as $\sim 1-2$ mag. These brightness variation are due to multiple factors such as: brightness modulations possibly related to orbital period (Woudt *et al.*, 2004), solar-type cycles activity of the secondary (Bianchini, 1987; Warner, 2003) on time scale of the orders of months/years (see inset in Fig. 17) but also to dwarf nova activity on time scales of 50-270 days (Della Valle and Rosino, 1987; Della Valle and Calvani, 1990).




Fig. 17 Q Cyg lightcurve. Inset: an example of modulation of the post-outburst lightcurve on time scale of hundreds days.




Fig. 18 AR Cir lightcurve. Inset: example of DN activity during the pre-outburst stage.

The distribution of the absolute magnitude at minimum, see fig. 19, is a distribution characterised by a mean brightness of $M_V = 3.8 \pm 0.24$ mag. Such distribution of luminosities requires a mass transfer rate between $\sim 10^{-8}$ and $10^{-9} M_\odot/\text{yr}$ (see fig. 5 in SG19). The brightness at minimum of the fastest declining novae implies mass transfer rates that are a full order of magnitude below the average value for most novae, close to the DNe values (Vogt, 1990).




Fig. 19 The distribution of absolute V magnitudes of GAIA DR2 novae at quiescence.

The link between DNe and CNe has been hypothesized long ago in the framework of the so called "Hibernation" scenario (Shara *et al.*, 1986) (see also Hillman *et al.*, 2020) which was originally elaborated to explain an apparent two orders of magnitude discrepancy between the density of classical novae inferred in the solar neighborhoods and those deduced from nova surveys in M31, nova theory and U, B photometric surveys. Intrinsically low CN space density would imply very short recurrence times between the outbursts, which in turn would imply high values of the \dot{M} , which are not observed. According to these authors the space density of CNe is much higher but novae would spend most time between two consecutive outbursts in a "hibernation" state. This stage would be characterised by low or very low mass transfer rates (10^{-10} - $10^{-12} M_{\odot}/yr$) as result of the increasing separation between the companion stars after each outburst, as consequence of the conservation of the angular momentum of the binary system. The canonical mass transfer rate of 10^{-8} - $10^{-9} M_{\odot}/yr$ would be recovered as soon as the magnetic braking and/or gravitational radiation emission decrease the separation between the companion stars. During this intermediate stage in which \dot{M} increases from hibernation to the canonical nova values, the system would exhibit DN activity. This scenario has received in the years some observational support.

First of all, the dwarf nova activity was detected not only after the nova outbursts, but it was also observed decades before the outbursts in V446 Her or IV Cep (Della Valle and Calvani, 1990) and AR Cir (see inset in Fig. 18). Duerbeck (1992) could measure an average decline of 10 ± 3 millimag per year in the post outbursts decline of nine old novae, admittedly with large individual scatter. However these data allowed Duerbeck to conclude that the observed trend was "compatible with the prediction of the hibernation hypothesis". Finally in more recent years, Shara *et al.* (2007, 2012) discovered Nova shells around two dwarf Novae: Z Cam and AT Cnc.

On the other hand the "Hibernation" scenario was introduced on the basis of a supposed existence of a two orders of magnitude discrepancy between the density

Density (pc^{-3})	Authors
10^{-5}	Patterson (1998)
3×10^{-6}	Warner (2001)
3×10^{-5}	Schwope <i>et al.</i> (2002)
$< 5 \times 10^{-7}$	Cieslinski <i>et al.</i> (2003)
$0.3 - 3.3 \times 10^{-5}$	Pretorius and Knigge (2007)
$< 2 \times 10^{-5}$	Hernandez <i>et al.</i> (2018)
$< 5 \times 10^{-6}$	Schwope (2018)

Table 3 Collection of measurements of CVs density close to the Sun.

of CVs occurring close to the Sun and the rate of CVs that was believed to be as high as $\rho \sim 10^{-4} \text{pc}^{-3}$ (Shara *et al.*, 1990). This density value of CVs has been the subject of many revisions in the last two decades that we have summarized in Tab. 3.

These values are 1-3 orders of magnitude smaller than the $\rho \sim 10^{-4} \text{pc}^3$, which originated the quest for the existence of a hibernation stage for CVs. Della Valle and Duerbeck (1993) found a density of nova outbursts in the galactic disk of:

$$\rho = 0.30 \pm 0.15 \times 10^{-10} \text{pc}^{-3} \text{yr}^{-1} \quad (7)$$

consistent with previous estimates of $\rho = 0.5 \times 10^{-10} \text{pc}^{-3} \text{yr}^{-1}$ (Duerbeck, 1984; Patterson, 1984), $\rho = 0.2 - 0.9 \times 10^{-10} \text{pc}^{-3} \text{yr}^{-1}$ (Naylor *et al.*, 1992) once we have removed from their original values the contribute of bulge novae. The next step is to derive the density of CNe from the density of nova outbursts by multiplying the observed density of nova outbursts for the recurrence time between the outbursts. Disk novae that are mostly fast and bright originate from massive WDs, say $1.1-1.3 M_{\odot}$. According to Sect. 1.2, these novae have recurrence times of the order of 2×10^4 years, which gives a density of Classical Nova progenitors of:

$$\rho_{CNe} = 0.6 \times 10^{-6} \text{pc}^{-3}. \quad (8)$$

This result is consistent with the lower density values reported in Tab. 3, which were derived with a broad variety of approaches including OGLE survey (Cieslinski *et al.*, 2003), ROSAT observations (Schwope *et al.*, 2002; Pretorius and Knigge, 2007; Schwope, 2018) and SDSS survey (Hernandez *et al.*, 2018). The still large range of uncertainties in the knowledge of the CV density does not exclude that there is some room for the existence of a "hibernation" stage in a "mild" version i.e. it should be active over a shorter interval of time than originally hypothesized.

3.4 The interaction of Nova with the Interstellar medium

The interaction of the expanding shells with the interstellar medium was discussed by Oort (1926) and Ambartsumian (1952). These models were developed on simple assumptions: the conservation of energy, average expansion velocities of the nova shell and ejecta masses of $\sim 10^{-5} M_{\odot}$. After assuming a range of interstellar densities, Oort could estimate: i) the "half-lifetimes" that is the time after which the expansion velocities of the ejecta have decreased to 50% of its initial value, to be 50-100 years and ii) 10,000 years the time needed to velocities of nova ejecta to

become indistinguishable from the chaotic motions of the circum-burst interstellar medium. Attempts to measure the deceleration of the nova ejecta (Duerbeck, 1987) were hampered by the inhomogeneities affecting the comparison between images of post-nova nebulae obtained decades apart. Also one should consider the possibility that the circumburst medium could have been swept away by previous outburst episodes. Downes and Duerbeck (2000) in their survey on nova shells, which included also the Slavin *et al.* (1995) data, decided to derive the distances via nebular parallaxes toward each nova by assuming no deceleration for the ejecta.

3.5 The Galactic Nova rate

The knowledge of the galactic nova rate is important for at least two reasons: i) some nova-like systems are believed to be progenitors of SNe-Ia (see sect. 11) therefore the correct measurement of the nova rate is an essential ingredient to be compared with the relatively well known SN-Ia rate in the Milky Way (Cappellaro *et al.*, 1999, 2015); ii) the role of novae as contributors to the Galactic nucleosynthesis has been well established on theoretical grounds (D’Antona and Matteucci, 1991; Romano *et al.*, 2001; Travaglio *et al.*, 2001; Romano and Matteucci, 2003; Prantzos, 2012). Recently observations of ^7Be (Tajitsu *et al.*, 2015) and ^7Li (Izzo *et al.*, 2015) have provided significant empirical support to these theoretical predictions. However, despite of the importance of these issues the correct value of the galactic nova rate is still poorly known (see Table 4). Just to give an idea here below we report a short summary of the estimates available in literature. Lundmark (1935) derived ~ 50 novae/yr. Allen (1954) inferred from his paper on the whole-sky statistics of celestial object about $R = 100 \div 200$ novae/yr, which is similar to the estimate obtained later on by Sharov (1972) of $R = 260$ novae/yr and at variance with $R \sim 50$ novae/yr estimated by Kopylov (1955). In more recent epochs Liller and Mayer (1987) have corrected the "observed" rate of 3.7 novae/yr for incompleteness and selection effects and they derived 74 novae/yr ± 24 . Hatano *et al.* (1997) have revised Liller and Mayer (1987) and assumed that most novae are produced in the disk and found $R = 41 \pm 20$. van den Bergh (1991) derived about $R = 16$ novae/yr by scaling the nova rate of M31 to the GCs population ratio Galaxy/M31=0.56. Ciardullo *et al.* (1990); Della Valle and Livio (1994); Darnley *et al.* (2006) found $R = 11 \div 46$ novae/yr and $R = 15 \div 24$ and $R = 34^{+15}_{-12}$ respectively after scaling the nova rates measured in extragalactic systems to the luminosity of the Milky Way. Shafter (1997) found $R = 35 \pm 11$ by studying the distribution of novae inside the Milky Way and applying a number of correction factors for incompleteness and observational bias. More recently, the same author (Shafter, 2017) has increased his previous estimate to $R = 50^{+31}_{-23}$ by revising the correction factors to the galactic nova counts and by assuming uniform properties between bulge and disk novae.

Fig. 20 shows the relationship between the nova rate and the K luminosity of the parent galaxy. A least squares fit to the data points gives:

$$\log N \approx 0.779 \times \log L_K - 7.09 \quad (9)$$

After assuming for the Milky Way $L_B \approx 2 \times 10^{10} B_\odot$ (van den Bergh, 1988), and (B-K)=3.50, corresponding to Sb Hubble type, we find ≈ 22 novae/yr, admittedly with a large attached error. However this result is consistent with the recent

Nova/yr	author
50	Lundmark (1935)
100- 200	Allen (1954)
50	Kopylov (1955)
260	Sharov (1972)
74	Liller and Mayer (1987)
11-46	Ciardullo <i>et al.</i> (1990)
16	van den Bergh (1991)
20 ± 5	Della Valle and Livio (1994))
41 ± 20	Hatano <i>et al.</i> (1997)
35 ± 11	Shafter (1997)
34^{+15}_{-12}	Darnley <i>et al.</i> (2006)
50^{+31}_{-23}	Shafter (2017)

Table 4 One century of Galactic Nova rate measurements.




Fig. 20 Extragalactic nova rates plotted against the K -band luminosity (data from Table 7 in sect. 6.6).

”direct” measurement of Mróz *et al.* (2015). These authors have measured the bulge nova rate of the Milky Way with the data of the OGLE survey. They find a rate for the Galactic bulge of 13.8 ± 2.6 novae/yr that according to these authors: *”...is much more accurate than any previous measurement of this kind, thanks to many years monitoring of the bulge by OGLE survey”*. A ”rounded” rate of about 14 ± 3 novae per year in a galaxy like the Milky Way, where the nova production is bulge dominated, implies a global rate that hardly can exceed 30 novae/yr in good agreement with the frequencies found in the 90’s (Ciardullo *et al.*, 1990; Della Valle and Livio, 1994; Shafter, 1997). Only if incompleteness affects dramatically the galactic nova counts in the disk we can relax the assumption that novae in the Milky Way are mainly produced in the bulge and therefore values of the galactic nova rate as large as ~ 50 novae/yr are still possible (Shafter, 2017). Presently the existence of this bias is not supported by the observations.




Fig. 21 The average lightcurve morphology of a Nova (from Warner (2003)). We have added in different colors the description of the main spectroscopic phases.

4 The Spectral evolution of Novae

Many contributions have been devoted to describe and to summarize the spectroscopic evolution of nova outbursts, mostly dedicated to identify the different nova stages, their absorption and emission lines and measuring the expansion velocities of the ejecta Payne-Gaposchkin (1957) and McLaughlin (1960). More recently, Shore (2012) has provided a general physical description of this sequence of stages that helped a lot to enlighten the physics underlying the spectroscopic evolution of Novae. A similar description based on the color-color evolution of nova outbursts has been also provided by Hachisu and Kato (2014).

In the following we have linked the "classical" photometric stages with the "old" and "new" spectroscopic descriptions of nova phases (see fig. 21). The spectral evolution of Novae is basically driven by two physical parameters: the mass density of the ejecta and their expansion velocities. In spite of the broad phenomenology observed in nova spectra six spectral stages have been defined and in the following we provide an essential description for each of them.

Initial rise - Pre-maximum spectrum: it corresponds to the adiabatic free expansion of the ejecta that is rapidly cooling since the expansion regime is much shorter than the characteristic cooling time of the ejecta. This phase is also referred as "hot fireball" spectrum and it is observed very seldom because it implies: i) the discovery of a nova during its rise at maximum, that normally lasts from a few hours to few days and ii) the immediate availability of telescope time, two conditions that normally do not occur frequently. Some novae show a pre-halt maximum (the so-called shoulder phase) in the light curve, which is characterized by bluer colours and that anticipates the final rise to the maximum brightness. This feature in the light-curve has been caught in very few novae, such as T Pyx (Shore




Fig. 22 Initial rise/pre-maximum spectrum of T Pyx (adapted from Izzo *et al.*, 2012).

et al., 2011) and V1974 Cyg (Shore *et al.*, 1993). The spectra bear some similarity to F supergiants (Kodaira, 1970). Balmer and non-Balmer lines are characterized by high expansion velocities (2,000-3,000 km/s) as measured from the minimum of the respective P-Cygni profiles. These features are produced during the initial expansion of the optically thick envelope ejected by the nova. (see Fig. 22).

Maximum light - Principal spectrum: this stage corresponds to the rise to the maximum brightness in V and R magnitudes and a contemporaneous decrease in luminosity at blue and UV wavelengths. This is the time when the fireball reaches the critical cooling temperature, the pseudo-photosphere starts to recede and the spectrum is being dominated by the free-free emission from thinner plasma outside the photosphere (Hachisu and Kato, 2006). Consequently, the nova ejecta itself acts now as a passive media, by converting the UV flux into longer wavelengths emission and then showing low-ionization transitions in the spectra, such as the Fe II group species (hence the name "*iron-curtain*"), Na I D and Ca II. These lines are flanked by broad P-Cyg profiles with blue-shifted velocities generally lower than those observed during the fireball phase (see Fig. 23). At the same time, the optically-thick stage in UV bands leads also to the presence of "pumped" transitions, like the O I 8446 Å, whose upper level is over-populated by the H Ly- β 1025.72 Å transition, which is almost coincident with the transition O I 1025.76 Å (Bowen, 1947; Williams, 2012). High-resolution spectroscopic observations have also revealed the presence of narrow and low-ionization heavy element absorptions (THEA - Williams *et al.* (2008), see Sec. 4.2), which are detected during the brightest epochs of several nova outbursts with expanding velocities relatively low (~ 500 km/s or less).

During the Principal phase, the so called **Diffuse Enhanced** absorption systems appear during the very early decline. The only feature that differentiates the Diffuse Enhanced from the Principal spectrum is the presence of multiple P-Cyg profiles (Fig. 23). All these absorption components show an apparent accelera-




Fig. 23 Principal spectrum phase of V1369 Cen (Izzo *et al.*, 2015), showing the presence of diffuse enhanced absorptions in the P-Cygni of the brightest emission lines.

tion of the ejecta with time, as inferred from the P-cygni absorptions. However, this acceleration is not real, it rather originates in the downturn density of the ejecta and the contemporaneous recession of the hot pseudo-photosphere. Indeed, a decreasing density implies a more optically thin ejecta, which implies a fragmentation of the broad P-Cygni absorption lines into several weaker structures. Very recently, it has been proposed that the appearance of new and faster absorption systems in the P-Cygni profiles of Balmer lines correlates with the presence of rebrightenings, or flares, in the light curves (Li *et al.*, 2017; Aydi *et al.*, 2019). These new systems would be the result of fast wind ejection from the underlying WD and their interaction with the pre-existing ejecta will lead to strong shocks, originating the high-energy emission observed in some novae (Metzger *et al.*, 2015; Cheung *et al.*, 2016; Mukai and Sokoloski, 2019; Aydi *et al.*, 2020), see also Sec. 12. Another explanation for the presence of the rebrightenings is that nova ejecta is actually composed by strong density inhomogeneities created during the TNR (Williams, 2016).

Toward the end of the early decline and at the beginning of the transition stage (that is the transition from optically thick to optically thin condition) another spectral system appears, the so called **Orion spectrum**. The optical spectra now show features due to high-ionization lines (He I, O II, N II), and the CIII/NIII lines Bowen blend ("4640 band") see Fig. 24. Their appearance in the spectrum is due the low density of the ejecta combined with the intense UV flux. An important characteristics of this stage is that the expansion velocities measured from these transitions do not generally coincide with the ones inferred during the iron-curtain phase at optical wavelengths. As the density further decreases the spectrum develops nebular lines which increase their brightness with time (Osterbrock & Ferland, 2006). During this stage the absorption systems disappear.




Fig. 24 Orion phase spectrum observed in ASASSN-16kt (Izzo *et al.*, 2018).

The enhancement in flux of the He II marks the end of the optically thick phase e.g. (Munari *et al.*, 2014) and the beginning of the **Nebular phase** followed by the increase in luminosity of forbidden transitions ((Rafanelli & Rosino, 1978)) of oxygen auroral and nebular [O III] 4363 and 4959, 5007, respectively and nitrogen [N II] 5755, 6548, 6584 at optical wavelengths, see Fig. 25. In the spectra of ONeMg novae we also observe bright [Ne III] 3869, 3968 and [Ne v] 3346, 3426. It is well known (McLaughlin, 1943; Vainu Bappu and Menzel, 1954; Payne-Gaposchkin, 1957) that spectroscopic observations obtained during the Nebular stage, when the ejecta are optically thin, are useful to study the geometrical and ionization structure of the ejecta (e.g. Shore *et al.*, 1996; Shore, 2013; Vanlandingham *et al.*, 2001; Iijima, 2012; De Gennaro Aquino *et al.*, 2014, 2015; Mason *et al.*, 2018). Nebular lines show flat-top or saddle shape or split profiles, often characterized by fine structures on their top, then suggesting both the presence of density variations in the ejecta and in the geometric distributions as polar caps or equatorial rings. During the nebular phase, the peak of the emission shifts to UV wavelengths, and finally to X-rays. The UV wavelength range was studied in very details thanks to the International Ultraviolet Explorer (IUE) satellite during the '80s and the first half of the '90s (see e.g. SG19). In this phase emission lines of ionised helium and high ionisation of carbon, nitrogen, oxygen and neon freshly-produced elements are observed (see e.g. Cassatella *et al.*, 1979; Stickland *et al.*, 1981; Shore *et al.*, 1996; Cassatella *et al.*, 2002, 2004, and references therein). Optical observations combined with UV spectra obtained at these epochs, allowed the determination of the ejecta abundances (Vanlandingham *et al.*, 1996, 1997; Schwarz, 2002; Shore *et al.*, 2003; Schwarz *et al.*, 2007). These results have definitely proven that element abundances in novae are generally super-solar, up to 1-2 orders of magnitude ((Livio and Truran, 1994; Gehrz *et al.*, 1998, 2008), see also José, Shore and Gehrz in Bode and Evans (2008)). During the nebular stage, Novae can also achieve their highest degree of excitation, when the ejecta are completely optically thin and the underlying hot WD surface is revealed as a bright source in soft X-rays (Oegelman




Fig. 25 Nebular spectrum observed in V1369 Cen (Izzo *et al.*, 2015; Mason *et al.*, 2018).

et al., 1993) due to the burning of residual hydrogen-rich material near its surface (see also sect. 12). In these conditions soft X-rays and the strong UV photon flux are capable to photo-ionize heavy elements, but the low density of the nova environment at these epochs will avoid their complete recombination, resulting in the presence of highly-excited coronal lines in the spectra such as [Ni VIII] 4446, 4493; [Ni IX] 4332, 4404; [Fe VII] 6086; [Fe x] 6376; [Fe XIV] 5303; and [Ar X] 5535 (Rosino *et al.*, 1989).

Finally, few years after the outburst the object develops the **Post-nova spectrum** that is characterized by the gradual disappearance of coronal and then forbidden lines, due to the decrease of the ionization level (see Tappert *et al.* 2014). Residual emission lines remain strong or at least still detectable, e.g. $H\alpha + [N II]$, NIII 4640 and He II 4686 emission line originating from the accretion disc around the WD. The latter two lines, for example, were still visible in the spectra of FH Ser 10 years after the outburst (Rosino *et al.*, 1986).

4.1 The Tololo Spectroscopic Taxonomy

Williams (1992, 1994); Williams *et al.* (1994) after studying two dozen of galactic and Magellanic novae concluded that novae can be broadly divided into two spectroscopic classes: the Fe II and He/N novae. The former are characterized by slow spectroscopic evolution with expansion velocities $\lesssim 2500$ km/s (FWZI) and the presence of Fe II lines in the early emission spectrum as the strongest non-Balmer lines. The latter are fast spectroscopically evolving novae, characterized by high expansion velocities ejecta $\gtrsim 2500$ km/s (FWZI) with He and N lines being the strongest non Balmer lines in the emission spectrum near maximum. Hybrid objects (e.g. V1500 Cyg), which evolve from Fe II to He/N, are classified as Fe II-b

(b=broad) and are physically related to the He/N rather than to Fe II class (see Fig. 26).

In Fig. 27 we report the heights above the galactic plane of the novae of our bona fide sample (S18 + SG19) after being classified according the Williams' criteria. The plot shows that novae belonging to the He/N and Fe II-b classes tend to concentrate close to the Galactic plane with a typical scale height ~ 100 pc, whereas Fe II novae are distributed more homogeneously up to $z \sim 300$ pc and possibly beyond (Della Valle and Livio, 1998). Fig 28 shows an upgraded version of the relationship between the rate of decline and the average expansion velocity of the nova ejecta measured at early stages, first observed by McLaughlin (1940). Filled dots refer to the He/N novae and they are characterized by the largest expansion velocities. On the basis of these plots we can infer that objects previously classified as 'disk' and 'bulge/thick-disk' novae tend to correspond to the He/N (+ Fe II-b) and Fe II spectroscopic classes. To explain this behavior one should consider the following argument: the more massive the WD (for a given \dot{M} and T_{WD}) the smaller is the mass of the accreted envelope (in view of $\Delta M_{acc} \propto R_{WD}^4 / M_{WD}$) the more violent is the outburst and the larger the fraction $\Delta m_{shock} / \Delta m_{wind}$ where Δm_{shock} is the shell mass ejected at the maximum and Δm_{wind} is the fraction of the shell mass ejected in the subsequent continuous optically-thick wind phase. Since He/N spectra are formed in the shell ejected at maximum light it is very likely that He/N novae are generally associated with massive WDs, which eject the less massive envelopes (see Sect. 1.2). In the framework of this interpretation the distributions reported in Fig. 27 and Fig. 28 are simple consequences of the trend exhibited by the rate of decline vs. height above the galactic plane (see Fig. 14).

4.2 The THEAs: Transient Heavy Elements Absorptions

The previous spectroscopic classifications were significantly enriched by high resolution observations of novae ($R \sim 50,000$) carried out at ESO La Silla Observatory at the end of 90's (Della Valle *et al.*, 2002; Mason *et al.*, 2005; Ederoclite *et al.*, 2006) and at other facilities (Sadakane *et al.*, 2010). These observations have revealed (Williams *et al.*, 2008) the presence around maximum light of a new spectroscopic system formed by transient heavy element absorptions (THEAs). These authors were able to identify more than 300 absorption lines of single ionized elements with $E_{ion} < 6$ eV belonging to the iron group, such as: Sc, Ti, V, Cr, Fe, Sr, Y, Zr. The FWHMs of THEAs show a broad range of velocities between 30-300 km/s. They all progressively weaken and disappear over timescales of days or weeks.

The observations show that THEAs are superimposed on the principal absorption systems of Novae and this implies that THEA systems originate outside the expanding photosphere, so they are not a direct product of the nova outburst. After days or weeks when the nova ejecta reach the THEAs, the transient systems are not observed any longer. Another property of THEA lines is that the time at which these lines are not observed depends on the rate of decline of Novae. The three fastest novae in the sample of Williams *et al.* (2008) do not show THEA, indeed, while the slowest objects can exhibit THEAs up to 1-2 months past maximum. The excellent temporal coverage suggests as most plausible explanation




Fig. 26 Typical "Fe-II" nova spectra obtained close to maximum light (top panel). "He/N" spectra of novae observed a few days past maximum (bottom panel) (from Williams (1992).)




Fig. 27 Histogram of the heights of novae above the galactic plane according to the Tololo spectroscopic classification. Data obtained from Gaia DR2 distances published in S18 and SG19, while spectral type obtained from Della Valle and Livio (1998)




Fig. 28 Relationship between the rate of decline and average expansion velocity (obtained from the FWHM) at early stages. Filled dots refer to the He/N and Fe II-broad (hybrid novae) classes. Light dots to Fe II novae. The filled dot at small rate of decline is V5558 Sgr a Fe II-b nova. Data from S18, SG19, Della Valle & Livio (1998) and (Özdönmez *et al.*, 2018)




Fig. 29 A selected wavelength range showing the optical spectral evolution in V1369 Cen within the first three weeks from the nova outburst (from Izzo *et al.* 2015). The THEA narrow absorptions are clearly detected within the first two weeks, when the ejecta are still optically thick. The ejecta density decreases with time and then the faint and narrow absorptions are the first features to disappear.

that the faster ejecta, which are associated with the fastest declining novae, have reached the THEA absorptions very few days past maximum, before our observations could start. As an alternative one should figure out a mechanism for which the THEA are not developed around fast declining novae. The origin of this material can be ascribed to activity of the companion star. The chemical species observed in THEA systems and the contemporaneous lack of H would seem to exclude that the observed material is a residual of previous outbursts. In the paper by Williams *et al.* (2008), it was proposed that they originate from a continuous mass ejection from the secondary star and some empirical estimates suggest that the involved amount of gas is $\sim 10^{-5} M_{\odot}$ located at 10-100 AU from the binary system.

5 The Maximum Magnitude vs. Rate of Decline Relationship in the Milky Way (MMRD)

The existence for this relation was first suggested by Zwicky (1936), who named it as "life-luminosity" relation. It was expressed as: $M_{max} = -5 \times \log \tau_{\Delta m} + \text{const}$, where $\tau_{\Delta m}$ is the time necessary to decrease Δm magnitudes from maximum. Unfortunately he calibrated the relation by using two galactic novae, Nova Per 1901 and Nova Aql 1918, and three Supernovae, SN 1936A in NGC 4273, SN 1926A in NGC 4303 and SN 1885A in M31, so the results were misleading. The first calibration of the MMRD relation entirely based on nova events was carried out by McLaughlin (1945). He used 13 Galactic Novae whose distances were determined with the nebular parallaxes, the intensities of interstellar absorption lines or the




Fig. 30 The maximum magnitude vs. rate of decline relation calibrated by McLaughlin (1945). Double circles: nebular expansion; open circles: intensity of the interstellar Calcium; solid circles: T Coronae and T Scorpii; crosses: galactic rotation; solid triangles: novae in Andromeda; open triangles: novae in the LMC; squares: novae in Sagittarius.

residual velocities from interstellar lines interpreted as due to galactic rotation, combined with novae coming from the Hubble survey in M31 and three novae from LMC (Fig. 30).

In the following decades a number of relations based on different nova samples, calibrators and assumptions on the galactic absorption in the interval 0.8 mag/kpc (McLaughlin, 1945) up to 3.5 mag/kpc (Kopylov, 1952) have been provided by many authors:

- $M_o = 2.0 \times \log t_3 - 10.1$ (Vorontsov-Velyaminov, 1947);
- $M_o = 3.7 \times \log t_3 - 13.8$ (Kopylov, 1952);
- $M_{pg} = 2.5 \times \log t_3 - 11.8$ (Schmidt, 1957);
- $M_{pg} = 2.5 \times \log t_3 - 11.5$ (Schmidt, 1957);
- $M_V = 2.5 \times \log t_3 - 11.75$ (Schmidt, 1957);
- $M_B = 1.8 \times \log t_2 - 11.5$ (Pfau, 1976);
- $M_{pg} = 2.4 \times \log t_3 - 11.3$ (de Vaucouleurs, 1978).

In the 80's the calibration of the MMRD relation received its first breakthrough. Cohen and Rosenthal (1983) e Cohen (1985) were able to measure the size of 19 shells around post-novae. These observations, together with the i) expansion velocities measured from the spectra, ii) the assumption of spherical symmetry in the expansion of the ejecta and iii) individual estimates of the galactic absorption toward each nova, provided for the first time a set of nova distances, which were homogeneously measured, and a new calibration of the MMRD relationship. The result, reported in Fig. 31, yielded:




Fig. 31 The MMRD relation obtained via nebular parallaxes of 19 novae (Fig. 7 from Cohen 1985).

$$M_V = 2.41 \pm 0.23 \times \log t_2 - 10.70 \pm 0.30. \quad (10)$$

Downes and Duerbeck (2000) have increased this sample to 30 objects and improved the quality of "old" measurements of nova shells surrounding nine post-novae with ground-based and Hubble Space Telescope (HST) data. Their linear fits gives:

$$M_V = 2.54 \pm 0.35 \times \log t_3 - 11.99 \pm 0.56 \quad (11)$$

$$M_V = 2.55 \pm 0.32 \times \log t_2 - 11.32 \pm 0.44 \quad (12)$$

The GAIA DR2 distances (Gaia Collaboration *et al.*, 2018) give the opportunity to check the reliability of nova distances measure via nebular parallaxes. Fig. 32 shows that nebular parallaxes are certainly consistent with GAIA DR2 measurements below 1.5 kpc. The scattered points at distances larger than 1.5 kpc is likely due to failures on the assumptions of the nebular parallaxes methodology (e.g. spherical symmetry expansion, no deceleration of the ejecta) and in particular to the poor quality of the A_V estimates.




Fig. 32 Distances obtained via nebular parallaxes (data from Downes & Duerbeck 2000) vs. GAIA DR2 distances (data from S18 and SG19).

The existence of the MMRD can be broadly outlined from simple physical arguments (Bath and Shaviv, 1978; Kantharia, 2017) or theoretically modelled (Hachisu and Kato, 2015). At first order one can estimate the luminosity at maximum of a nova as: $L = 4\pi R^2 \sigma T^4$, where $R \sim v \times t$ with v being the average expansion velocity of the ejecta and t the time from the outburst. This simplified approach shows in very intuitive way why novae characterized by the largest expansion velocities can achieved the highest luminosity at maximum light. For this reason their envelopes become optically thin very soon and their decline in luminosity proceed faster than in intrinsically fainter (and slowly expanding) novae. The tacit assumption in this argument is that the masses of the ejecta have roughly the same mass for all CNe: and this is true within a factor about ten, between 10^{-4} and $10^{-5} M_{\odot}$. For example some RNe or *ultra-fast* novae (Yaron *et al.*, 2005; Shara *et al.*, 2017a) that expel envelopes, about 100-1000 times lighter than observed in CNe, achieve the transparency condition ($\tau \ll 1$) in the ejecta very soon, therefore their maxima are fainter and also they decay much more quickly than CNe. The MMRD relation was analytically derived by Shara (1981) and more recently computed from the theoretical modeling of different lightcurves calculated for a large variety of white dwarf chemical composition and masses (Hachisu and Kato, 2007; Hachisu *et al.*, 2008; Hachisu and Kato, 2019; ?; Kato *et al.*, 2009). After normalizing the lightcurves for an appropriate time scale factor, the latter authors derive a "universal" behavior that can be described as : $M_V = 2.5 \times \log t_3 - 11.6$, which fit the GAIA data within a sufficient degree of approximation (see Fig. 33).

However, even after the papers by Cohen (1985) and Downes and Duerbeck (2000) the large scatter exhibited by the MMRD, about 3 magnitudes (see Fig. 31) has hampered the use of novae as effective distance indicators. This scatter is essentially produced by three factors. The first one is connected to the nebular




Fig. 33 The absolute V magnitude plotted against the t_3 value for the SG19 and S18 nova sample. The solid line corresponds to the best-fit found by Hachisu and Kato (2009)

parallax methodology that assumes symmetric expansion and no deceleration for the nova ejecta. By neglecting the role of opening angles and inclination of the ejecta it can bring into the distance measurements significant errors (Ford and Ciardullo, 1988; Shore, 2012; Wade *et al.*, 2000) at least for more distant and old novae. The second element is the lack of a method to measure with the necessary accuracy the visual extinction toward the nova. We observe novae from inside our Galaxy and the estimates of the reddening toward each nova are very crude (see Sect. 5.2). They are based on empirical relations based on the intensity of interstellar lines (Munari and Zwitter, 1997) and from the comparison of predicted and observed emission lines ratios (Robbins, 1968a,b). For example, unblended H and He recombination lines can be used for this goal if they are optically thin and not affected by collisions (Della Valle *et al.*, 2002). Yet the attached errors are large and the final estimate of absorptions very uncertain.

5.1 The Shape of the MMRD Relationship

Local effects as the reddening toward each nova and the difficulty of determining distances accurately suggest to look at other galaxies to calibrate and to determine the "true" shape of the MMRD relationship. These drawbacks are well detectable by comparing Fig. 31 with Fig. 34.

The former plot is the MMRD derived through nebular parallaxes by Cohen (1985). Here the uncertainties on the distances and absorptions, inflate the intrinsic scatter of the MMRD, which can well fitted by a linear regression. Fig. 34 represents the MMRD derived by studying the nova population in M31 and LMC (data from Della Valle and Livio, 1995; Shafter, 2013, all data have been collected in Tab. 13 and Tab. 14). This plot shows a linear trend only between $0.04 < v_d < 0.2$ ($10 < t_2 < 50$ days). For the entire range of v_d ($=2/t_2$) the analytic representation is a reverse S-shaped function (Capaccioli et al. 1989), (Della




Fig. 34 M31 and LMC novae in the MMRD plane (data from Della Valle and Livio, 1998; Shafter, 2013). The best fit is the solid line described by eq. 15. With black and grey dashed lines we report the $\pm 3\sigma$ strips, with $1\sigma = 0.25$. A χ^2 test indicates that the M31 and LMC nova data samples are better fitted by a S-shape formulation than a linear function: $\chi^2_S = 1.70$ while for the linear model $\chi^2_{lin} = 1.87$.

Valle and Livio, 1995). A $\tilde{\chi}^2$ test indicates that M31 and LMC nova data are better fitted by a S-shape function (see eq. 15 in Sect.5.2) than a simple linear fit: $\tilde{\chi}^2_S = 1.70$ while for the linear model $\tilde{\chi}^2_{lin} = 1.87$. The linear fits performed on novae in the Milky Way are the result of scanty statistics combined with the large uncertainties associated to the measurements of the interstellar absorption along the line of sight to each single nova.

Theoretical attempts to explain the MMRD relation also provide evidence that a linear fit to the nova data only provides a rough first order approximation. Hartwick and Hutchings (1978); Shara (1981); Livio (1992) were able to derive simple relationships between the absolute magnitude at maximum and the mass of the underlying WD. For example Livio (1992) derives:

$$M_{max} \approx -8.3 - 10 \times \log M_{WD}/M_{\odot} \quad (13)$$

and in turn a theoretical MMRD relation of the form:

$$t_3 = 51.3 \times 10^{0.2 \times (M_B + 9.76)} \times [10^{0.067 \times (M_B + 9.76)} - 10^{-0.067 \times (M_B + 9.76)}]^{3/2} \quad (14)$$

that is expressed in days. The red dashed line in Fig. 35 shows the theoretical relation, superimposed on the GAIA MW data, which is expressed in absolute magnitudes. The flattening of the observed distribution at high luminosities is real and can be interpreted as a physical threshold due to the fact that the WDs in nova systems which result in super-Eddington novae are approaching the $\sim 1.4M_{\odot}$




Fig. 35 (Upper panel) Absolute magnitudes derived from GAIA DR2 distances and rate of decline for the t_2 parameter, fitted by the S-shape reverse relation. The solid line is the best fit eq. 15, where the two dashed lines identify the $\pm 3\sigma$ strip, with $\sigma = 0.26$ mag. The dashed gray lines correspond to the fit obtained with a linear model. The magnitude of the intercept of the MMRD with the y-axis is $M_V = -8.85$ mag for the t_2 parameterization. (Lower panel) Same analysis, but for the t_3 parametrization. The two dashed lines identify the $\pm 3\sigma$ strip, with $\sigma = 0.23$ mag. The dashed red line is the theoretical track (eq. 14). The magnitude of the intercept of the MMRD with the y-axis is $M_V = -9.02$ mag. Data have been obtained from Tab. 9 and Tab. 10. $\chi^2_{S\text{-shape}} = 1.02$, $\chi^2_{\text{linear}} = 1.05$.

Chandrasekhar limit. The flattening at faint level of luminosity, characterized by $m_{pg} \sim 19$ mag, is relatively near to the photographic detection limit of the M31 surveys by Rosino, and may well represent the bright wing of Eddington novae which are populating the bottom of the MMRD relation. Therefore we cannot exclude that the faint flattening might be the result of an observational bias. However we note that the nature of the outburst for very low mass WD has not been fully explored and therefore it is still possible that the observed flattening is a consequence of the physics of the outburst.

5.2 How good is the MMRD?

Recently, by taking advantage from nova distances determined by GAIA DR2, the MMRD has been tested by S18 and SG19 and they have achieved very different conclusions. The former author concluded that *"...the MMRD should no longer be used"* whereas the latter authors reached opposite conclusions. This conflicting results are even more puzzling after considering that the two samples share 15 novae and the parameters used to compute the MMRD are very similar. Then we have formed a "bona fide" sample of 29 objects to test the MMRD relation. This sample includes the "golden sample" of S18, the SG19 novae and V1500 Cyg, whose parallax measured by GAIA DR2 is $p = 0.777 \pm 0.187$ mas (see Table 9). Two novae have been removed from S18 "golden sample" sample: BT Mon and V1330 Cyg, whose magnitudes at maximum and rates of decline reported by the author are incorrect. Correct values can be inferred from original data reported in the literature. The maximum light of BT Mon was certainly much brighter than the value of $m(\max) = 8.5$ mag reported by S18. This conclusion can be reached after noting that spectroscopic observations of BT Mon (McLaughlin, 1941) revealed that this nova was in the so called "Orion" phase when it was observed at $m=8.5$ mag. Since Orion phase is subsequent to the principle and diffuse enhanced spectra (see Sect. 4), this magnitude value cannot correspond to the magnitude at maximum of the nova. On the basis of these arguments, Smith *et al.* (1998) suggested that the maximum of nova BT Mon could be as bright as $m_v \approx 4$ (see SG19 for a deep discussion of the BT Mon case). Also the estimate of the magnitude at maximum of V1330 Cyg, $m(\max)=9.9$ mag, reported by S18 appears incorrect. According to Ciatti and Rosino (1974):

"the maximum was lost. At the time of the discovery the nova was just at the end of the early decline, in the transition phase, showing the typical fluctuations of a moderately fast nova.... Since in general the transition takes place when the nova has fallen about 3.5 mag below maximum ... it is likely that the maximum was attained near May 15 with 7.5."

The same authors estimate $t_2 = 12^d$ which is much shorter than $t_2 = 161^d$ reported by S18. In view of these uncertainties we have exclude from our "bona fide" sample both BT Mon and V1330 Cyg.

After applying the reverse S-shape function, derived from M31 and LMC novae, to our "bona fide" sample, we get the following absolute calibration for the MMRD of the MW:

$$M_V = -7.78(\pm 0.22) - 0.81 \times \arctan((1.32 - \log t_2)/0.23) \quad (15)$$




Fig. 36 GAIA DR2 distances vs. MMRD distances. The red point marks CI Aql. MMRD distances are computed using the MMRD formulation of Eq. 15. Data from Tables 9.

In Fig. 36 we compare the distances obtained via GAIA DR2 with the distances obtained via MMRD. The plot shows a good agreement between GAIA DR2 and MMRD distances. Basically the most deviating object is the Recurrent Nova CI Aql, which is characterized by one of the highest correction for absorption ($A_V = 2.6$ mag). The inspection of the residuals indicates that above ~ 2 kpc the MMRD tends to overestimate the distances up to $\sim 30\%$. Unless to figure out some bizarre mechanism for which the MMRD relation is followed only by novae nearby the Sun, the most simple explanation of this behavior is not a failure of the MMRD but a generalized underestimation of the measure of A_V toward the more distant novae. Our analysis support the SG19 results, contrary to S18 conclusions.

6 Nova populations in external galaxies

The use of galactic data to study the properties of the Milky Way nova populations has been often questioned because of the observational bias due to both interstellar absorption in the galactic disk and our position within the Galaxy. These effects can be largely minimized by studying the spatial distribution of nova populations in external galaxies where novae are all at essentially the same distance.

6.1 Novae in M31

Due to its closeness and the relatively high frequency of nova events, the M31 nova population has been the only one extensively studied. However a simple skimming through past literature reveals the existence of different ideas on the nova population assignment: intermediate populations (Arp, 1956; Rosino, 1964), halo population (Wenzel and Meinunger, 1978), mainly bulge population (Ciardullo




Fig. 37 Surface density profiles of M31 nova population (light and filled dots) as a function of the semimajor-axis distance from the center. The solid and dashed lines represent the photometric $r^{1/4}$ models of the bulge (Walterbos, 1986; de Vaucouleurs, 1958). The long dashed is the density profile of GC systems. The nova data (circles) follows the light profile of the bulge (Fig. 7 from Capaccioli *et al.* 1989).

et al., 1987; Capaccioli *et al.*, 1989; Shafter and Irby, 2001), mostly disk population (Hatano *et al.*, 1997). Although it is possible that a fraction of M31 novae assigned to the bulge by Ciardullo *et al.* (1987) and Capaccioli *et al.* (1989) are in fact physically related to the disk and their allotment to the bulge was a simple consequence of neglecting the geometrical projection effects (Hatano *et al.*, 1997), there are two arguments that strongly suggest that most novae in M31 mainly belong to the bulge. First of all Capaccioli *et al.* (1989) show a close similarity between the projected density distribution of the nova system with the decomposed light profile of M31 into the two main components: bulge and disk (see Fig. 37). In this plot, novae follow the solid line, i.e. the $r^{1/4}$ model for the bulge along the major axis resulting from the decomposition in bulge/disk components of the B band luminosity profile (Walterbos, 1986), while the dotted line is the exponential model of the major axis B light profile of the disk from the same author. It is apparent that the nova surface density follows the light profile of the bulge and unless severe selection effects, due to dust for example prevent the discovery of disk novae in M31, one can conclude that novae ignore the trend of the light of the disk. We note that Kaur (2016) results on M31 seem to exclude that dust can induce such a selection effect.

Shafter and Irby (2001) have reappraised the Hatano *et al.*'s conclusions, after comparing the spatial distribution of novae with the background light of M31. They indeed confirm that most novae in M31 originate in the bulge: likely about 70% and not less than 50%. Some interesting hints come also from spectroscopic observations (Shafter *et al.*, 2012). These authors find that about $\sim 80\%$ of M31




Fig. 38 The MMRD relation for 91 M31 novae. The two dashed lines mark the $\pm 3\sigma$ strips, with $\sigma = 0.17$ mag strip. The magnitude of the intercept of the MMRD with the y-axis is $M_{pg} = 15.76$ mag. Data have been obtained from Table 13

novae belong to the Fe II spectral class. In view of what we discussed in paragraph 4.1, we can conclude that most novae in M31 are "bulge" novae.

The first estimate of nova rate in M31 dates back to Hubble (1929) who derived from a sample of about eighty Novae an observed rate of 21.5/yr. After applying a number of correction factors he concluded: "*that thirty per year is a reasonable estimate of the frequency of novae in M31*". Not far from this result appears the estimate by Arp (1956). He obtained from an almost continuous 1.5 year survey carried out with the 60-inch telescope at Mount Wilson an observed rate of 19.85 per year, which he corrected to a "real" frequency of 26 ± 4 novae per year to account for systematic bias. By taking advantage of two nova surveys carried out at Asiago Observatory with the 1.22m (Rosino, 1964, 1973) and 1.82m (Rosino *et al.*, 1989) telescopes, Capaccioli *et al.* (1989) derived from a global sample of 142 novae, an observed rate of 29 ± 4 novae per year. Later analysis of Shafter and Irby (2001) based on 53 new discovered novae found a rate slightly larger than previous estimates: 37^{+12}_{-8} novae per year, and about 70% of them produced in the bulge. Darnley *et al.* (2006) reported in the framework of POINT-AGAPE microlensing survey a nova rate in M31 of 65^{+16}_{-15} novae/yr from an "observed" sample of 20. This is a full factor two larger than previously estimated. All nova rate data are summarized in Tab. 5.

A technical analysis of the completeness factors and criteria adopted by Darnley *et al.* (2006) to derive the latter result is outside the scope of this review. Nevertheless, we point out the following: according to Shafter and Irby (2001) the nova rate in M31 can be as high as ~ 50 novae/yr "*if the luminosity-specific disk and bulge nova rates are comparable*". On the other hand, Darnley *et al.* (2006) concluded that "*M31 bulge CN eruption rate per unit r flux is more than five times*

Novae	rate (yr^{-1})	author	method
85	30	Hubble (1929)	observations
30	26 ± 4	Arp (1956)	observations
142	29 ± 4	Capaccioli <i>et al.</i> (1989)	observations
72	$37 + 12 - 8$	Shafter and Irby (2001)	observations
22	$65 + 16 - 15$	Darnley <i>et al.</i> (2006)	observations
-	97	Chen <i>et al.</i> (2016)	Theory
-	106	Soraisam (2016)	Theory+ analysis of the bias

Table 5 Collection of nova rate measurements for M31.

greater than that of the disc.”. It looks like that the two conclusions are inconsistent with each other. A possible way out is that the nova rate in M31 is significantly higher than reported above. For example, Chen *et al.* (2016) by using different star formation histories such as a starburst case for early type galaxies, a constant star formation rate case for spiral galaxies and a composite case for ”early” spirals such as M31, estimated a new rate of 97 novae per year for Andromeda galaxy. These authors also found other interesting results. The nova rate at 10 Gyr in an elliptical galaxy is up to 20 times smaller than a spiral galaxy with the same mass as result of the different density of nova progenitors and ”selection effects” discussed in Sect. 6.6. Therefore the specific nova rate per unit of luminosity (or mass) in the ”Ellipticals” peaks around 1 Gyr and declines by 2 orders of magnitude at 10 Gyr. In ”Spirals” the specific nova rate peaks around 2 Gyr and declines by a factor of 4 after 10 Gyr. The specific nova rate for elliptical galaxies at 10 Gyr is $1 \div 2$ per $10^{10} M_{\odot} yr^{-1}$, which is consistent with previous predictions (Matteucci *et al.*, 2003) and observations (see Fig. 42). The normalized nova rate for Spiral at 10 Gyr is $\sim 20-40$ per $10^{10} M_{\odot} yr^{-1}$, significantly higher than measured in early type galaxies. This result is qualitatively in agreement with observations (Della Valle and Livio, 1994; Della Valle *et al.*, 2002; Alis and Saygac, 2014) although the ratio is much larger than observed. According to Chen *et al.* (2016) this mismatch may be due to either the incompleteness of past surveys or incorrect assumptions of the SF history in their models. By applying the normalized nova rates in spirals derived by Chen *et al.* (2016), to the LMC, one should observe between 6 and 12 novae/yr, which are unrealistic values for the nova rate in our satellite galaxy (see Sect. 6.3). Therefore we tend to downsize the case for incompleteness and to agree with the authors that the mismatch is possibly due to some incorrect assumptions on the SF history, although the trend of the normalized nova rates exhibited by early-type to late-type hosts is qualitatively consistent with observations.

Similar high frequency have been obtained by Soraisam (2016) after assuming that specific nova rates in the bulge and disc are the same and by correcting the Arp’s and Darnley *et al.*’s rate for an hypothetical bias against the discovery of fast novae due to the poor observing cadence of this survey. These authors estimated 106 novae yr^{-1} . We note that Arp survey consisted of about 1000 plates taken on 290 nights distributed over ~ 18 months, June 1953 - Jan 1955. These data imply about one observing night every due days and about three plates per night. With such a ”dense” sampling it is rather difficult to think that Arp might have missed a significant number of ”fast” and ”bright” novae. Arp indeed concluded: *”Consequently of the 30 novae discovered, for only 5 were the maxima missed by more than a day”*. Therefore the paucity of detection of fast and bright novae in M31 does not seem the result of an observational bias rather being due to the

poor efficiency of old stellar populations to produce fast novae, which originate by massive WDs. These novae are characterized by lifetimes of the order of a few $\times 10^7$ years (see Sect. 1.2) and therefore we do not expect that they are predominant in $\sim 10^9$ years old bulge stellar population. In the following we adopt $R_{M31} = 40^{+20}_{-10}$ novae/yr as "educated guess" for the frequency of occurrence of novae in M31.

6.2 Novae in M33

Until 90's the only existing estimates of the nova rate in M33 were based on the Hubble and Sandage (1953) and Arp (1956) observations who derive an approximate rate of about 1 nova/yr and an estimate by Sharov (1993) of 0.5 novae/yr. However, in the 60's a nova survey on M33 started at Asiago Observatory, although sparser than the M31 survey carried out in the same years. Those observations allowed to discover 5 novae and to conclude that the nova rate in M33 was not "so low as previously assumed" (Rosino and Bianchini, 1973). A new surveys was started a few years later at the 1.82m telescope and five more novae were discovered (Della Valle, 1988). Altogether these observations have provided the database to measure a rate of $R_{M33} = 4.6 \pm 0.9$ novae/yr (Della Valle *et al.*, 1994). A few years later Williams and Shafter (2004) started an H_α survey which led to the discovery of four novae brighter than $m_{H_\alpha} = 18$ mag, and to derive a Nova rate of 2.5 ± 0.7 novae/yr. In Tab. 6 we summarize the results reported by these three independent surveys. A grand total of 14 novae have been discovered during a total control-time of 5.5 yr. After extrapolating the observed number of novae to the whole M33 field, according to the correction factors provided by the different authors we find: $R_{M33} = 3.5^{+0.95}_{-0.8}$ (1σ poissonian error).

Nova Survey Telescope	C-Time yr	Novae obs	Corr. field	Novae
1.22m	0.85	5	1	5
1.82m	1.9	5	1.4	7
1.00m	2.75	4	1.75	7
	5.5	14		19

Table 6 Summary of Nova surveys in M33. Col. 2: control time of the survey; col.3: "observed" number of novae; col.4: correction factor to the whole galaxy field; col. 5: "observable" number of novae. Data from Rosino and Bianchini (1973); Della Valle (1988); Williams and Shafter (2004).

We note that to compute the control time of their survey Williams and Shafter (2004) assumed a set of template H_α lightcurves based on the peak mag and rate of decline derived for M31 novae. The authors are well aware of the possible drawbacks of their approach and indeed they warn the readers:

"in our analysis is the assumption that the lightcurve properties of novae in M31 which were used in the Monte Carlo and mean H_α lifetime nova rate calculations are representative of novae in M33. If M33 contains a higher percentage of "fast" novae compared with M31, then we may be overestimating the mean nova lifetime in M33 and thus underestimating the true nova rate."




Fig. 39 Novae in M33 galaxy (Della Valle *et al.*, 1994).

Their assumption is fully justified if one compares nova populations having similar properties (e.g. novae of the bulge of M31 with novae in an elliptical galaxy in Virgo), but it is not applicable here because M33 is an almost bulgeless galaxy (Bothun, 1992) and therefore its nova production necessarily originates in the disk, in particular, Fig. 39 shows that novae appear superimposed on the arms of the parent galaxy. Shafter *et al.* (2012) were able to carry out a spectroscopic survey on novae in M33, a notable piece of work given the small number of novae that can be discovered during a campaign. They found that only 2 out of 8 discovered objects were members of the Fe II Williams' spectroscopic class. This result is different (at a confidence level of $\sim 95\%$) from what expected on the basis of Milky Way (Della Valle and Livio, 1998) and M31 (Shafter *et al.*, 2011) results, where most nova populations ($\sim 80\%$) are Fe II objects. Therefore the assumption that the properties of novae in M33 were similar to those of novae in M31 is not anymore justified and the rate of 2.5 novae/yr should be regarded as a robust lower limit for the nova rate in M33. Fast Novae close to maximum light evolve more quickly than slow novae also if observed in H_α light although at slower rate than exhibited in m_{pg} light, possibly a factor ~ 1.5 rather than ~ 5 has derived from the comparison of t_3 for slow and fast novae. As a consequence the control time of Williams and Shafter (2004) is likely to be overestimated by $\sim 30\%$ and the nova rate in M33 underestimated by an equal amount. Therefore the nova rate in M33 is larger than previously estimated: $R_{M33} = 4.1^{+1}_{-0.8}$.

6.3 Novae in the Magellanic Clouds

The properties of the outbursts of LMC novae and their frequency of occurrence have been well established only recently. Until 90's only 20 objects - the first one




Fig. 40 Spectra of M33 novae. M33N 2006-09a appears to be a Fe II nova, while 2007-09a and 2009-01a are He/N systems. From Shafter *et al.* (2012).

observed in 1926 (Buscombe and de Vaucouleurs, 1955) - were available and a reliable estimate of the nova rate was jeopardized by the fragmentary nature of observations, in some cases even one discovery every ten years. A first summary of the available data until Nova LMC 1988 has been provided by Capaccioli *et al.* (1990). In spite of the paucity of the data these authors could compile the MMRD relation that showed a remarkable percentage excess of fast novae compared to M31.

The data could be fitted with the same MMRD derived from M31 novae, obviously after applying a different zero point. In other words the MMRD relation seems universal in terms of relation between peak magnitude and rate of decline but not as it concerns the relative percentage of the bright and fast novae with respect to faint and slow. This piece of observation has an important consequence. We have previously commented about the rates of decline from maximum light as a tracer for differences in the parameters of nova outbursts (Sect. 3.2), particularly




Fig. 41 The MMRD relation for 27 LMC. The two dashed lines mark the $\pm 3\sigma$ strip. 1σ corresponds to ≈ 0.32 mag. The magnitude of the intercept of the MMRD with the y-axis is $V = 10.03$ mag. Data have been obtained from Tab. 14

for the mass of the underlying WD. Most novae in the LMC are bright and fast declining then suggesting a membership to disk nova population which originates from more massive WDs. However other parameters are at play. For example Starrfield *et al.* (2000) explain this behavior on the basis of the metallicity of the accreted material. Kato *et al.* (2013) after studying the effects of low Z on the properties of nova outbursts achieved similar conclusions. These authors concluded that LMC novae should be brighter and eject more material than Galactic novae of the same type since the LMC metallicity is about one-third that of the Galaxy. The age of the parent stellar populations of fast and slow novae were also studied in details by Subramaniam and Anupama (2002) on a sample of 15 LMC novae. They were able to prove an upper and lower limits, for the age of parent stellar population of fast and moderately fast novae of 3.2 and 1 Gyr, unlike the stellar regions around slow novae mostly in the range 4-10 Gyrs.

More recently Shafter (2013) revised and collected data for 43 novae and through a detailed analysis of 29 lightcurves and 18 spectra, was able to confirm that LMC novae: i) have faster rates of decline than novae in the Milky Way and in M31; ii) show a high percentage of He/N and Fe IIb novae compared with M31. Therefore bright He/N disk-novae dominate the LMC nova production unlike M31 where faint Fe II bulge novae are predominating.

The measurement of the nova rate in the LMC is a recent story. The first estimate was provided by Graham (1979) at the end of a ~ 10 years survey which allowed to discovery 10 novae. He suggested a rate between 2 and 3 novae/yr, and today we can say that he was right. Capaccioli *et al.* (1990) found 2 ± 1 novae/yr,

Della Valle *et al.* (2002) found 2.5 ± 0.5 novae/yr and recently Mróz *et al.* (2016) from the OGLE survey derived 2.4 ± 0.8 novae/yr.

The estimate of the rate in the SMC is a more critical issue because a continuous monitoring of this galaxy for nova searches has never been carried out if not very recently thanks to the OGLE surveys (Mróz *et al.*, 2016). Here below, we describe four different approaches:

i) we can derive a first order estimate after assuming that the nova rate scales as the K-band luminosity of the host. After assuming $K_{LMC} = -2.17$ and $K_{SMC} = -0.43$, we derive a nova rate of ~ 0.5 novae/yr.

ii) A second measurement derives from the discovery of 4 novae in the SMC during 1999 (Welch and MACHO Collaboration, 1999a; Glicenstein, 1999b,a; Welch and MACHO Collaboration, 1999b). Although none of the objects has been spectroscopically confirmed, their lightcurves show sudden changes of the brightness by more than ~ 7 mag on time scales of days/weeks. This behaviour is typical for novae, although dwarf-nova bursts originated by the rare sub-class of WZ Sge (Ortolani and D’Odorico, 1980) show a similar photometric behaviour. However their intrinsic luminosity excludes this possibility. After assuming that these 4 events are novae, a simple application of poissonian statistic implies that the nova rate in the SMC has to be larger than 0.7 novae/yr (Della Valle *et al.*, 2002) at $\sim 99\%$ confidence level.

iii) A third approach is based on the intensive monitoring of the SMC operated in the early 2000s, for example by Bill Liller, which led to the discovery of 6 novae. We derive an "observed" rate of $R_{SMC} = 0.75$ novae yr^{-1} which should be regarded as a robust lower limit.

iv) Finally, Mróz *et al.* (2016) further extended the SMC nova database and derive a rate of $R_{SMC} = 0.9 \pm 0.4$ novae yr^{-1} fully consistent with the rates derived above. This is the rate that will be adopted hereinafter.

6.4 Novae in M81

The analysis of 79 plates obtained in the 1950-55 Palomar campaign for the discovery of Cepheids and Novae in M81 has been published many years later by Shara *et al.* (1999). These authors concluded that the spatial distribution of 23 discovered novae is *consistent with the two populations divisions as argued previously*. Particularly they estimate that the fraction of novae belonging to the bulge is not larger than $\sim 60\%$. From the data reported in the paper it has been possible to derive the control time of the survey and a rate of 24 ± 8 novae/yr in agreement within the errors with the rate estimated by Neill and Shara (2004) of 33_{-8}^{+13} novae per year. The average value that we adopt hereinafter is: 28_{-6}^{+7} novae per year.

6.5 Novae in Virgo and Fornax Ellipticals

First observations of novae in Virgo galaxies were reported by Hubble (Bowen, 1954) but only after Pritchett and van den Bergh (1987) the study of novae outside the Local Group of galaxies changed from sporadic interest into a mature field of research. These authors estimated on the basis of 9 detections in the B band

a rate of $\sim 160 \pm 57$ novae/yr, which almost corresponds to the rate in M49 because 8 out of the 9 novae were discovered in this galaxy. Shafter *et al.* (2000) used the Kitt Peak 4m telescope to monitor M87 in H_α filter and they discovered 9 novae from which a rate of 91 ± 34 novae/yr was estimated. Ferrarese *et al.* (2003) reported after a 55 days campaign carried out with HST/WFPC2 in V and I colors the discovery of 9 novae in M49, from which they derived a rate of 100^{+35}_{-30} novae/yr. Curtin *et al.* (2015) have monitored in H_α M87, M49, and M84 with the Canada-France-Hawaii 3.6 m telescope over about 1 year of control time and they estimated 154^{+23}_{-19} , 189^{+26}_{-22} and 95^{+15}_{-14} , respectively. By averaging these measurements we find $R_{M49} = 158^{+20}_{-17}$ and $R_{M87} = 136^{+19}_{-17}$ for M87. These rates are consistent with the frequency of occurrence of $\sim 90 - 180$ novae/yr estimated by Della Valle and Gilmozzi (2002) for the giant elliptical NGC 1316 in Fornax cluster. Recently Shara *et al.* (2016) have derived from a new survey on novae in M87 a higher rate of 363^{+33}_{-45} (see next paragraph).

6.6 Nova Rates as function of the Hubble type of the hosts

Several authors (Della Valle and Livio, 1994; Shafter *et al.*, 2000) have studied the trend of the nova rates as a function of the total and normalized luminosity of the parent galaxy. They suggested, though with different views that there may exist a systematic difference in the nova rate per unit of K luminosity in galaxies of different colors (Hubble types) and/or classes of luminosity (see their Fig. 3 and 4). One result on which both teams seem to converge is that "blue" galaxies of the sample, characterized by low luminosity classes are more prolific nova producers than early-type spirals and ellipticals by a factor ~ 3 . The current available data are summarized in Tab. 7 with obvious meaning of the acronyms.

Analysis of the data reported in Fig. 42 confirms that "red" galaxies, say, redder than $(B-K) \sim 3.2$ show a constant trend of the normalized nova rate, close to ~ 2 while blue systems, belonging to low luminosity class galaxies, exhibit a normalized luminosity a factor at least ~ 3 times larger.

The existence of an overproduction of nova outbursts in the LMC and SMC and to a smaller extent in M33 might be a consequence of the so called 'selection effects' on the nova frequency, investigated by Truran and Livio (1986) and Ritter *et al.* (1991). It is due to the different recurrence times and number of nova progenitors per unit of luminosity in the respective hosts. The argument is the following: from Della Valle and Livio (1998) we derive an average magnitude at maximum for a typical nova originating in an old stellar population ("bulge nova") of $M_B \sim -7$ mag. The same authors derive $M_B \lesssim -9$ mag for the average nova produced in a star forming stellar population ("disk nova"). After using the relation obtained by Livio (1992) between the magnitude at maximum and the mass of the underlying white dwarf, we obtain: $M_{WD} \sim 0.8 M_\odot$ and $M_{WD} \sim 1.2 - 1.3 M_\odot$ respectively (see also SG19). We know that nova systems in disk dominated galaxies result in more frequent nova events due to the shorter nova recurrence time associated with massive WDs. For example, from Yaron *et al.* (2005) we obtain for Novae characterized by the above reported masses, recurrence times of $\sim 4.5 \times 10^4$ years (by interpolating data of their Tab. 3 for $M=0.65$ and $M=1 M_\odot$, $\dot{M}=10^{-9}$ and $T=50,000K$) and 3×10^3 yr for a $M=1.25 M_\odot$, $\dot{M}=10^{-9}$ and $T=50,000K$. On the other hand, according to Cummings *et al.* (2018) and assuming a Salpeter IMF

Galaxy	B (mag)	A_{gal} (mag)	A_{int} (mag)	(B-K) (mag)	(m-M) (mag)	Rate (#/yr)	L_K/L_\odot (10^{10})	ν_K (*)	Ref.
M31	4.36±0.02	0.46	0.2	3.85	24.38	40 $^{+20}_{-10}$	8.0	5.0	1,3,10,16
N5128	7.84±0.06	0.50	0.0	3.38	28.12	28 $^{+7}_{-7}$	6.2	4.5	4
M33	6.27±0.03	0.18	0.40	2.87	24.64	4.1 $^{+1}_{-0.8}$	0.7	6.1	6,14
LMC	0.91±0.05	0.26	0.13	2.74	18.50	2.4 $^{+0.8}_{-0.8}$	0.3	8.4	5,20
SMC	2.70±0.10	0.20	0.41	2.71	18.99	0.9 $^{+0.4}_{-0.4}$	0.1	10.4	11,20
M49	9.37±0.06	0.10	0.00	4.30	31.06	158 $^{+20}_{-17}$	51.1	3.1	2,13,19
M100	10.05±0.08	0.11	0.06	3.84	31.04	25 $^{+5}_{-5}$	17.7	1.4	7
M87	9.59±0.04	0.10	0.00	4.17	31.03	136 $^{+19}_{-17}$	36.0	3.8	9,19
						363 $^{+33}_{-45}$		10.1	21
M51	8.96±0.06	0.15	0.12	3.43	29.42	18 $^{+7}_{-7}$	7.5	2.4	9
M101	8.31±0.09	0.04	0.03	3.24	29.34	12 $^{+4}_{-4}$	10.4	1.1	9,17
M81	7.89±0.03	0.35	0.37	3.99	27.80	28 $^{+7}_{-6}$	7.9	3.5	8,15
N1316	9.42±0.08	0.09	0.00	4.15	31.66	135 $^{+45}_{-45}$	73.8	1.8	12
M84	10.01±0.12	0.18	0.00	3.76	31.32	95 $^{+15}_{-15}$	22.1	4.3	19
M94	8.99±0.05	0.08	0.09	3.72	28.21	5 $^{+1.8}_{-1.4}$	3.1	1.6	18

Table 7 Extragalactic Nova Data. References: Arp (1956) (1) Pritchett and van den Bergh (1987) (2), Capaccioli *et al.* (1989) (3) Ciardullo *et al.* (1990) (4), Capaccioli *et al.* (1990) (5), Della Valle and Livio (1994) (6), Ferrarese *et al.* (1996) (7), Shara *et al.* (1999) (8) Shafter *et al.* (2000) (9) Shafter and Irby (2001) (10), Della Valle *et al.* (2002) (11), Della Valle and Gilmozzi (2002) (12), Ferrarese *et al.* (2003) (13), Williams and Shafter (2004) (14), Neill and Shara (2004) (15), Darnley *et al.* (2006) (16), Coelho *et al.* (2008) (17), Güth *et al.* (2010) (18), Curtin *et al.* (2015) (19), Mróz *et al.* (2016) (20), Shara *et al.* (2016) (21).

the number of nova progenitors characterized by $\sim 0.8 M_\odot$ white dwarf is about 5 times larger than the number of novae with $\sim 1.2/1.3 M_\odot$ white dwarf. Therefore the observable outburst ratio between disk and bulge novae becomes:

$$\rho_{disk}/\rho_{bulge} \sim \Delta T_{bulge}/\Delta T_{disk} \times N_{disk}^{prog}/N_{bulge}^{prog} \approx 15/5 \sim 3, \quad (16)$$

in good agreement with observations.

Recently, Shara *et al.* (2016) derived from 41 novae discovered in a HST survey on M87 a global rate of 363^{+33}_{-45} novae/yr. If confirmed this high rate implies a specific (normalized) nova rate of about $7.9^{+2.3}_{-2.6}$ which is comparable to those exhibited by Magellanic Clouds (see Fig. 42). Therefore the density of nova outbursts in giant elliptical would be similar to that observed in dwarf irregular star forming galaxies. In turn this implies that the recurrence times of novae harbored in binary systems including light WDs - the only ones active in the old stellar populations of ellipticals - should be considerably shorter than estimated by several authors (Yaron *et al.*, 2005; Starrfield *et al.*, 2019), or that the number of potential progenitors of CNe vastly exceed the value expected from stellar evolution arguments.

One possible explanation suggested by Shara *et al.* (2016) is that all previous ground-based surveys for novae in external galaxies have missed most novae, especially the ones belonging to the hypothetical class of "fast" and "faint". There are two arguments against this hypothesis: i) Pritchett and van den Bergh (1987) survey on Virgo ellipticals was densely sampled with 11 observing runs distributed on 15 nights. None faint and fast novae were discovered but seven faint and slow




Fig. 42 The specific nova rate (nova rate per unit of luminosity of the host) plotted as a function of galaxy color. The red circle corresponds to the nova rate value in M87 as inferred by Shara *et al.* (2016). Data from Table 7.

declining objects. This behavior is reminiscent of the Arp’s result described in Sect. 6.1 and it is suggestive that old bulge stellar population are inefficient producers of bright and fast novae due to the lack of active nova systems containing massive WDs; ii) Ferrarese *et al.* (2003) survey was carried out with HST as well, so it is difficult to call for a massive observational bias affecting this survey. Recently Shafter (2017) through an independent analysis of the HST data resize the result of Shara *et al.* (2016) and concluded that

“...the global nova rate in M87 remains uncertain, both due to the difficulty in identifying bona fide novae from incomplete lightcurves, and in extrapolating observations near the center of M87 to the entire galaxy. We conclude that nova rates as low as 200 per year remain plausible.”

A speculative possibility is represented by a further develop of an interesting finding of Livio *et al.* (2002). After the serendipitous discovery of a SN Ia near the jet of the active galaxy 3C 78, these authors have examined the issue of whether jets can enhance accretion onto white dwarfs via Bondi accretion process. They also present evidence for observations of 11 novae in close proximity to the jet. From a simple application of the Bondi (1952) formula for accretion on a star, we derive that this process is efficient only for very low stellar velocities (a few km/s). If we assume typical star velocities of the order of 10-100 km s⁻¹ as measured by Williams *et al.* (1994) for nova systems in the Milky Way and the jet has a width of about 0.1 kpc, the transit time is less than 100 Myr during which the accreted mass onto massive WDs would be only 10⁻⁵-10⁻⁷ M_⊙ similar to the mass of nova accreted envelopes. This makes the accretion process from the ISM likely inefficient to trigger SN Ia explosions while it may play an important role to trigger nova explosions. If true, this result may account for the high frequency of nova events observed near the nucleus/jet of M87 (Madrid *et al.*, 2007) that might be considerably higher than observed in the rest of the galaxy.

6.7 Nova outbursts as a function of the Metallicity content and class of luminosity of the Hosts

The observed trend of the normalized nova rate might be also interpreted in terms of differences in the metallicity of the parent stellar population. For example Starfield *et al.* (1998) have discovered, on theoretical grounds, that novae originating in low metallicity stellar environments are more brighter at maximum light, than novae originating in populations characterized by solar metallicity and this fact is observed in the LMC, which is characterized by a metallicity that is about 1/3 of that of the Milky Way. The presence of a nova overproduction in blue galaxies with respect to red galaxies may also hint for the existence of a link between star formation and the frequency of occurrence of nova outbursts. For example, Yungelson *et al.* (1997) have suggested that in Irregulars and late dwarf spirals galaxies the nova rates is mainly determined by the current star formation rate in the disk, while the nova rate in elliptical galaxies depends mainly on the mass of the galaxy. Since the star formation rate in giant spirals was, several Gyr ago, much higher than the present one (Gavazzi and Scodreggio, 1996) it turns out that most stars in the disk of giant spirals are old and their nova population characterized by a frequency of occurrence of nova systems not too different from that occurs in their bulges. In addition since the present SFR decreases moving from late to early spirals, it is not unexpected to find that the integrated colors of giant spirals should be redder than the integrated colors of dwarf spirals where the SFR is still active. This scenario provides a simple explanation of the fact that giant late spirals, as M51 and M101 have redder colors than dwarf galaxies of comparable Hubble type and a nova frequency of occurrence similar to that of the bulges.

7 Novae in Globular Clusters

The researches for novae in globular clusters (GCs) have always given a very limited number of detections. Two Novae have been so far discovered in Milky Way GCs: M80 and M14. The former, T Sco 1861, occurred in M80 (Sawyer, 1938) and it was a moderately fast nova with a t_3 of about three weeks. The latter was a nova discovered by (Sawyer Hogg and Wehlau, 1964) on photographic plates taken with the 72 inches telescope at the Dominion Observatory in 1938. M14 has been imaged on 124 nights from 1932 to 1963 and the nova was detected during its final decline in all eight plates obtained in 1938. The post-nova candidate was possibly identified by Shara *et al.* (1986). The whole collection was formed by 2500 plates that have imaged 50 GCs. In a zero order approximation we can set a lower limit for the rate of novae in Milky Way GCs, after considering an effective survey time of about 30 years on 50 GCs and a single detection. By combining these data the rate of novae should be larger than 6.6×10^{-4} novae $GC^{-1}yr^{-1}$. In modern times nova outbursts have been unsuccessfully searched by Ciardullo *et al.* (1990) in 54 globular clusters of M31 on a mean effective survey time of about 2 years. Similar results were obtained by Tomaney and Shafter (1992) that carried out a survey on 200 M31 GCs over 1 year of effective surveillance time. The null result for both surveys can provide some constraints on the Nova rate in GCs. In particular, from simple Poisson statistics we find that the probability of obtaining a null result are 95% and 5% when the expected values are 0.05 and 3.0, respectively. Therefore one

should expect that the nova rate in GCs may be larger than the 95% probability value, i.e. 1.6×10^{-4} novae $GC^{-1}yr^{-1}$ and hardly higher than the 5% probability value, i.e. 10^{-2} novae $GC^{-1}yr^{-1}$. The lower limit is consistent with the result of Shara *et al.* (2004), who estimated a rate for the M87 of $\sim 4 \times 10^{-3}$ novae $GC^{-1}yr^{-1}$. The results based on X-ray surveys on M31 GCs (Henze *et al.*, 2013) of: $1.5 \times 10^{-2}GC^{-1}yr^{-1}$ matches the upper limit on the GCs nova rates obtained via optical surveys. Finally Curtin *et al.* (2015) and Doyle *et al.* (2019) find a GC nova rate of: $\sim 4 \times 10^{-4}$ and 5×10^{-4} novae $GC^{-1}yr^{-1}$, respectively.

The scanty statistics results in a global rate affected by almost two orders magnitude uncertainty: between a few $\times 10^{-4}$ and 10^{-2} novae $GC^{-1}yr^{-1}$.

During their nova survey on M49, M87 and M84, Curtin *et al.* (2015) found that two out of 84 novae were discovered in the GC systems of these galaxies. The GC systems correspond to 2.4% of detected novae, while the mass of GCs comprise only the 0.23% of the combined mass of the galaxies. Thus, these authors speculate that the nova rate in GCs might be enhanced by a factor 10 relatively to the nova rates measured in the host galaxies. Such a high nova rate in GCs might receive some support by the study of (Kato *et al.*, 2013) who found on theoretical grounds that the nova population in GCs should be mainly formed by fast novae. These novae are easily missed during low-cadence surveys that characterized the search for CNe in such stellar systems. On the other hand, Cao *et al.* (2012) by studying the M31 globular clusters system concluded that the M31 GC specific nova rate is not significantly higher than the Andromeda specific nova rate. We should take both conclusions with some degree of caution because they are still dominated by small number statistics.

8 The MMRD Relationship in Galaxies of different Hubble types

Fig. 43 shows the maximum magnitude vs. rate of decline relationship for LMC, M31 and Virgo novae. This plot indicates that the nova production in the LMC is clearly biased towards fast and bright novae, whereas the M31 nova population exhibits a prominent ‘slow’ component. By the same token the trend exhibited by novae discovered in Virgo. These trends are confirmed by a Kolmogorov-Smirnov test carried out on the rates of decline distributions of the respective nova populations. We find that the lightcurves of M31 novae mimic well the trend exhibited by the Virgo nova population (KS Probability = 0.7), but deviate strongly from the behaviour of the LMC novae (KS Probability = 0.005) which are on average faster. It is not obvious to explain this behavior in terms of an observational bias: indeed the brightest novae are detected in the nearest galaxy (LMC) and they would be missed in the more distant ones (Virgo), while faint novae would be detected in 15-20 Mpc away systems and missed in nearby galaxies, such as the MCs. The apparent magnitudes at maximum of novae belonging to the faint tail of the LMC, reach $V=12-13$ mag which are well within the capability of surveys currently in progress (Mróz *et al.*, 2016), therefore faint and slow novae in MCs are intrinsically rare objects. Obviously it is possible that some fast nova is missed in poor sampled surveys (Rosino *et al.*, 1989) but this doesn’t seem the case for the Arp (1956), Pritchett and van den Bergh (1987) and Ferrarese *et al.* (2003) surveys which observations were carried out - on average - every other night.




Fig. 43 The comparison among the MMRD relationships for LMC, M31 and Virgo novae. The best fit is the solid line described by eq. 15. The two dashed lines mark the $\pm 3\sigma$ strips.

Since the luminosity at maximum and the speed class of a nova depends on the physical parameters that determines the strength of the outburst (Yaron *et al.*, 2005) the rate of decline should be regarded as a valuable tracer of intrinsic differences in the nova populations. The differences in the MMRDs find a simple explanation in the framework of the two nova populations scenario: nova systems in disk dominated galaxies are associated with more massive WDs then resulting in faster, intrinsically brighter and more frequent nova events (see Sect. 6.6). Given the relatively short lifetime for the nova activity of these systems (see Sect. 1.2) they can be detected in galaxies where the star formation is still active, while the nova population in old stellar systems is mainly formed by "old" nova systems that contain light WD primaries and are characterized by lifetime activity of the order of $\sim 10^9$ years. Therefore the lack of fast and bright novae in old stellar systems as elliptical galaxies appear to be real.

8.1 Outliers of the MMRD

The existence of outliers of the MMRD was first reported by Arp 1956, see his Fig. 32. Arp himself suggested that nova number 4 was a heavy absorbed object, "obscured" by 2.2 mag. Rosino (1973) found two more deviating objects Nova 48 and Nova 66 which were found to be spatially coincident with nova 79 and nova 81, therefore he classified them as RNe. Both novae have fallen below the MMRD, then providing the first suggestion that some RN (not all) might be outlier of the MMRD. Della Valle (1991) found another faint outlier of the LMC MMRD: Nova LMC 1951 which was about two magnitudes dimmed than expected from the measured rate of decline. He also found evidence for the existence of a family of superbright novae (see for example Fig. 44) brighter than expected up to ~ 1 mag above the $+3\sigma$ strip of the MMRD: Nova LMC 1990-1 (Schwarz *et al.*, 2001), Rosino's novae in M31, nr. 28, 57, 53, 36, 85 and one object in Virgo. In the following we have assumed as outliers the points that fall outside the $\pm 3\sigma$ strip. Faint and fast Novae are rare objects, so they may well be patchy absorbed objects or intrinsically faint novae (such as RNe, for example) which eject very light envelopes. More difficult is to explain the existence of "super-bright"




Fig. 44 The MMRD relation for the LMC reported in Della Valle (1991) showing the bright outliers of the relation, as open circles, and the faint ones, the filled circles. .

novae, because it would imply the existence of an extra-energy source other than nuclear burning. The spectroscopic and photometric evolution of nova LMC 1991 excludes its membership to the class of "red luminous novae" (Rich *et al.*, 1989; Mould *et al.*, 1990; Martini *et al.*, 1999). According to Iben and Livio (1993) one possible explanation is the dissipation of the orbital energy motion of the secondary engulfed in the expanding nova ejecta during the Nova common envelope phase. An alternative may be the relatively rare super-bright nova explosions which occur in CV-like systems formed by a cold low-mass degenerate dwarf accreting matter from a close low-mass companion (Iben and Tutukov, 1992).

More recently a paper by Kasliwal *et al.* (2011) claim on the basis of 20 novae discovered in M31 (10), M81 (6), M82 (3) and NGC 2403 (1) that this sample is inconsistent with the canonical MMRD. A close inspection of their fig. 11, reveals that only 2 objects out of 17 (for which was possible to determine the MMRD parameters) match the MMRD. Other 4 objects are borderline and consistent with the relation within the errors. In other words between 65% and 90% of novae in Kasliwal *et al.* (2011) sample appear to be outliers of the relation. This is a surprising finding if compared with the percentage of outliers discussed above. Taking these figures at their face values the MMRD should be a very poor relation, but this seems out of question after the recent GAIA DR2 results (see SG19 and fig. 36 in this paper). The robustness of the relation is confirmed by the fact




Fig. 45 Blue dots: M31 novae from Arp (1956) and Rosino (1964, 1973); Rosino *et al.* (1989). Red triangles are Kasliwal *et al.* (2011) novae for which the MMRD parameters were computed with sufficient accuracy. The two dashed lines mark the $\pm 3\sigma$ strips. Data from Table 15.

that the "canonical" MMRD is based on 119 M31 and LMC nova lightcurves, see Fig. 34, and only $\sim 5\%$ of them are outliers. The M31 data come from three different surveys obtained with three different telescopes (60-inch, mount Wilson and 1.22 and 1.82 m at Asiago Observatory) from two experienced observers as Arp (1956) and Rosino (1964, 1973); Rosino *et al.* (1989). LMC data are a collection of relatively sparse observations, but in spite of this, the MMRD relation for the LMC, see Fig. 41, shows only two (7%) obvious outliers. Therefore it is difficult to understand where the Kasliwal *et al.* (2011) high percentage of outliers is produced. The daily cadence of the Kasliwal *et al.*'s survey - which is not very different from the Arp one - does not seem at the origin of this high percentage of outliers. Kasliwal *et al.* (2011) finds t_2 values not unusual for nova systems and much longer than the t_2 of a few hours hypothesized for the ultra-fast nova population. In the attempt to clarify this point, in Fig. 45 we have added to the Arp and Rosino novae the eight M31 Kasliwal *et al.*'s objects for which the authors reported the magnitude at maximum and the t_2 rate of decline (see their Tab. 5). The use of apparent magnitudes rather than absolute magnitudes has the advantage to minimize the corrections to the original observations by avoiding the magnitude revision for background and foreground absorptions and removing the systematics due to the adoption of the distances toward each galaxy. The g mag have been transformed to m_{pg} through $m_{pg} = g - 0.05$ derived from the photometric corrections reported in Capaccioli *et al.* (1989) and Jordi *et al.* (2006).

The lack of errorbars do not allow to be more quantitative but we think that five objects match the relation and another one might be consistent within the uncertainties with it. The two faintest novae might well be patchy absorbed novae and/or RNe caught in outburst. Admittedly on scanty statistics the fraction of




Fig. 46 Blue dots: M31 novae from Arp (1956) and Rosino (1964, 1973); Rosino *et al.* (1989). Green crosses: M31 novae from Hubble (1929) whose magnitudes turned out to be systematically too bright by ~ 0.7 mag. The two dashed lines mark the $\pm 3\sigma$ strips. Data from Table 16.

outliers objects in M31 (2 out of eight) turns out to be consistent with previous findings.

Finally we note the following: the discovery of a new photometric sub-class of "faint" and "fast" classical novae by Kasliwal *et al.* (2011) was essentially based on their Fig. 11, where it appears that most of newly discovered novae fall into the faint and moderately fast region of the MMRD after revision of their apparent magnitudes for internal and foreground absorptions toward each galaxy and by assuming the corresponding distance moduli. Systematic behaviours exhibited by data can hide pitfalls and other explanations. For example a similar systematic behaviour, even if observed in a different region of the MMRD plane, is exhibited by Hubble (1929) novae. In Fig. 46 we have added the Hubble data to Arp and Rosino novae (Hubble data from Tab. VI of Capaccioli *et al.* 1989).

Most Hubble novae are systematically above the $+3\sigma$ strip of the MMRD. They seem to identify a sub-class of "bright" and "slow" CNe. In fact these data points simply suffer of a ~ -0.7 mag offset in the zero point of the photometric calibration.

However, the observational evidence for the existence of a new photometric sub-class of "faint" and "fast" classical novae has vanished after Cao *et al.* (2012) have published the data of their new survey on M31 novae. Fig. 47 has been obtained, by adding the Cao *et al.* (2012) dataset to the Arp and Rosino data, by using the photometric correction term $m_{pg} = R + 0.1$. We confirm all parameters reported by Cao *et al.* (2012) with one exception: the t_2 value for nova 2010-06b. The first maximum of this "slow nova" is followed by a sharp and fast decline after which we can observe secondary maxima followed by strong fluctuations in brightness and finally a more gentle decline. To compute the correct rate of decline

Nova	$t_{2,Cao}$	$t_{2, revised}$	V_{max}
2009-08b	> 35	45	17.76
2009-11b	> 37	90	18.25
2009-11d	> 17	25	17.72
2010-06a	> 31	40	16.88
2010-10a	> 9	15	17.03
2010-10b	> 41	60	17.44
2010-12b	> 13	17	15.56

Table 8 Fair quality data from the Cao *et al.* (2012) sample and revised estimates.




Fig. 47 Blue dots: M31 novae from Arp (1956) and Rosino (1964, 1973); Rosino *et al.* (1989). Red triangles: M31 novae from Cao *et al.* (2012) data. With the exception of three obvious outliers, Cao *et al.*'s data match the MMRD within $\pm 3\sigma$ (dashed lines). Data from Table 15.

we have applied the smoothing procedure described in sect. 3.1. We wish to point out that this "procedure" is standard (i.e. not "ad hoc") for measuring t_2 or t_3 in nova lightcurves. This is the case for nova 2010-06b (that shows some resemblance to the nova n. 36 of Rosino) for which a $t_2=25^d$ is a more realistic choice. The MMRD parameters for the fair quality data of the Cao's sample for which only lower limits to the t_2 were provided by the authors, have been directly derived from the published lightcurves (see Tab. 8 in Cao *et al.* 2012) and our results are fully consistent with the original estimates of Cao *et al.* (2012). All these data points match satisfactorily the MMRD relation and there are only two obvious outliers. The very bright nova 2009-10b, which achieved a magnitude at maximum of 14.9 mag (almost -10 mag in terms of absolute magnitude at maximum). Such bright novae are unusual in M31 but not unique events. Nova 2009-10b is a "twin" of novae nr. 28 (Rosino, 1964) and 57 (Rosino, 1973) indeed. The other one is nova 2010-06b that falls in the region of patchy absorbed CNe or RNe.

The third paper claiming the existence of a significant fraction of outliers in the MMRD relation is by Shara *et al.* (2017a). These authors based their claim on the Kasliwal *et al.* (2011) results and on their own survey on M87 Novae (Shara *et al.*,




Fig. 48 Blue dots: ten M49 and one NGC 4365 novae from Pritchett and van den Bergh (1987) and Ferrarese *et al.* (2003). Red dots: nineteen M87 novae from Shara *et al.* (2016). The magnitude of the intercept of the MMRD with the y-axis is $V = 22.54$ mag. Dashed lines mark the $\pm 3\sigma$ strip, with $\sigma = 0.36$ mag.

2016). They discovered 41 novae and determine the MMRD parameters for 21 lightcurves. A quantitative inspection of each lightcurve confirms the parameters derived by Shara *et al.* (2016) with the exception of nova 8 for which we find $t_2 = 7$ rather than 3.7 and novae 18 and 22 for which is impossible to derive the respective t_2 due to the incompleteness of the lightcurves.

Fig. 48 shows the positions of novae discovered in Virgo ellipticals (Pritchett and van den Bergh, 1987; Ferrarese *et al.*, 2003; Shara *et al.*, 2016) into the MMRD plane after assuming that M49 and M87 are basically at the same distance: $(m-M) = 31.10$ mag and 31.12 mag, respectively³. Virgo Novae are still consistent with the MMRD, but their distribution is characterized by a larger dispersion than it is observed in M31 and LMC. Indeed out of 30 objects (see Table 17) about 20% are outliers at level of $\pm 3\sigma$. The larger dispersion is likely due to the presence of systematics related to photometric measurements at very faint luminosity levels in three different galaxies carried out by three different teams (Pritchett and van den Bergh, 1987; Ferrarese *et al.*, 2003; Shara *et al.*, 2016).

The distribution of the data points is consistent with the MMRD relation, particularly the HST data do not show the existence of any faint and fast subclass of CNe, but possibly the existence of a superbright outlier similar to nova 2 in NGC 4472 discovered by Pritchett and van den Bergh (1987). Obviously, due to the time sampling of the lightcurves we cannot exclude that these surveys have missed the ultra-fast Novae with t_2 as short as a few hours (Yaron *et al.*, 2005; Shara *et al.*, 2017a). However due to the tiny masses of the accreted envelopes, $10^{-8} M_{\odot}$ (Shara

³ <http://http://leda.univ-lyon1.fr/>

et al., 2017a) their peak magnitudes should be considerably dim with respect to CNe and so they are rather hard to discover in such distant galaxies.

An important result obtained by Shara *et al.* (2016) is the discovery of a conspicuous faint and slow nova population, which is typical of an old parent stellar populations. From their published lightcurves we can infer that Novae from n. 25 through n. 41 peak between $V \sim 24$ mag and $V \sim 26$ mag and are characterized by t_2 values that unfortunately can be only roughly estimated, but they very likely range in the interval $40^d - 200^d$. These 17 data points would consistently populate the faint and slow region of the M87 MMRD in the range of rates of decline corresponds to 0.7-0.01, in terms of $\log(100 \times v_d)$ units.

9 Novae as Distance Indicators

There are several reasons for which novae were believed to be good distance indicators. They can be summarized as follows:

- i) Novae are bright, they can reach $M_B \sim -10$ mag, about 3 magnitudes more luminous than Cepheids of longer periods.
- ii) Unlike Cepheids that can be detected only in spirals, novae can be "easily" recognized also in ellipticals.
- iii) Intrinsic differences in the outburst properties, existing between novae associated with the bulge or spiral arms stellar populations do not affect the zero point of the MMRD relationship, but only change the relative percentage of fast and bright novae. Particularly we note that the luminous plateau of the MMRD relationship, characterized by an absolute magnitude of $M_V = -8.85 \pm 0.10$ mag (calibrated with GAIA DR2 nova distances), seems to be a "standard candle", because it is constant along the different Hubble types of the host galaxies.
- iv) There exists a good theoretical understanding of the MMRD, i.e. of the tool that it is used.

On the other hand, there are also some drawbacks for using novae as distance indicators:

- i) The cosmic scatter of the MMRD relationship is relatively small, $\sigma = 0.25$ mag, which brings about 12% uncertainty in the distance. However it is not so small as it would be required in the 1-2% cosmology accuracy era (Riess *et al.*, 2019).
- ii) Novae are transient objects so the time of their occurrence in the sky is unpredictable and therefore their observations are not easy to be scheduled. This aspect is not considered with favor by the various Time Allocation Committees.

9.1 The distance to M31, LMC and Virgo

To test the reliability of novae as distance indicators we measure the distances to M31, LMC and Virgo ellipticals by comparing the respective MMRD relations with the Galactic MMRD (Eq. 15 and Fig. 35) obtained using GAIA DR2 distance measurements, i. e. the "bona fide" sample described in Sect. 5.2 with the

additional case of V1500 Cyg⁴. Particularly we compare the absolute magnitude of the high luminosity plateau of the Galactic MMRD: $M_V = -8.85$ (see Fig. 35) with the apparent magnitudes of the high luminosity plateau of M31, LMC and Virgo MMRDs, which are $m_{pg} = 15.76$ mag, $M_V = 10.03$ mag and $M_V = 22.54$ mag, respectively. The distance modulus for M31 can be derived by combining the new data with Eq. 4 from Capaccioli *et al.* (1989):

$$(V - M_v)_o + A_B = [m_{pg} - M_V] - 0.26 = \quad (17)$$

$$= [15.76 - (-8.85)] - 0.26 = 24.35 \pm 0.31(1\sigma)$$

which is in excellent agreement with the Cepheids measurement of 24.38 mag obtained by Riess *et al.* (2012). Obviously, Novae due to the dispersion of the MMRD relation, provide the distance to M31 with an accuracy that is three times less than that provided by Riess *et al.* (2012).

The comparison between Galactic and LMC MMRD relations finds a distance modulus:

$$(V - M_v)_o = 10.03 - (-8.85) - 3.16 \times E(B - V) = 18.44 \pm 0.41(1\sigma) \quad (18)$$

The color excess $E(B - V) = 0.14 \pm 0.05$ mag has been derived from 6 novae reported by Shafter (2013) and it implies $A_V = 0.44$ mag. Despite of scanty statistic this value is in good agreement with $A_V = 0.3 \pm 0.1$ mag derived by Capaccioli *et al.* (1990) from observations of reddened LMC Cepheids. The distance modulus of $(m - M) = 18.44$ mag is in excellent agreement with the eclipsing binary distance of $(m - M) = 18.477 \pm 0.004$ (statistical) ± 0.026 (systematic) mag, obtained by Pietrzyński *et al.* (2019) by studying 20 eclipsing binaries in the LMC. Novae exhibit an uncertainty on the distance to the LMC of $\sim 19\%$ which is much larger than the one derived by Pietrzyński *et al.* (2019). Finally the comparison between the Galactic and Virgo MMRDs finds, after assuming $A_V = 0.07$ mag:

$$(V - M_v)_o = 22.54 - (-8.85) - 0.07 = 31.32 \pm 0.44(1\sigma) \quad (19)$$

in agreement within $\sim 1\sigma$ with recent measurements of Virgo cluster ellipticals of $(m - M)_{M49} = 31.10$ mag, and $(m - M)_{M87} = 31.12$ mag.⁵

9.2 Novae as Standard Candles

The concepts of "distance indicators" and "standard candles" are often interchangeably used. In fact, they are different. The observation of a SN-Ia (standard candle) allows, at least in principle, to immediately compare the apparent and absolute magnitudes, and then to compute the distance to the host galaxy by

⁴ The GAIA DR2 distance of V1500 Cyg used in this paper, $d = 1.29 \pm 0.31$, can be derived using the parallax measurement as provided in the GAIA DR2 data archive: <https://gea.esac.esa.int/archive/>

⁵ see Hyperleda archive - <http://leda.univ-lyon1.fr>

observing a single object. On the other hand if we observe a single Cepheid (distance indicator) this is not enough to make a direct distance measurement of the distance to the host galaxy. We need first to observe a number of Cepheids with different periods in such a way to determine the "P-L" or the "P-L-C" relations, and then we use these relations to determine the distance. In this respect when we use the MMRD relation, novae behave as distance indicators.

The most popular method for using novae as "standard candles" is the so called Buscombe and de Vaucouleurs (1955) law. They first pointed out that all novae, irrespective of their rates of decline, have about the same absolute magnitude 15 days past maximum. This is a simple consequence of the MMRD relationship for which the most luminous novae decline faster and viceversa. As a consequence there is an epoch in which the lightcurves of fast and intrinsically bright novae cross the lightcurves of intrinsically faint and slowly declining objects. There are several calibration of the $\langle M_{15} \rangle$ magnitude: $\langle M_{V15} \rangle = -5.24 \pm 0.15$ mag (van den Bergh and Younger, 1987), $\langle M_{V15} \rangle = -5.60 \pm 0.43$ mag (Cohen, 1985) $\langle M_{V15} \rangle = -5.69 \pm 0.42$ mag (Capaccioli *et al.*, 1989). Downes and Duerbeck (2000) measured the distances of a dozen of galactic novae via nebular parallaxes and obtained $\langle M_{V15} \rangle = -6.05 \pm 0.44$ mag. Darnley *et al.* (2006) found $M_{I15} = -6.3 \pm 1$ mag. We know that a nova at maximum light has a color index of $(B - V)_0 = 0.23$ mag (van den Bergh and Younger, 1987) that corresponds to a spectral type between A7 and F0, which in turn it is characterized by $(V - I)_0 \approx 0.4$ (Allen, 1973). This color index implies $M_{V15} \approx -5.66 \pm 1.00$ mag, after correcting for $A_V = 0.23$ mag derived from $A_B = 0.31$ mag (Burstein and Heiles, 1984). SG19 from a sample of 17 galactic novae have recalibrated the M_{V15} with GAIA DR2 distances and used a homogeneous treatment of the reddening correction (Selvelli and Gilmozzi, 2013). They find a weighted mean of $M_{V15} = -5.71 \pm 0.10$ mag.

From the Arp's high quality sample of 23 M31 novae we compute $m_{pg15} = 18.98 \pm 0.58$ mag. The distance modulus can be obtained by the usual formula (see above) and $A_B = 0.31$ mag (Burstein and Heiles, 1984):

$$(V - M_v)_0 + A_B = -(m_{pg15} - M_{V15}) - 0.03 = 24.35 \pm 0.59 \quad (20)$$

which agrees with the MMRD distance within a few hundredths of magnitude, admittedly with a lower accuracy. For the LMC we assume $M_{V15} = 13.05 \pm 0.25$ mag (Capaccioli *et al.*, 1990) and $A_V = 0.44$ mag (see above) and we find:

$$(m - M) = 18.32 \pm 0.27 \quad (21)$$

in agreement within about a tenth of magnitude with the MMRD distance.

Ferrarese *et al.* (2003) by using 5 novae discovered in M49 find a $M_{V15} = 24.77 \pm 0.19$ mag, which would implies $M_{V15} = -6.26 \pm 0.19$ mag, which is about 2.5σ brighter than the SG19 calibration. This result is borderline but it is not totally off the wall after considering the small number (five) of objects in the Ferrarese *et al.*'s sample.

10 The role of Novae in the Galactic Nucleosynthesis

Novae may significantly contribute to the galactic nucleosynthesis (Gehrz *et al.*, 1998) because they produce considerable concentrations of β^+ isotopes such as ^{13}C




Fig. 49 Histogram of the nova mass envelopes. Data from Warner (2003); Della Valle *et al.* (1997, 2002); Mason *et al.* (2005); Ederoclite *et al.* (2006); Izzo *et al.* (2015); Molaro *et al.* (2016).

¹⁵N (Sparks *et al.*, 1978; Peimbert and Sarmiento, 1984; Williams, 1985) ²²N, ²⁶Al (Hillebrandt and Thielemann, 1982; Kolb and Politano, 1997), ²²Ne (Livio and Truran, 1994), ⁷Li (Arnould and Norgaard, 1975; Starrfield *et al.*, 1978; D’Antona and Matteucci, 1991; Izzo *et al.*, 2015) and ⁷Be (Tajitsu *et al.*, 2015; Molaro *et al.*, 2016). The abundances of these elements in nova ejecta are significantly enhanced with respect to solar concentrations. The quantitative assessment of the contribution of novae to the chemical evolution of the Milky Way needs three quantities: i) the galactic nova rate; ii) the mass of the ejected envelopes; iii) the abundances of heavy elements in the nova ejecta. The Galactic nova rate has been extensively discussed in Sect. 3.5 and in the following we adopt a frequency of occurrence for Novae in the Milky Way of about 25 novae/yr.

10.1 The mass of the ejecta

We present in Fig. 49 the estimates of the mass distribution of the ejecta collected from literature. The data distribution peak at $\times 10^{-4}$ M_⊙. Fig. 50 shows that the ejected mass marginally correlates (95% c.l.) with the class of speed of the nova, according to:

$$\log M_{env} = 0.45 \times \log t_2 + 0.24 \quad (22)$$

This result is not unexpected. Since the more massive white dwarfs need, on average, to accrete less material to achieve the conditions to trigger the outburst,




Fig. 50 The correlation between the nova envelopes mass vs. rate of decline. Mass of the envelopes in units of $\log (10^{-5})M_{\odot}$. Data from Gehrz *et al.* (1998) and Warner (2003).

they should eject less massive envelopes. Now we are in the position to assess - at least as order of magnitude - the contribute of a nova system to the chemical enrichment of the host galaxy. We derive that about $0.25M_{\odot}$ are processed through the nova ejecta over 100 years which is the characteristic time scale for Supernova explosions in the Milky Way. This figure has to be compared with $\sim 10 \div 20M_{\odot}$ ejected by a typical core-collapse SN. It follows that only nuclei enriched, in the nova ejecta, by a factor $\sim 50 \div 100$ can contribute significantly to the galactic nucleosynthesis. In old and passive stellar systems like the elliptical galaxies where the formation of new stars has ceased long ago and therefore the rate of core-collapse is practically zero, the contribute of novae may be more significant. In this systems only SNe-Ia occur and these SN explosions are the result of the disruption of a CO WDs that release about $1.3M_{\odot}$ into the interstellar gas of the parent galaxy every 100 years or so.

10.2 The abundances in the nova ejecta

Recognition of unusual chemical abundances in the nova ejecta dates back many years (Payne-Gaposchkin, 1957; Williams, 1977; Gallagher and Starrfield, 1978). This topic has been reviewed several times (Livio and Truran, 1994; Gehrz *et al.*, 1998; José and Hernanz, 1998) and more recently by José (2012) and (Hernanz, 2015). One point stands clearly: all novae, for which reasonable data are available (even admittedly characterized by large uncertainties), appear to be enriched in either He or/and heavy elements. This result is not unexpected because TNR proceed in the accreted H-rich envelope by means of the CNO burning chain. Fig. 51 illustrates the average trend of the heavy elements abundance in the nova ejecta as a function of the rate of decline. Two facts emerges: 1) the available data suggest that the enrichment into the nova envelopes is normally below the




Fig. 51 Abundances in the nova ejecta vs. rate of decline. The data do not show a significant correlation. The grey horizontal line represents an over abundance of 50 times the solar one. Data from Gehrz *et al.* (1998) and Warner (2003).

~ 50 time threshold 2) the heavy elements abundance in the ejecta do not seem to correlate with the speed class of the novae.

10.3 Novae as Lithium Factories

During the TNR phase preceding the nova outburst, the synthesis of light chemical elements, and some of their isotopes, takes place. As soon as the the matter piles up onto the WD and the degeneracy pressure is reached at the bottom of the accreted layer the thermonuclear reactions (TNRs) start. The first reaction to be ignited is the proton-proton chain. As the temperature increases, convective motions bring unburned elements from the interior of the WD into the burning region, allowing the ignition of the CNO cycle. This nuclear reaction increases the production of β -unstable isotopes like ^{13}N , ^{15}O and ^{18}F , that will finally lead to the outburst and matter ejection (Sparks *et al.* 1977; Starrfield *et al.* 2009). During these processes, helium can also produce ^7Be , via the $^3\text{He} - ^4\text{He}$ reaction channel, an isotope which decays into the stable ^7Li after a half-life time decay of ~ 53 days (Cameron and Fowler, 1971).

Abundance studies of some of the best observed novae led to the evidence that CNO elements are very abundant in nova ejecta (Gehrz *et al.*, 1998), in particular of the order of ten times the Solar value for the nitrogen, oxygen, neon and magnesium (Vanlandingham *et al.*, 1996; Schwarz, 2002). On the other hand, we had little information about the lithium production. In classical novae, convective motions can transport a large quantity of freshly-produced ^7Be in its most external regions. In these external layers, lithium can survive and then ejected once the nova outburst is triggered by the TNR. An immediate consequence is that CNe may represent one of the most efficient astrophysical sources of lithium (Starrfield




Fig. 52 The distribution of the abundance of lithium ($A(\text{Li})$) as measured for stars with different metallicities ($[\text{Fe}/\text{H}]$) in the Galaxy (Spite and Spite, 1982; Spite, 1990). The black continuous line corresponds to the predictions of the chemical evolution models (Romano *et al.*, 2001) assuming the average yield of lithium provided by novae of $M_{\text{Li}} = 2.55 \times 10^{-10} M_{\odot}$ and with a primordial abundance of $A(\text{Li}) = 2.3$. The grey dashed region marks the upper and lower limit considering the uncertainties on the lithium yield. The red line represent the predictions of the chemical evolution model when neglecting the novae lithium yield. Dashed black line represents the evolution of the lithium if we start from a primordial abundance as observed by Planck of $A(\text{Li}) = 2.7$ (Coc *et al.*, 2014). Data from Charbonnel and Primas (2005); Lambert and Reddy (2004); Lodders *et al.* (2009); Sbordone *et al.* (2010); Ramírez *et al.* (2012). Plot from Izzo *et al.* (2015).

et al., 1978). However, long-term spectroscopic observations of CNe in outburst have never revealed the presence of Li in the nova ejecta (Friedjung, 1979).

Lithium is one of the most fragile element in the Universe, given its low binding energy per nucleon (Conti, 1968). Consequently, lithium is easily destroyed in many astrophysical environments, mainly via proton collision, at a temperature of $\sim 2 \times 10^6$ K. Lithium is also the only stable "metal" formed during the Big Bang Nucleosynthesis, given the absence of a stable element with atomic number $A=8$ (the only possible one, ${}^8\text{Be}$, has a very short life, $t \approx 10^{-17}$ s). Within the standard BBN theory, it is possible to predict the primordial lithium abundance from the estimate of the baryon-to-photon ratio parameter (Dunkley *et al.*, 2009; Fields, 2011; Coc *et al.*, 2013), which shows a discrepancy with the lithium abundance inferred from old stars observed in the Galactic halo, the so-called "Cosmological Lithium problem" (Spite and Spite, 1982; Sbordone *et al.*, 2010). Similarly, the lithium abundance inferred from younger Galactic stellar populations is much larger than the abundance estimated from the BBN (Spite and Spite, 1982; Spite, 1990), then suggesting the presence of unknown lithium factories in the Galaxy, this is the so-called: "Galactic Lithium problem" ((D'Antona and Matteucci, 1991; Matteucci *et al.*, 1995)). In recent years different astrophysical sources have been proposed for the lithium nucleosynthesis such as: the cosmic-ray spallation (Reeves

et al., 1990), the hot bottom burning in low-mass red giant branch (RGB) stars (Sackmann and Boothroyd, 1999), intermediate asymptotic giant branch (AGB) stars (Sackmann and Boothroyd, 1992; Travaglio *et al.*, 2001), carbon stars (Abia *et al.*, 2001) but it turned out that they were unable to produce the full amount of Lithium observed in the Milky Milky ((Romano *et al.*, 2001)).

10.4 "The Hunt for Red Lithium"

The idea that Novae could be efficient Lithium producers has been predicted long ago (Arnould and Norgaard, 1975; Starrfield *et al.*, 1978) but it has never been confirmed (e.g. (Friedjung, 1979; Della Valle *et al.*, 2002)). The typical signature of lithium in the spectra of CNe is the resonance transition of its neutral state, which is a very tight absorption doublet centered at ${}^7\text{Li I } 6707.8 \text{ \AA}$. Although Friedjung (1979) did not detect any absorption line corresponding to ${}^7\text{Li I}$, he was able to put upper limits on the lithium abundance in nova ejecta. Lithium was extensively and unsuccessfully searched in the following years. A possible evidence for the presence of a transient ${}^7\text{Li}$ emission line was made for the case of Nova Vel 1999 (Della Valle *et al.*, 2002), but further analysis using UV spectral data from Hubble Space Telescope showed its origin as due to neutral nitrogen (Shore *et al.*, 2003).

The presence of Lithium in CNe remained a mystery until 2013. Two very bright novae were discovered and then observed with high-resolution spectrographs by two distinct groups: V339 Del and V1369 Cen. The presence of blue-shifted absorption lines due to ${}^7\text{Be II}$ was reported in the late (more than 50 days from the outburst) spectra of V339 Del (Tajitsu *et al.*, 2015), while early spectra of V1369 Cen revealed the presence of an absorption feature that was confirmed to be due to the resonance transition of neutral Lithium at the expanding velocity of $V_{exp} = -550 \text{ km/s}$ (Izzo *et al.*, 2015). For this latter case, by using an approach similar to the one used by Friedjung (1979), the total mass of lithium ejected was computed to be $M_{Li} = (0.3 - 4.8) \times 10^{-10} M_{\odot}$. After assuming this amount of lithium as ejected by all slow novae in the Galaxy, and using accurate numerical codes developed to compute the chemical evolution of the Galaxy (Romano *et al.*, 2001), it was proved that novae can explain the observed over-abundance of lithium observed in young stellar populations (Spite and Spite, 1982; Fields *et al.*, 2014; Izzo *et al.*, 2015). The detection of Li I in the early spectra of V1369 Cen does not represent the unique case so far. In 2015 an other bright slow nova was discovered in the direction of the Sagittarius constellation, V5668 Sgr and similarly to V1369 Cen, early spectra showed the presence of a faint absorption line that corresponds to the Li I $\lambda 6708$ transition blue-shifted at a velocity of $v_{exp} \sim 500 \text{ km/s}$ (Wagner *et al.*, 2018). The same velocity is displayed by other typical transition observed in novae spectra as Na I D2 $\lambda 5890$ and Fe II $\lambda 5169$. This nova also showed the presence of blue-shifted absorptions related to the resonance doublet of ${}^7\text{Be II}$ $\lambda\lambda 3130/31$ (Molaro *et al.*, 2016). By following a similar approach Molaro *et al.* (2016) have estimated the ejected mass of ${}^7\text{Be}$ to be $M_{Be, Sgr} = 7 \times 10^{-9} M_{\odot}$. In passing we note that Prantzos (2012) in a detailed review on possible lithium factories in the Galaxy, explicitly mentioned that:

"if novae provide $\sim 50\text{-}65\%$ of solar lithium, they ought to have a Li yield of $\sim 10^{-9} M_{\odot}$."




Fig. 53 The spectrum of V1369 Cen observed 6 days from the discovery with FEROS. The upper panel shows the spectrum centered at the lithium resonance transition wavelength of Li I 6708 Å, while the lower panel shows the spectrum centered at the Na I 5890 Å. The presence of Li I 6708 is confirmed by its observed ejecta velocity, which is the same of Na I D and other absorption features in the spectrum (Izzo *et al.*, 2015).

A prediction in excellent agreement with the result derived from observations. After assuming a typical nova life-time of 10^{10} yrs, Molaro *et al.* (2016) suggest that a dozen of novae/yr like V5668 Sgr can produce the total amount of ${}^7\text{Li}$ observed in the Galaxy.

In recent years, additional evidences of ${}^7\text{Be}$ have been found in other novae (Tajitsu *et al.*, 2016; Izzo *et al.*, 2018; Selvelli *et al.*, 2018; Molaro *et al.*, 2020).

11 Novae as Supernova-Ia progenitors

SNe-Ia are commonly referred as standard candles even if it would be more appropriate to describe these objects as "standardizable" candles because their high luminosity at maximum correlates with their rate of decline. Differently from CNe the intrinsically brighter SNe-Ia are characterized by the slowest rates of decline (Phillips, 1993). The process of standardization works well on empirical basis (Perlmutter *et al.*, 1998; Riess *et al.*, 1998; Schmidt *et al.*, 1998) but still it is far from being completely understood (Mazzali *et al.*, 2001; Wygoda *et al.*, 2019) and the uncertain nature of the binary systems does not help to clarify this issue. SNe-Ia




Fig. 54 The spectrum of V5668 Sgr (left panel) and V407 Lup (right panel) observed 88 days and 8 days, respectively, after the nova discovery (Molaro *et al.*, 2016; Izzo *et al.*, 2018). The upper panel shows the spectrum centered at the ${}^7\text{Be}$ resonance transition wavelength of 3130.583 Å, while the lower panel shows the spectrum centered at the Na I 5890.93 Å. The presence of ${}^7\text{Be}$ is clearly marked by the combined identification of the two components of the doublet, at the same expanding velocity than other features, in this case of Na I 5890.

basics that can be directly derived from observations can be summarized as the following:

- SNe-Ia occur in early and late Hubble types;
- Occurrence in early type galaxies with SFR less or much less than a few $10^{-1} M_{\odot}/\text{yr}$ implies that SNe-Ia are not caused by the collapse of the core of stars with $M > 8M_{\odot}$;
- The lack of H and He in the ejecta points in the direction of highly evolved progenitors;
- The energy for unit of mass, i.e. $1/2 \times (10^4 \text{ km/s})^2$ is similar to the one obtained from the conversion of C (7.7 Mev b.e. per nucleon) into Fe (8.8 Mev b.e. per nucleon).

On the basis of these arguments, the common wisdom strongly suggest that SNe-Ia are the thermonuclear disruption of mass-accreting WDs in binary systems. Historically two scenarios are commonly accepted for producing SNe-Ia: the so called double degenerate (Webbink, 1984; Iben and Tutukov, 1984; Tornambe and Matteucci, 1986) where two CO WDs in a binary systems make coalescence as result of the lost of orbital energy for GWs and the single degenerate (Whelan and Iben, 1973; Nomoto, 1982a) in which a WD accretes material from a MS star (see Maoz *et al.* 2014; Livio and Mazzali 2018 for an update review on this topic). In the latter framework a SN-Ia can be produced by the explosions of WDs in recurrent- or symbiotic- nova systems, when the respective WDs approach the Chandrasekhar limit after having accreted material from the companion stars. Because these Cataclysmic Systems typically contain massive WDs characterized by a high accretion rates, they are regarded as possible SN-Ia progenitors. Recently Starrfield *et al.* (2019) have found on the basis of 1-D hydrodinamical simulations

that all C-O WDs in the range of masses $0.6\text{--}1.35M_{\odot}$, which accrete material from companions within a narrow interval of $\dot{M} \approx 1.6 \times 10^{-10} M_{\odot}/\text{yr}$ would grow in mass up to 80% of the accreted mass after each nova cycle. These authors conclude that novae can be an important channel for type Ia progenitors. These results are still preliminary and they should be taken with some degree of caution. We note for example that even if WDs are growing in mass because the ejected/accreted mass is smaller than 1, the Chandra mass might not be reached. This possibility depends among other things on the mass of the secondary and on the efficiency of the accretion onto the WD that is often defined as $\eta = \Delta m_{\text{growth}}/\Delta m_{\text{trans.}}$. Kato and Hachisu (1989) computed η for different sizes of the Roche lobes and they never found, even under the most favorable conditions (high value of M_{\odot}), a value larger than 0.3–0.4. For example a $1M_{\odot}$ WD which accretes material at $\dot{M} \sim 10^{-10} M_{\odot} \text{ yr}^{-1}$ and it is characterized by $\eta \lesssim 0.1$ requires a secondary star with a mass much larger than that of the primary, to reach the Chandrasekhar limit of the order of $M_2 \sim 0.35/\eta \approx 3.5M_{\odot}$. The CV configuration where M_2 is $> M_1$ might be not characterized by stable mass transfer (Warner, 2003).

Symbiotics as progenitors of SNe-Ia had some popularity in the 90's thanks to (Munari and Renzini, 1992), which showed that the density of progenitors of Symbiotic novae was vastly consistent with the rate of SN-Ia explosions. A few years later this scenario was questioned by radio observations of SN 1986G in Centaurus A (Boffi and Branch, 1995) that failed to reveal the radio synchrotron emission (Chevalier, 1988) that should have been produced during the interaction between the SN blast wave and the circumburst material. The impact of this negative result was partially softened by the fact that SN 1986G was somehow peculiar object and therefore may be it was not representative of the general behavior of whole SN-Ia class. The result from radio observations of 27 SNe-Ia obtained with the Very Large Array two decades later by Panagia *et al.* (2006) was much more sharp. There was not a single positive detection and the lack of radio signal implied the existence of a very low circumburst density. Thus Panagia *et al.* (2006) were able to set an upper limit of $\dot{M} \lesssim 3 \times 10^{-8} M_{\odot}/\text{yr}$ on the mass loss rate of the secondary star, which is 1-2 orders of magnitudes smaller than the \dot{M} expected from red giant donor. This upper limit was further constraint up to $\dot{M} \lesssim 5 \times 10^{-9} M_{\odot}/\text{yr}$ by Chomiuk *et al.* (2016).

The most direct pieces of observation, to date, which might rule out Symbiotics to form the main bulk of SN-Ia progenitors are the deep pre-explosion images, obtained with HST, of the site of the "standard" SN-Ia 2011fe. Li *et al.* (2011) did not find a suitable candidate progenitor at the SN location and by taking advantage of the proximity of the M101 host galaxy, they could set a firm upper limit on the magnitude of the progenitor system. They could conclude that the progenitor was: "10 ÷ 100 times fainter than previous limits on other SN-Ia progenitors. This directly rules out luminous red giants and the vast majority of helium stars as the mass donating companion. These observations favour a scenario where the exploding WD of SN 2011fe accretes matter either from another WD, or by Roche-lobe overflow from a subgiant or main-sequence companion star".

Finally we point out the vain search for a surviving companion in the SN-Ia remnant SNR 0509-67.5 by Pagnotta *et al.* (2014). These authors could concluded that: "This lack of any ex-companion star to deep limits ($M_V = 8.4 \text{ mag}$) rules out all published single-degenerate models for this supernova. The only remaining

possibility is that the progenitor of this particular type Ia supernova was a double-degenerate system."

Then the question becomes whether or not some room is left for the existence of the single degenerate scenario or is it completely ruled out?

Probably the most convincing piece of evidence that a single degenerate system can explode as SN-Ia is given by the detection of circumburst material around SN-Ia 2006X (Patat *et al.*, 2007a). The analysis of the evolution in the spectroscopic features of NaI-D doublet induced by the change of the ionization conditions of the SN radiation field have proven that this material was located at distances from the SN of about $D \sim 10^{16}$ cm and it was expelled by the Supernova progenitor about 50 yr before the explosion. The existence and the fast spectroscopic evolution of this circumburst material is reminiscent of the THEA absorptions observed by Williams *et al.* (2008), around novae shortly after the maximum light. According to Patat *et al.* (2007a) these erratic or recurrent in time emission episodes rule out a double degenerate scenario progenitor for SN 2006X. In passing we also note that a similar study carried out by (almost) the same team (Patat *et al.*, 2007b) on SN-Ia 2000cx was not able to detect a behavior similar to that exhibited by SN 2006X. Taking at their face values these results may suggest that both scenarios are at play. If this fact is true a second question arises: what is the nature of the single degenerate systems? According to general believe some of these systems might be Recurrent Novae e.g (Kato and Hachisu, 2012). Out of about 400 nova outbursts in the Galaxy, Fig. 5, only ten of them has been identified as RNe. The rigorous correction of these numbers for incompleteness is very questionable procedure because many assumptions are involved. However we can apply at first order approximation the following argument. From Sect. 3.5 we assume a galactic nova rate of 25 Novae/yr, therefore in the last 156 years (from U Sco in 1863) in the Galaxy have occurred about 4000 nova explosions. It means that, on average, the probability to detect in this period of time a nova outburst was about 10%. Since to identify a Recurrent Nova we need to observe at least two outbursts from the same system, the probability to have observed a RN is at first order only $\sim 1\%$. Since with observed in the MW about a dozen of RNe, their actual number has to be ~ 10 times as larger. This simple order of magnitude calculation would indicate a "true" ratio RNe/CNe of $\sim 30\%$ for our Galaxy cf. (Pagnotta *et al.*, 2014). We note that Della Valle and Livio (1996) have estimated for M31 and LMC a fraction RNe/CNe of 10% and 30% respectively and they concluded that "*RN-type systems can account for at most a few percent of the SN-Ia rate*". Shafter *et al.* (2015) consistently estimated that "*as many as one in three nova eruptions observed in M31 arise from progenitor systems having recurrence times < 100 yr...*" and they concluded that it is very unlikely that RNe could play a major role as SNe-Ia progenitors in a galaxy like M31.

These last results about the contribute of RNe to the SNe-Ia rates seem to relegate the SD to a completely secondary role with respect to double degenerate. However this story does not seem to be over yet. Della Valle and Livio (1994) and Mannucci *et al.* (2005) have found that the SNe-Ia rates, per unit of luminosity of the respective host galaxies, are a full order of magnitude higher in late type/*blue* galaxies than observed in E-S0/*red* galaxies. This result can be explained by assuming that a significant fraction of Ia events in late spirals/irregulars originates in a relatively young stellar component characterized by evolutionary times shorter than ~ 100 Myr (Mannucci *et al.*, 2006), while the SNe-Ia in passive elliptical

galaxies have a time evolution which is described by a much wider distribution, well characterized by an exponential function with a decay time of about 3 Gyr. Recently (Ikiewicz *et al.*, 2019) have studied a subset of Symbiotic binaries, which contain a Mira donor that is characterized by orbital periods of dozens of years. Obviously these symbiotic stars are not normally considered suitable SN-Ia progenitors because of their large separation (however see Hachisu *et al.* 1999 for a discussion of the symbiotic scenario). This team has analysed the evolution of V407 Cyg and they find that there is 90% probability that this system can explode as SN-Ia and this probability increases to 97% for WD with $\sim 1.35 M_{\odot}$, which incidentally coincides with the value of the mass suggested by (Hachisu and Kato, 2012). Ikiewicz *et al.* (2019) predict that the white dwarf in V407 Cyg will reach the Chandrasekhar limit in 40-200 Myr and eventually will explode as SN-Ia. This evolutionary time is consistent with the "prompt" SN-Ia population identified by Mannucci *et al.* (2006) (see also Greggio 2010). Very recently Darnley *et al.* (2006, 2019) have provided convincing evidence that the progenitor of the yearly erupting RN M31N 2008-12a is a single degenerate system with a red giant donor (see also Williams 2016). Kato *et al.* (2015) have determined that the central WD is of CO type and that its mass is $\sim 1.38 M_{\odot}$. Moreover, the WD is quickly growing in mass, by ejecting only $\sim 40\%$ of the matter accreted during each nova outburst cycle (which corresponds to $\sim 10^{-7} M_{\odot}$). Therefore, it will pass the Chandrasekhar limit in about 100,000 years and will terminate its life as a SN-Ia eruption.

12 Multi-wavelength Novae

In the last fifteen years, the possibility to investigate nova outbursts in other frequencies than the optical and near-infrared electromagnetic range has become a standard procedure, thanks to a new-generation of instruments and detectors operating at radio, infrared (IR) and high-energies as X- and γ -rays. These observations have revealed that novae undergo a complex evolution characterized by several emission phases and a non-spherical geometry for the nova ejecta. In the following we briefly review some of the most interesting results obtained in recent years.

12.1 Novae in the Infrared

Abundances from novae can also be inferred from IR observations, in particular for Ne, Mg and Al (Gehrz *et al.*, 2008, 2015). IR observations have also demonstrated that CNe, particularly of the CO class, are dust grain factories (Geisel *et al.*, 1970; Hyland and Neugebauer, 1970; Clayton, 1979; Gehrz *et al.*, 1980a,b; Gehrz, 1990; Gehrz *et al.*, 1995a; Mason *et al.*, 1996, 1998; Woodward and Starrfield, 2011; Chesneau *et al.*, 2012). In these objects the free-free emission that dominates the optical peak is often followed by a dust-dominated phase, while in ONe novae an Orion phase immediately follows the Principal stage (Gehrz *et al.*, 1995b; Woodward *et al.*, 1995, 1997). NIR observations of novae have also provided convincing evidence for bi-polar ejection of nova shells (Chesneau *et al.*, 2007, 2011, 2012). For example NIR-interferometric observations of V339 Del (Schaefer *et al.*, 2014) obtained during the earliest phases have detected from the second day an

aspherical distribution of the light that can be explained by assuming prolate or bipolar expanding structures. We address the reader to the papers by Chesneau and Banerjee (2012) and by Gehrz *et al.* (2008) for a deep insight into IR emission from Novae.

12.2 X-ray Novae

Classical Novae have been detected in X-rays since the first observations by the ROSAT satellite in the 80's (Krautter *et al.*, 1996; Orio *et al.*, 2001). During the bright outburst phases, X-ray radiation is emitted as a consequence of three physical processes: 1) a thermal emission from the underlying WD, which has been observed in several novae (Krautter *et al.*, 1996; Mukai and Ishida, 2001; Balman & Krautter, 2001; Schwarz *et al.*, 2011; Page *et al.*, 2015; Osborne, 2015) when the nova ejecta starts to become optically thin, what is called also as supersoft X-ray source phase (SSSs); 2) a hard X-ray component, originated in the shocks between the ejecta with material surrounding the WD progenitor, likely to be previous ejected shells or a dense interstellar medium (Mukai and Ishida, 2001; Metzger *et al.*, 2015); 3) emission from the re-established accretion during the quiescent phase (Takei *et al.*, 2011). ROSAT observed the supersoft emission in only a few objects, among them v1974 Cyg one of the most extensively studied Galactic novae by ROSAT (Balman *et al.* (1998)). The breakthrough in this field occurred with the commissioning of the Neil Gehrels Swift satellite (Gehrels *et al.*, 2004). Thanks to its capability to re-point a target within minutes/hours, this satellite is particularly suitable for observations of target of opportunities such as nova stars. For example one of the first and most complete X-ray analysis of a nova outburst is the case of the 2006 outburst of the recurrent nova RS Oph (Osborne *et al.*, 2011). Today we have observed the nova emission in X-rays from the very earliest phases during the whole visibility stage. Among ~ 80 novae observed in X-rays, more than half have been detected in X-rays (Ness *et al.*, 2007; Osborne, 2015). Its sensitivity allowed us to extend the nova studies to objects occurring in M31 and M33 and MCs (Aydi *et al.*, 2018). The results of a systematic survey on Novae in the Milky Way and M31 with Swift has provided us with a number of important results: i) there exists an early phase characterised by hard X-ray emission due to the presence of shocks within the different components of nova ejecta (Chomiuk *et al.*, 2014b) and with pre-existent circum-burst medium; ii) the onset of the supersoft phase begins soon in novae characterised by a low t_2 value, and then a low ejected mass (Henze *et al.*, 2014); iii) the duration of the supersoft phase is an indicator of the WD mass, being inversely related to it (Hachisu and Kato, 2006, 2010); iv) the temperature of the thermal component observed during the supersoft phase, modeled with a simple blackbody function, is directly related with the mass of the underlying WD; v) X-ray photons rising from the supersoft phase can destroy the dust produced at late phases (Gehrz *et al.*, 2018), but see also Shore *et al.* (2018); vi) some novae show a modulation in X-rays, whose characteristic period varies from few dozens of seconds like RS Oph, KT Eri and V5668 Sgr (Osborne *et al.*, 2011; Ness *et al.*, 2015) up to few days like Nova LMC 2009A. We address the reader to the paper on Swift nova observations by Osborne (2015).

In the last years, 13 novae belonging to the Milky Way (Orio *et al.*, 2005; Mukai *et al.*, 2008), M31 (Orio, 2006; Orio *et al.*, 2010; Henze *et al.*, 2010, 2011,




Fig. 55 The optical emission of V1369 Cen (lower panel) compared with the emission detected by the Fermi-LAT detector (upper panel, courtesy M. Kovacevic). Grey regions mark the time intervals corresponding to a positive detection of Fermi-LAT.

2012, 2014) and SMC (Orio *et al.*, 2018) have been monitored with XMM-Newton and Chandra. The main goal was the study of supersoft X-ray sources at high-resolution, aimed at both determining the duration of their SSS phase and the physical parameters of the observed atmosphere, including abundance studies (Ness *et al.*, 2009; Pinto *et al.*, 2012). Some theoretical implications have been discussed by (Hernanz and Sala, 2010). During these surveys was possible to discover several RNe (Henze *et al.*, 2018) and remarkably enough some very rare events such as a nova in outburst in a globular cluster of M31 (Henze *et al.*, 2013).

12.3 Gamma-ray Novae

In 2014, Classical Novae, were definitely revealed as a new class of gamma-ray sources (Ackermann *et al.*, 2014), although the first nova-like observed in gamma-rays was the symbiotic nova V407 Cyg (Abdo *et al.*, 2010). To date about a dozen of novae and symbiotic systems have been detected in gamma-rays at energy larger than 100 MeV, but below few GeV. The physical origin of this emission is not yet clear, but it is likely related to strong shocks driven by both the dense

nova ejecta when colliding with the tenuous interstellar medium or the wind from the secondary (Martin and Dubus, 2013). As an alternative the internal shocks are produced by the interaction of a fast wind radiatively driven by the nuclear burnings on the white dwarf (Martin *et al.*, 2018) with inhomogeneities formed into the expanding shells of the nova ejecta (see the case of v1369 Cen Mason *et al.* 2018). However, we cannot exclude that a combination of these two mechanisms is at play. Although these inhomogeneities are often observed as knots in some historical Novae, such as T Pyx (Shara *et al.*, 2015), only $\sim 15\%$ of the objects optically discovered show gamma emission. This rises the question whether or not gamma-ray novae are intrinsically rare among galactic novae or this small percentage results from a distance bias (Morris *et al.*, 2017; Franckowiak *et al.*, 2018), which prevents the discovery of high energy emission in most CNe because of their distances.

The high-energy radiation observed in novae does not generally show a direct link with the optical emission (Franckowiak *et al.*, 2018) with the exception of bright LAT novae such as: ASAS-SN 16ma (Li *et al.*, 2017), TCP J1734475-240942 (Munari *et al.*, 2017), V906 Car (Aydi *et al.*, 2020) and v1369 Cen (see Fig. 55) where a correlation between the observed optical light curve and the γ -ray flux was observed, then implying that a considerable fraction of the optical radiation comes from the re-processing of the gamma-ray radiation at longer wavelengths. This evidence also suggests that the optical radiation observed in the nova phenomenon is not only originating from the thermal energy released in the TNR (Metzger *et al.*, 2015), but it is partly due to the re-processing of high-energy radiation coming from acceleration of particles likely originating by magnetic field amplification in the shocks (Li *et al.*, 2017) or radiation processes in leptonic and/or hadronic scenarios (Vurm and Metzger, 2018).

After connecting the nova high-energy radiation with the multiple absorption of the principal spectra (see sect. 4), the modelling of nearby nova remnants (Slavin *et al.*, 1995; Bode, 2002; Moraes and Diaz, 2009; Ribeiro *et al.*, 2009, 2013b,a), IR interferometry (Chesneau and Banerjee, 2012; Schaefer *et al.*, 2014), and the delayed emission from radio observations (see next subsection) a more clear scenario of nova phenomenon emerges. In this framework immediately after the ignition of the TNR, a first component with an oblated/prolate matter geometry distribution expands with velocities of few hundreds of km/s. At later times, a faster ($\sim 1,000$ - $5,000$ km/s) wind, is emitted from the poles of the central WD, with relatively high opening angles, and then it catches up with the dense earlier component. The interaction between these two ejecta can lead to particle acceleration that give origin to the observed gamma-ray radiation. The resulting structure of the remnant will show years after the outburst the consequences of the interaction, as for example observed in FH Ser 1970 an ellipsoid structure characterised by denser ring-like component along the minor axis (Della Valle *et al.*, 1997; Gill and O’Brien, 2000).

Novae are also factories of isotopes, like ^7Be and ^{22}Na , which decay via electron capture or through the β^+ -process emitting very high-energy photons at specific energies (see Hernanz 2012 for a review). For example, ^7Be decays into ^7Li emitting a photon with an energy of 478 keV. The recent discovery of a consistent amount of beryllium produced in novae implies that we should be able to detect this line with space-based detectors like INTEGRAL. However, a detection of this emission line is still missing, likely due to the large distance of the novae for which beryllium




Fig. 56 The European VLBI Network (EVN) radio image of the remnant of the V959 Mon as obtained on September 18, 2012. The bi-polar shape of the remnant is clearly shown by the darker blobs, where the red circles mark the centered position on September 18 while the green circles mark the observed positions ~ 22 days later (Chomiuk *et al.*, 2014a).

and/or lithium have been observed (see e.g. Siegert *et al.*, 2018). The detection of this 478 keV line would represent the conclusive confirmation that novae are one of the main factories of lithium in the Milky Way.

12.4 Novae at Radio frequencies

One of the most efficient methods to measure the mass ejected in Novae consists in the analysis of radio emission emitted by the thermal bremsstrahlung free-free emission, in particular for the case of an impulsive ballistic shell ejection (see e.g. Seaquist and Palimaka, 1977, and Seaquist & Bode in Bode and Evans 2008). Novae have been observed in radio since the 70's, with FH Ser being one of the first studied case (Hjellming and Wade, 1970; Wade and Hjellming, 1971), where Hjellming *et al.* (1979) succeeded in reproducing the phenomenology observed in radio with a spherical shell linear gradient model characterised by only three variable parameters. Other novae, like HR Del and V1500 Cyg can be very well explained by this ionized shell model (Ennis *et al.*, 1977). Spatially-resolved radio observations have also allowed to resolve the ejecta which are characterised by a non-spherical emission and in several cases by a bi-polar geometry and lobes (Taylor *et al.*, 1988; Hjellming, 1995). The first nova that was resolved in radio was V1974 Cyg (Eyres *et al.*, 1996), which showed the presence of an evolving remnant only ~ 80 days after the outburst. At these epochs, radio images already showed deviation from spherical symmetry. These observations have contributed to improve the theoretical models that were used to reproduce the observed shape of the nova remnants (Lloyd *et al.*, 1996).

Recent observations show a more complex structure. While a non-spherical geometry characterized by bi-polar ejecta, is observed in almost all cases ((Rupen *et al.*, 2008; Sokoloski *et al.*, 2008; Chomiuk *et al.*, 2014a), see also Fig. 56) with some exceptions like V351 Pup (Wendeln *et al.*, 2017), additional pieces of

the jigsaw have been discovered. First, the presence of a non-thermal emission in few novae like RS Oph (Rupen *et al.*, 2008) v1723 Aql (Weston *et al.*, 2016a) and v5589 Sgr (Weston *et al.*, 2016b) and in the gamma-ray nova V1324 Sco (Finzell *et al.*, 2018). These observations were explained as synchrotron radiation rising from shocks originating in the interaction between a faster wind with a oblated/toroidal slower components (Vlasov *et al.* 2016. Another interesting result consists in the presence of a delayed (with respect to the initial optical outburst) event of mass ejection observed in the recurrent nova T Pyx (Chomiuk *et al.*, 2014b), characterised by an amount of mass larger than expected from recurrent novae (Nelson *et al.*, 2014). This result may suggest that the impulsive ejection model of novae needs some adjustments. On the other side it confirms that radio observations are more efficient in measuring the masses of the nova ejecta than optical observations (e.g. the case of v458 Vul (Roy *et al.*, 2012b)). We address the reader to (Roy *et al.*, 2012a) for a paper which summarize the radio studies on novae. The future for radio studies on nova stars is very promising in view of the new facilities that will be come operational in the next decades such as: SKA (O’Brien *et al.*, 2015) or the next generation Very LArge Array (ngVLA Linford *et al.*, 2018).

13 Conclusions

The new distances to galactic novae provided by GAIA DR2 have allowed us to re-analyze the numerous collections of data for novae in our Galaxy and in external systems. From our analysis the following results emerge:

- 1) The analysis of the nova counts, their heights above the galactic plane and rates of decline of novae inside the Milky Way has allowed to derive, on more robust empirical basis, the notion of disk and bulge nova populations. Fig. 14 shows strong evidence ($> 3\sigma$) for the existence of a correlation between the rate of decline and the height above the galactic plane. This trend indicates that the fastest nova systems, which contain the most massive WDs, are concentrated close to the galactic plane. We then derive that the typical *disk nova* is a fast evolving object whose lightcurve exhibits a bright peak at maximum $M_V \sim -9$ ($t_2 \lesssim 13^d$ or $t_3 \lesssim 20^d$) a smooth early decline and preferentially belongs to the He/N (or Fe II broad) spectroscopic class. The progenitor is preferentially located at small heights above the galactic plane ($\lesssim 150\text{pc}$) and the associated WD is rather massive, $M_{WD} \gtrsim 1 M_\odot$. The typical *bulge/thick disk nova* is a slow evolving object whose lightcurve exhibits a fainter peak at maximum, $M_V \sim -7.2$ mag ($t_2 \gtrsim 13^d$ or $t_3 \gtrsim 20^d$), often multiple maxima, dust formation, maximum standstill and belongs to the Fe II spectroscopic class. The progenitors extend up to 500 pc and beyond from the galactic plane and are likely to be related to a Pop II stellar population of the galactic thick-disk/bulge. Therefore they are associated (on average) with less massive WDs, $M_{WD} \lesssim 1 M_\odot$. The characteristic recurrence times between outbursts for nova systems formed by massive WDs are $\Delta T_{rec} \lesssim 10^{3\div 4}$ years and their duty cycle lasts \approx a few $\times 10^7$ years. For nova systems formed by light WDs $\Delta T_{rec} \lesssim 10^{5\div 6}$ years and their lifetime activity lasts $\approx \times 10^9$ years.

- 2) Analysis of the MMRD relationship for LMC, M31 and Virgo novae confirm the existence of systematic differences in the distributions of the rates of decline of

the respective nova populations. At least 70% of novae in the LMC are bright and fast and therefore associated with massive WDs while fast novae in M31 are $\lesssim 25\%$. The two distributions are significantly different (K-S gives $\gtrsim 99\%$). The analysis of the spatial distributions suggests that nova populations in M31 and M81 are a mixture of disk and bulge novae, $\sim 30\% - 70\%$ and $40\% - 60\%$ respectively. Novae in the LMC and M33 originate from disk population whereas novae in M87 and NGC 1316 are obviously related to an old bulge stellar population.

3) Analysis of the distributions of the rates of decline as a function of the absolute magnitude at maximum of LMC, M31 and Virgo, reveals the existence of a scant group of deviating objects. This sub-sample includes super-bright novae which extra energy input might be due to dissipation of orbital energy motion of the secondary engulfed in the nova ejecta during the nova common envelope stage. Also there are objects which fall below the 3σ strip. They can well be RNe that eject very light envelopes (10^{-7} - $10^{-8} M_{\odot}$), hundreds or thousands times lighter than envelopes associated with CNe or patchy absorbed novae. Outliers of the M31 MMRD found by Kasliwal *et al.* (2011) might well be classified into the previous classes of deviating objects without calling for the existence of a sub-class of faint and fast CNe that at the moment appear only predicted. Cao *et al.* (2012) observations of novae in M31 are fully consistent with the MMRD relation. If we consider novae observed in the nearest galaxies, such as M31 and LMC, we have a total number of 144 data points (see Tables 13, 14, 15, 16), 12 of which (8%) are obvious outliers of the MMRD at 3σ level (see Fig. 45 and Fig 47).

4) Comparison between the distances obtained for galactic novae with the MMRD relation and GAIA DR2 agree to better (or much better) than 30% then confirming that the MMRD relation can be used to estimate the distance to novae in the Milky Way with an interesting accuracy, contrary to recent claims. Most of the error (still) associated to distance measurements obtained via MMRD comes from the uncertainty in estimating: i) the absorption A_V toward each nova; ii) the correct computation of the MMRD parameters, such as the observed magnitude at maximum and the rate of decline; iii) misuse of the MMRD to intrinsically faint objects such as RNe.

5) GAIA DR2 distances have allowed a new absolute calibration of the MMRD relation. The comparison between the absolute and observed MMRD relations of M31, LMC and Virgo finds: $(V - M_v)_{M31} = 24.35 \pm 0.31$ mag, $(V - M_v)_{LMC} = 18.44 \pm 0.41$ mag, and $(V - M_v)_{Virgo} = 31.32 \pm 0.44$ mag, which are in excellent agreement with the current distance measurements for Local Group and Virgo galaxies, respectively. Obviously, given the intrinsic dispersion of the MMRD it appears hard to measure the distances to extragalactic systems to better than $\sim 10\% - 20\%$ which make novae not longer competitive distance indicators in the era of "high precision" cosmology (Pietrzyński *et al.*, 2019; Riess *et al.*, 2019). This situation could improve in the next years if both the sample of "calibrators" (that is the number of novae inside the Milky Way with GAIA DR2 distance and characterized by good measurement of A_V) and the samples of novae discovered in external galaxies with reliable measurements of the MMRD parameters, will significantly increase.

6) The nova rate in the Milky Way is currently known within a factor of two: 20-40 Novae/yr. Recent new measurements ranging between a bulge rate of 13.8 ± 2.6

novae/yr (Mróz *et al.*, 2015) and a global rate 50 novae/yr (Shafter, 2017) do not help to clarify this discrepancy. On the other hand we have shown that novae in the Milky Way and M31 exhibit similar properties such as the distributions of the rates of decline and the spectroscopic classification (most MW and M31 novae belong to Fe II type). Therefore if most Galactic novae are produced in the bulge, it is unlikely that the global nova rate of the Milky Way might be significantly larger than it was measured by Mróz *et al.* (2016). So, a rate of $R_{MW} = 25^{+15}_{-5}$ /yr seems an appropriate educated guess.

7) A few sporadic observations have allowed to provide an estimate of the nova rate in Globular Clusters systems. An analysis of the present data finds a rate ranging between: a few $\times 10^{-4}$ and 10^{-2} novae $\text{GC}^{-1}\text{yr}^{-1}$.

8) While there is a substantial agreement on LMC, SMC and M33 rates, we find a factor two discrepancy for M31, $R_{M31} = 30 - 60/\text{yr}$ and a full factor three uncertainty for the giant elliptical galaxies in Virgo cluster $R_{Virgo} = 100 - 360/\text{yr}$. A simple explanation is that old nova survey suffer of severe incompleteness. For example the rather sparse surveys carried out by Rosino in the 70's on M31 might suffer of such observational bias. Despite the control time has been duly estimated and the correction for field incompleteness have been taken into account, a factor $\sim two$ may be still hidden in fragmented and not homogeneous material collected over 25 years. Much more difficult is to explain a factor two incompleteness in the Arp (1956) data. From his paper we can read some impressive numbers

"Nearly 1000 plates were taken on 290 nights between June 1953 and January 1955. Observations were as continuous as possible..... Consequently of 30 novae discovered for only 5 were the maxima missed by more than one day."

This summary doesn't leave too much room for having missing one nova out of two, unless 50% of novae in M31 are completely extinguished by absorption. We have adopted a rate of $R_{M31} = 40^{+20}_{-10}$ novae/yr.

As far as concern nova observations in Virgo ellipticals the most straight explanation is that nova surveys from ground based telescopes suffer of severe incompleteness. However this hypothesis seems not supported by Ferrarese *et al.* (2003) survey on M49 which was carried out with HST. They found a rate of $\sim 70 \div 135$ novae/yr. Another possibility discussed in this paper is that the nova rate of M87 might be enhanced by the jet. The rate of this galaxy remains uncertain, likely in the range 100-200 novae/yr.

The mismatch among the nova rates represents a good rationale for undertaking new dedicated surveys on novae in Local Group galaxies and beyond. Particularly new surveys in MCs, M33 and M31 characterized by a cadence of few hours (rather than a few days) would be capable to discover the possible sub-class of ultra-fast novae predicted on theoretical grounds (e.g. (Yaron *et al.*, 2005)). There is no reason to look for these intrinsically faint transients in distant galaxies. The use of telescopes characterized by large fields of view will help to minimize the number of correction factors that are normally applied to the observed rates.

9) Whether or not the nova rate depends on the Hubble type of parent galaxy is a point not yet fully clarified. One result about which all authors seem to converge




Fig. 57 The Nova "time consuming" index, that is expressed in terms of telescope time (computed in hours) needed to make the "average" nova lightcurve in extragalactic systems. It is plotted vs. the epochs of the observations.

is that the specific nova rate (i.e. the normalized nova rate per unit of luminosity of the host) exhibited by blue low luminosity class galaxies, such as M33, LMC and SMC, is higher by a factor about 3 than observed in giant Spirals and Ellipticals. This result can be interpreted in terms of "selection effect" due to the different recurrence times of the nova outbursts combined with the different density of nova progenitors belonging to different parent stellar populations.

10) High resolution spectroscopic observations of galactic and MCs novae have produced two interesting results. i) In 2008, Williams and collaborators have discovered the so called THEAs (Transient Heavy Element Absorptions), which are signatures of a toroidal component of material surrounding the primary WD that is observed at low velocities during the nova outburst. ii) Izzo *et al.* (2015) have observed for the first time ^7Li in the ejecta of V1369 Cen and in the same years, Tajitsu *et al.* (2015, 2016), Molaro *et al.* (2016) and Izzo *et al.* (2018) have identified also the presence of ^7Be , which decays in ^7Li with an half-life time decay of ~ 53 days, finally confirming 70's predictions that novae were Lithium factories.

11) Novae and SNe-Ia connection. Several pieces of evidence point out that RNe and symbiotic stars should not be the major class of progenitors of Type Ia supernovae. Particularly, the radio observations of two dozen SNe-Ia with no detections and pre-explosion images of the SN-Ia 2011fe prototype seem to exclude that stellar systems containing red giant donors are the main channel for SN-Ia production. Nevertheless the SD does not appear completely ruled out. Particularly, objects such as V407 Cyg and M31N 2008-12a that contain red giant donors seem to be promising candidates to explode as SNe-Ia. We tentatively conclude that the both channels are open, although DD seems to be able to provide the major contribution.

12) Nova survey in extragalactic systems, despite their scientific interest, have not been so popular among astronomers and even less among the various time allocation committees spread over the world (although some remarkable exceptions do exist). The main reason for this is the unpredictable nature of nova events, which made nova observations considerably time consuming and difficult to schedule. However two important facts may help to boost the interest for nova studies: i) the observing time requests at 2-4m class telescopes have decreased in the last 20 years by about a factor 5 (survey on ESO telescopes, Patat *et al.* (2019) but see also Patat and Hussain 2013). Telescope of these sizes are suitable to study the nova properties in Local Group galaxies and therefore the general decrease for their quest can play in favor of proposal dedicated to novae; ii) large field of view telescopes, such as LSST that will be come operational in the next few years, have the potential to change dramatically the future of nova studies. Fig 57 shows the trend of the "time consuming" index, that we have computed in terms of hours at the telescope needed to make the "average" nova lightcurve in extragalactic systems vs. the respective observing epochs. Obviously, this parameter depends on many factors (e.g distance to the galaxy, size and field of view of the telescope, seeing, photographic plates or CCDs, ground-based vs. space instrumentation, intrinsic rate of events...), however we can recognize a general trend. From the 20-40 hours typical of the surveys carried out during the photographic era on nearby galaxies, we drop to ~ 8 hours per lightcurve with 4-meter class telescope + CCD for galaxies in Virgo cluster and to 1.5h/lightcurve with 8-meter telescope + CCDs for a survey in giant ellipticals. This index has further decreased with the use of HST to less than 1h per lightcurve. In principle the study of extragalactic novae with future large size and large fields telescope such as LSST will dramatically decrease this "index" to a few minutes/lightcurve. It will be essentially limited by observing overheads (e.g. slewing telescope to target, CCD readout...) and the intrinsic frequency of nova outbursts of the observed galaxies.

Acknowledgements The authors thank the referees for their constructive criticisms that have significantly improved the presentation and discussion of the data. We are also very grateful to Bob Williams for giving us valuable comments on the manuscript and to Roberto Gilmozzi for helpful discussions. MDV is deeply grateful to Prof. Leonida Rosino for having introduced him to the realm of Nova physics and to Hilmar Duerbeck for the many nights shared at the La Silla telescopes. MDV thanks ESO-Garching, where this manuscript was drafted for the creative and friendly atmosphere. LI was supported by grants from VILLUM FONDEN (project number 16599 and 25501). LI also acknowledges support from funding associated with Juan de la Cierva Incorporacion fellowship IJCI-2016-30940.

References

- Abbott, B. P., Abbott, R., and Abbott, T. D. et al. (2017). Multi-messenger Observations of a Binary Neutron Star Merger. *ApJ*, **848**(2), L12.
- Abdo, A. A., Ackermann, M., Ajello, M., Atwood, W. B., Baldini, L., Ballet, J., Barbiellini, G., Bastieri, D., Bechtol, K., Bellazzini, R., Berenji, B., Blandford, R. D., Bloom, E. D., Bonamente, E., Borgland, A. W., Bouvier, A., Brandt, T. J., Bregeon, J., Brez, A., Brigida, M., Bruel, P., Buehler, R., Burnett, T. H., Buson, S., Caliandro, G. A., Cameron, R. A., Caraveo, P. A., Carrigan, S., Casandjian, J. M., Cecchi, C., Celik, O., Charles, E., Chaty, S., Chekhtman, A., Cheung, C. C., Chiang, J., Ciprini, S., Claus, R., Cohen-Tanugi, J., Conrad, J., Corbel, S., Corbet, R., DeCesar, M. E., den Hartog, P. R., Dermer, C. D., de Palma, F., Digel, S. W., Donato, D., do Couto e Silva, E., Drell, P. S., Dubois, R., Dubus, G., Dumora, D., Favuzzi, C., Fegan, S. J., Ferrara, E. C., Fortin, P., Frailis, M., Fuhrmann, L., Fukazawa, Y., Funk, S., Fusco, P., Gargano, F., Gasparrini, D., Gehrels, N., Germani, S., Giglietto, N., Giordano, F., Giroletti, M., Glanzman, T., Godfrey, G., Grenier, I. A., Grondin, M. H., Grove, J. E., Guiriec, S., Hadasch, D., Harding, A. K., Hayashida, M., Hays, E., Healey, S. E., Hill, A. B., Horan, D., Hughes, R. E., Itoh, R., Jean, P., Jóhannesson, G., Johnson, A. S., Johnson, R. P., Johnson, T. J., Johnson, W. N., Kamae, T., Katagiri, H., Kataoka, J., Kerr, M., Knödlseeder, J., Koering, E., Kuss, M., Lande, J., Latronico, L., Lee, S. H., Lemoine-Goumard, M., Garde, M. L., Longo, F., Loparco, F., Lott, B., Lovellette, M. N., Lubrano, P., Makeev, A., Mazziotta, M. N., McConville, W., McEnery, J. E., Mehault, J., Michelson, P. F., Mizuno, T., Moiseev, A. A., Monte, C., Monzani, M. E., Morselli, A., Moskalenko, I. V., Murgia, S., Nakamori, T., Naumann-Godo, M., Nestoras, I., Nolan, P. L., Norris, J. P., Nuss, E., Ohno, M., Ohsugi, T., Okumura, A., Omodei, N., Orlando, E., Ormes, J. F., Ozaki, M., Paneque, D., Panetta, J. H., Parent, D., Pelassa, V., Pepe, M., Pesce-Rollins, M., Piron, F., Porter, T. A., Rainó, S., Rando, R., Ray, P. S., Razzano, M., Razzaque, S., Rea, N., Reimer, A., Reimer, O., Reposeur, T., Ripken, J., Ritz, S., Romani, R. W., Roth, M., Sadrozinski, H. F., Sander, A., Parkinson, P. M., Scargle, J. D., Schinzel, F. K., Sgrò, C., Shaw, M. S., Siskind, E. J., Smith, D. A., Smith, P. D., Sokolovsky, K. V., Spandre, G., Spinelli, P., Stawarz, L., Strickman, M. S., Suson, D. J., Takahashi, H., Takahashi, T., Tanaka, T., Tanaka, Y., Thayer, J. B., Thayer, J. G., Thompson, D. J., T. L., Torres, D. F., Tosti, G., Tramacere, A., Uchiyama, Y., Usher, T. L., Vandenbroucke, J., Vasileiou, V., Vilchez, N., Vitale, V., Waite, A. P., Wallace, E., Wang, P., Winer, B. L., Wolff, M. T., Wood, K. S., Yang, Z., Ylinen, T., Ziegler, M., Maehara, H., Nishiyama, K., Kabashima, F., Bach, U., Bower, G. C., Falcone, A., Forster, J. R., Henden, A., Kawabata, K. S., Koubsky, P., Mukai, K., Nelson, T., Oates, S. R., Sakimoto, K., Sasada, M., Shenavrin, V. I., Shore, S. N., Skinner, G. K., Sokoloski, J., Stroh, M., Tatarnikov, A. M., Uemura, M., Wahlgren, G. M., Yamanaka, M., and Fermi LAT Collaboration (2010). Gamma-Ray Emission Concurrent with the Nova in the Symbiotic Binary V407 Cygni. *Science*, **329**(5993), 817–821.
- Abia, C., Domínguez, I., Straniero, O., Limongi, M., Chieffi, A., and Isern, J. (2001). The Implications of the New $Z=0$ Stellar Models and Yields on the Early Metal Pollution of the Intergalactic Medium. *ApJ*, **557**(1), 126–136.
- Ackermann, M., Ajello, M., Albert, A., Baldini, L., Ballet, J., Barbiellini, G., Bastieri, D., Bellazzini, R., Bissaldi, E., Blandford, R. D., Bloom, E. D., Bottacini, E., Brandt, T. J., Bregeon, J., Bruel, P., Buehler, R., Buson, S., Caliandro, G. A., Cameron, R. A., Caragiulo, M., Caraveo, P. A., Cavazzuti, E., Charles, E., Chekhtman, A., Cheung, C. C., Chiang, J., Chiaro, G., Ciprini, S., Claus, R., Cohen-Tanugi, J., Conrad, J., Corbel, S., D’Ammando, F., de Angelis, A., den Hartog, P. R., de Palma, F., Dermer, C. D., Desiante, R., Digel, S. W., Di Venere, L., do Couto e Silva, E., Donato, D., Drell, P. S., Drlica-Wagner, A., Favuzzi, C., Ferrara, E. C., Focke, W. B., Franckowiak, A., Fuhrmann, L., Fukazawa, Y., Fusco, P., Gargano, F., Gasparrini, D., Germani, S., Giglietto, N., Giordano, F., Giroletti, M., Glanzman, T., Godfrey, G., Grenier, I. A., Grove, J. E., Guiriec, S., Hadasch, D., Harding, A. K., Hayashida, M., Hays, E., Hewitt, J. W., Hill, A. B., Hou, X., Jean, P., Jogler, T., Jóhannesson, G., Johnson, A. S., Johnson, W. N., Kerr, M., Knödlseeder, J., Kuss, M., Larsson, S., Latronico, L., Lemoine-Goumard, M., Longo, F., Loparco, F., Lott, B., Lovellette, M. N., Lubrano, P., Manfreda, A., Martin, P., Massaro, F., Mayer, M., Mazziotta, M. N., McEnery, J. E., Michelson, P. F., Mitthumsiri, W., Mizuno, T., Monzani, M. E., Morselli, A., Moskalenko, I. V., Murgia, S., Nemmen, R., Nuss, E., Ohsugi, T., Omodei, N., Orienti, M., Orlando, E., Ormes, J. F., Paneque, D., Panetta, J. H., Perkins, J. S., Pesce-Rollins, M., Piron, F., Pivato, G., Porter, T. A., Rainó, S., Rando, R., Razzano,

- M., Razzaque, S., Reimer, A., Reimer, O., Reposeur, T., Saz Parkinson, P. M., Schaal, M., Schulz, A., Sgrò, C., Siskind, E. J., Spandre, G., Spinelli, P., Stawarz, L., Suson, D. J., Takahashi, H., Tanaka, T., Thayer, J. G., Thayer, J. B., Thompson, D. J., Tibaldo, L., Tinivella, M., Torres, D. F., Tosti, G., Troja, E., Uchiyama, Y., Vianello, G., Winer, B. L., Wolff, M. T., Wood, D. L., Wood, K. S., Wood, M., Charbonnel, S., Corbet, R. H. D., De Gennaro Aquino, I., Edlin, J. P., Mason, E., Schwarz, G. J., Shore, S. N., Starrfield, S., Teyssier, F., and Fermi-LAT Collaboration (2014). Fermi establishes classical novae as a distinct class of gamma-ray sources. *Science*, **345**(6196), 554–558.
- Alis, S. and Saygac, A. T. (2014). *Investigating the Nova Rate Connection to Galaxy Type using Archival Hubble Space Telescope Images*, volume 490 of *Astronomical Society of the Pacific Conference Series*, page 95.
- Allen, C. W. (1954). Whole-sky statistics of celestial objects. *MNRAS*, **114**, 387.
- Allen, C. W. (1973). *Astrophysical quantities*.
- Amati, L. (2006). The $E_{p,i}$ - E_{iso} correlation in gamma-ray bursts: updated observational status, re-analysis and main implications. *MNRAS*, **372**(1), 233–245.
- Ambartsumian, V. A. (1952). *Teoreticheskaya astrofizika*.
- Arnould, M. and Norgaard, H. (1975). The Explosive Thermonuclear Formation of ${}^7\text{Li}$ and ${}^{11}\text{B}$. *A&A*, **42**, 55.
- Arp, H. C. (1956). Novae in the Andromeda nebula. *AJ*, **61**, 15–34.
- Aydi, E., Page, K. L., Kuin, N. P. M., Darnley, M. J., Walter, F. M., Mróz, P., Buckley, D. A. H., Mohamed, S., Whitelock, P., Woudt, P., Williams, S. C., Orio, M., Williams, R. E., Beardmore, A. P., Osborne, J. P., Kniazev, A., Ribeiro, V. A. R. M., Udalski, A., Strader, J., and Chomiuk, L. (2018). Multiwavelength observations of nova SMCN 2016-10a - one of the brightest novae ever observed. *MNRAS*, **474**(2), 2679–2705.
- Aydi, E., Chomiuk, L., Strader, J., Swihart, S. J., Bahramian, A., Harvey, E. J., Britt, C. T., Buckley, D. A. H., Chen, P., Dage, K., Darnley, M. J., Dong, S., Hamsch, F.-J., Holoien, T. W. S., Jha, S. W., Kochanek, C. S., Kuin, N. P. M., Li, K. L., Monard, L. A. G., Mukai, K., Page, K. L., Prieto, J. L., Richardson, N. D., Shappee, B. J., Shishkovsky, L., Sokolovsky, K. V., Stanek, K. Z., and Thompson, T. (2019). Flaring, Dust Formation, And Shocks In The Very Slow Nova ASASSN-17pf (LMCN 2017-11a). *arXiv e-prints*, page arXiv:1903.09232.
- Aydi, E., Sokolovsky, K. V., Chomiuk, L., Steinberg, E., Li, K. L., Vurm, I., Metzger, B. D., Strader, J., Mukai, K., Pejcha, O., Shen, K. J., Wade, G. A., Kuschnig, R., Moffat, A. F. J., Pablo, H., Pigulski, A., Popowicz, A., Weiss, W., Zwintz, K., Izzo, L., Pollard, K. R., Handler, G., Ryder, S. D., Filipović, M. D., Alsaberi, R. Z. E., Manojlović, P., De Oliveira, R., Walter, F. M., Vallely, P. J., Buckley, D. A. H., Brown, M. J. I., Harvey, E. J., Kawash, A., Kniazev, A., Kochanek, C. S., Linford, J., Mikolajewska, J., Molaro, P., Orio, M., Page, K. L., Sokoloski, J. (2020). Direct evidence for shock-powered optical emission in a nova. *Nag. Astron.*, **4**, 4.
- Baade, W. (1944). The Resolution of Messier 32, NGC 205, and the Central Region of the Andromeda Nebula. *ApJ*, **100**, 137.
- Baade, W. (1957). Galaxies and their stellar population. *The Observatory*, **77**, 165–171.
- Baade, W. (1958). The Population of the Galactic Nucleus and the Evidence for the Presence of an Old Population Pervading the Whole Disk of our Galaxy. *Ricerche Astronomiche*, **5**, 303.
- Baade, W. and Zwicky, F. (1934). On Super-novae. *Proceedings of the National Academy of Science*, **20**(5), 254–259.
- Bailer-Jones, C. A. L., Rybizki, J., Fouesneau, M., Mantelet, G., and Andrae, R. (2018). Estimating Distance from Parallaxes. IV. Distances to 1.33 Billion Stars in Gaia Data Release 2. *AJ*, **156**(2), 58.
- Balman, S. & Krautter, J. 2001, MNRAS, 326, 1441
- Balman, S., Krautter, J., & Ögelman, H. 1998, ApJ, 499, 395
- Bath, G. T. and Shaviv, G. (1978). The space density, recurrence rate and classification of novae. *MNRAS*, **183**, 515–522.
- Bianchini, A. (1987). Detection of solar-type cycles in cataclysmic variables. *Mem. Soc. Astr. It.*, **58**, 245–251.
- Bianchini, A., Hamzaoglu, E., and Sabbadin, F. (1981). The old-nova GK Per (1901). I. Determination of the orbital period. *A&A*, **99**, 392–393.
- Bode, M. F. (2002). The Evolution of Nova Remnants. In M. Hernanz and J. José, editors, *Classical Nova Explosions*, volume 637 of *American Institute of Physics Conference Series*,

- pages 497–508.
- Bode, M. F. and Evans, A. (2008). *Classical Novae*, volume 43.
- Boffi, F. R. and Branch, D. (1995). Radio Emission from Type IA Supernovae as a Test of Symbiotic Star Progenitor Systems. *PASP*, **107**, 347.
- Bondi, H. (1952). On spherically symmetrical accretion. *MNRAS*, **112**, 195.
- Bothun, G. D. (1992). The Bulge/Halo Luminosity of M33 and its Globular Cluster Population. *AJ*, **103**, 104.
- Bowen, I. S. (1947). Excitation by Line Coincidence. *PASP*, **59**(349), 196–198.
- Bowen, I. S. (1954). Edwin P. Hubble: 1889-1953. *Science*, **119**(3085), 204.
- Burstein, D. and Heiles, C. (1984). Reddening estimates for galaxies in the second reference catalog and the Uppsala General Catalog. *ApJS*, **54**, 33–79.
- Buscombe, W. and de Vaucouleurs, G. (1955). Novae in the Magellanic Clouds and in the Galaxy. *The Observatory*, **75**, 170–175.
- Cameron, A. G. W. and Fowler, W. A. (1971). Lithium and the s-PROCESS in Red-Giant Stars. *ApJ*, **164**, 111.
- Cao, Y., Kasliwal, M. M., Neill, J. D., Kulkarni, S. R., Lou, Y.-Q., Ben-Ami, S., Bloom, J. S., Cenko, S. B., Law, N. M., Nugent, P. E., Ofek, E. O., Poznanski, D., and Quimby, R. M. (2012). Classical Novae in Andromeda: Light Curves from the Palomar Transient Factory and GALEX. *ApJ*, **752**(2), 133.
- Capaccioli, M., Della Valle, M., D’Onofrio, M., and Rosino, L. (1989). Properties of the Nova Population of M31. *AJ*, **97**, 1622.
- Capaccioli, M., Della Valle, M., D’Onofrio, M., and Rosino, L. (1990). Distance of the Large Magellanic Cloud through the Maximum Magnitude versus Rate of Decline Relation for Novae. *ApJ*, **360**, 63.
- Cappellaro, E., Evans, R., and Turatto, M. (1999). A new determination of supernova rates and a comparison with indicators for galactic star formation. *A&A*, **351**, 459–466.
- Cappellaro, E., Botticella, M. T., Pignata, G., Grado, A., Greggio, L., Limatola, L., Vaccari, M., Baruffolo, A., Benetti, S., Bufano, F., Capaccioli, M., Cascone, E., Covone, G., De Cicco, D., Falocco, S., Della Valle, M., Jarvis, M., Marchetti, L., Napolitano, N. R., Paolillo, M., Pastorello, A., Radovich, M., Schipani, P., Spiro, S., Tomasella, L., and Turatto, M. (2015). Supernova rates from the SUDARE VST-OmegaCAM search. I. Rates per unit volume. *A&A*, **584**, A62.
- Cassatella, A., Benvenuti, P., Clavel, J., Heck, A., Penston, M., Selvelli, P. L., and Macchetto, F. (1979). On the ultraviolet spectrum of Nova Cygni 1978. *A&A*, **74**, L18–L19.
- Cassatella, A., Altamore, A., and González-Riestra, R. (2002). Classical novae in outburst: The early evolution of the ultraviolet continuum. *A&A*, **384**, 1023–1029.
- Cassatella, A., Lamers, H. J. G. L. M., Rossi, C., Altamore, A., and González-Riestra, R. (2004). A study of the expanding envelope of Nova V1974 Cyg 1992 based on IUE high resolution spectroscopy. *A&A*, **420**, 571–588.
- Cecchini, G. and Gratton, L. (1942). *Le stelle nuove*.
- Charbonnel, C. and Primas, F. (2005). The lithium content of the Galactic Halo stars. *A&A*, **442**(3), 961–992.
- Chen, H.-L., Woods, T. E., Yungelson, L. R., Gilfanov, M., and Han, Z. (2016). Modelling nova populations in galaxies. *MNRAS*, **458**(3), 2916–2927.
- Chesneau, O. and Banerjee, D. P. K. (2012). Interferometric studies of novae in the infrared. *Bulletin of the Astronomical Society of India*, **40**, 267.
- Chesneau, O., Nardetto, N., Millour, F., Hummel, C., Domiciano de Souza, A., Bonneau, D., Vannier, M., Rantakyro, F., Spang, A., Malbet, F., Mourard, D., Bode, M. F., O’Brien, T. J., Skinner, G., Petrov, R. G., Stee, P., Tatulli, E., and Vakili, F. (2007). AMBER/VLTI interferometric observations of the recurrent Nova RS Ophiuchii 5.5 days after outburst. *A&A*, **464**(1), 119–126.
- Chesneau, O., Meilland, A., Banerjee, D. P. K., Le Bouquin, J. B., McAlister, H., Millour, F., Ridgway, S. T., Spang, A., ten Brummelaar, T., Wittkowski, M., Ashok, N. M., Benisty, M., Berger, J. P., Boyajian, T., Farrington, C., Goldfinger, P. J., Merand, A., Nardetto, N., Petrov, R., Rivinius, T., Schaefer, G., Touhami, Y., and Zins, G. (2011). The 2011 outburst of the recurrent nova α ASTROBJ γ T Pyxidis/ASTROBJ γ . Evidence for a face-on bipolar ejection. *A&A*, **534**, L11.
- Chesneau, O., Lagadec, E., Otulakowska-Hypka, M., Banerjee, D. P. K., Woodward, C. E., Harvey, E., Spang, A., Kervella, P., Millour, F., Nardetto, N., Ashok, N. M., Barlow, M. J., Bode, M., Evans, A., Lynch, D. K., O’Brien, T. J., Rudy, R. J., and Russell, R. W. (2012).

- The expanding dusty bipolar nebula around the nova V1280 Scorpi. *A&A*, **545**, A63.
- Cheung, C. C., Jean, P., Shore, S. N., Stawarz, L., Corbet, R. H. D., Knödlseider, J., Starrfield, S., Wood, D. L., Desiante, R., Longo, F., Pivato, G., and Wood, K. S. (2016). Fermi-LAT Gamma-Ray Detections of Classical Novae V1369 Centauri 2013 and V5668 Sagittarii 2015. *ApJ*, **826**(2), 142.
- Chevalier, R. A. (1988). The interaction of supernova 1987A with dense circumstellar gas. *Nature*, **332**(6164), 514–516.
- Chomiuk, L., Linford, J. D., Yang, J., O’Brien, T. J., Paragi, Z., Mioduszewski, A. J., Beswick, R. J., Cheung, C. C., Mukai, K., Nelson, T., Ribeiro, V. A. R. M., Rupen, M. P., Sokoloski, J. L., Weston, J., Zheng, Y., Bode, M. F., Eyres, S., Roy, N., and Taylor, G. B. (2014a). Binary orbits as the driver of γ -ray emission and mass ejection in classical novae. *Nature*, **514**(7522), 339–342.
- Chomiuk, L., Nelson, T., Mukai, K., Sokoloski, J. L., Rupen, M. P., Page, K. L., Osborne, J. P., Kuulkers, E., Mioduszewski, A. J., Roy, N., Weston, J., and Krauss, M. I. (2014b). The 2011 Outburst of Recurrent Nova T Pyx: X-ray Observations Expose the White Dwarf Mass and Ejection Dynamics. *ApJ*, **788**(2), 130.
- Chomiuk, L., Soderberg, A. M., Chevalier, R. A., Bruzewski, S., Foley, R. J., Parrent, J., Strader, J., Badenes, C., Fransson, C., Kamble, A., Margutti, R., Rupen, M. P., and Simon, J. D. (2016). A Deep Search for Prompt Radio Emission from Thermonuclear Supernovae with the Very Large Array. *ApJ*, **821**(2), 119.
- Ciardullo, R., Ford, H. C., Neill, J. D., Jacoby, G. H., and Shafter, A. W. (1987). The Spatial Distribution and Population of Novae in M31. *ApJ*, **318**, 520.
- Ciardullo, R., Ford, H. C., Williams, R. E., Tamblyn, P., and Jacoby, G. H. (1990). The Nova Rate in the Elliptical Component of NGC 5128. *AJ*, **99**, 1079.
- Ciatti, F. and Rosino, L. (1974). Photographic and spectroscopic observations of N Aql 1970, N Cyg 1970, and N Sct 1970. *A&AS*, **16**, 305–322.
- Cieslinski, D., Diaz, M. P., Mennickent, R. E., and Pietrzyński, G. (2003). Identification of New Eruptive Cataclysmic Variables toward the Galactic Bulge and Magellanic Clouds Using the OGLE-II Database. *PASP*, **115**(804), 193–211.
- Clayton, D. D. (1979). Sudden Grain Nucleation and Growth in Supernova and Nova Ejecta. *Ap&SS*, **65**(1), 179–189.
- Coc, A., Uzan, J.-P., and Vangioni, E. (2013). Standard Big-Bang Nucleosynthesis after Planck. *arXiv e-prints*, page arXiv:1307.6955.
- Coc, A., Uzan, J.-P., and Vangioni, E. (2014). Standard big bang nucleosynthesis and primordial CNO abundances after Planck. *JCAP*, **10**, 50.
- Coelho, E. A., Shafter, A. W., and Misselt, K. A. (2008). The Rate and Spatial Distribution of Novae in M101 (NGC 5457). *ApJ*, **686**(2), 1261–1268.
- Cohen, J. G. (1985). Nova shells. II. Calibration of the distance scale using novae. *ApJ*, **292**, 90–103.
- Cohen, J. G. and Rosenthal, A. J. (1983). Nova shells. *ApJ*, **268**, 689–697.
- Conti, P. S. (1968). Lithium Destruction and Rotational Braking. *ApJ*, **152**, 657.
- Cummings, J. D., Kalirai, J. S., Tremblay, P. E., Ramirez-Ruiz, E., and Choi, J. (2018). The White Dwarf Initial-Final Mass Relation for Progenitor Stars from 0.85 to 7.5 M_{\odot} . *ApJ*, **866**(1), 21.
- Curtin, C., Shafter, A. W., Pritchett, C. J., Neill, J. D., Kundu, A., and Maccarone, T. J. (2015). Exploring the Role of Globular Cluster Specific Frequency on the Nova Rates in Three Virgo Elliptical Galaxies. *ApJ*, **811**(1), 34.
- Curtis, H. D. (1917a). Finding List for General Catalog Numbers. *PASP*, **29**(170), 180–180.
- Curtis, H. D. (1917b). Novae in the Spiral Nebulae and the Island Universe Theory. *PASP*, **29**(171), 206–207.
- Curtis, H. D. (1917c). Two Novae in N.G.C. 4321. *Harvard College Observatory Bulletin*, **642**, 1–1.
- D’Antona, F. and Matteucci, F. (1991). Galactic evolution of lithium. *A&A*, **248**, 62.
- Darnley, M. J., Bode, M. F., Kerins, E., Newsam, A. M., An, J., Baillon, P., Novati, S. C., Carr, B. J., Cr    , M., Evans, N. W., Giraud-H    ud, Y., Gould, A., Hewett, P., Jetzer, P., Kaplan, J., Paulin-Henriksson, S., Smartt, S. J., Stalin, C. S., and Tsapras, Y. (2004). Classical novae from the POINT-AGAPE microlensing survey of M31 - I. The nova catalogue. *MNRAS*, **353**(2), 571–588.
- Darnley, M. J., Bode, M. F., Kerins, E., Newsam, A. M., An, J., Baillon, P., Belokurov, V., Calchi Novati, S., Carr, B. J., Cr    , M., Evans, N. W., Giraud-H    ud, Y., Gould, A.,

- Hewett, P., Jetzer, P., Kaplan, J., Paulin-Henriksson, S., Smartt, S. J., Tsapras, Y., and Weston, M. (2006). Classical novae from the POINT-AGAPE microlensing survey of M31 - II. Rate and statistical characteristics of the nova population. *MNRAS*, **369**(1), 257–271.
- Darnley, M. J., Ribeiro, V. A. R. M., Bode, M. F., Hounsell, R. A., and Williams, R. P. (2012). On the Progenitors of Galactic Novae. *ApJ*, **746**(1), 61.
- Darnley, M. J., Hounsell, R., O’Brien, T. J., Henze, M., Rodríguez-Gil, P., Shafter, A. W., Shara, M. M., Vaytet, N. M. H., Bode, M. F., Ciardullo, R., Davis, B. D., Galera-Rosillo, R., Harman, D. J., Harvey, E. J., Healy, M. W., Ness, J. U., Ribeiro, V. A. R. M., and Williams, S. C. (2019). A recurrent nova super-remnant in the Andromeda galaxy. *Nature*, **565**(7740), 460–463.
- De Gennaro Aquino, I., Shore, S. N., Schwarz, G. J., Mason, E., Starrfield, S., and Sion, E. M. (2014). The spectroscopic evolution of the recurrent nova T Pyxidis during its 2011 outburst. III. The ultraviolet development from iron curtain through the post-X-ray turnoff. *A&A*, **562**, A28.
- De Gennaro Aquino, I., Schröder, K. P., Mittag, M., Wolter, U., Jack, D., Eenens, P., González-Pérez, J. N., Hempelmann, A., Schmitt, J. H. M. M., Hauschildt, P. H., and Rauw, G. (2015). High spectral resolution monitoring of Nova V339 Delphini with TIGRE. *A&A*, **581**, A134.
- de Maupertuis, P. L. M. (1732). *Discours sur les differentes figures des astres*.
- de Vaucouleurs, G. (1958). Photoelectric photometry of the Andromeda Nebula in the UB system. *ApJ*, **128**, 465.
- de Vaucouleurs, G. (1978). The extragalactic distance scale. II. Distances of the nearest galaxies from primary indicators. *ApJ*, **223**, 730–739.
- Della Valle, M. (1988). *Distance modulus and rate of novae in M33*, volume 4 of *Astronomical Society of the Pacific Conference Series*, page 73.
- Della Valle, M. (1991). Nova LMC 1991 : evidence for a super-bright nova population. *A&A*, **252**, L9.
- Della Valle, M. and Calvani, M. (1990). *Quasi-Periodic Outburst Activity of Novae at Minimum*, volume 369, page 48.
- Della Valle, M. and Duerbeck, H. W. (1993). The space density of classical novae in the galactic disk. *A&A*, **271**, 175–179.
- Della Valle, M. and Gilmozzi, R. (2002). Rebirth of Novae as Distance Indicators due to Efficient Large Telescopes. *Science*, **296**, 1275.
- Della Valle, M. and Livio, M. (1994). On the Progenitors of Type IA Supernovae in Early-Type and Late-Type Galaxies. *ApJ*, **423**, L31.
- Della Valle, M. and Livio, M. (1995). The Calibration of Novae as Distance Indicators. *ApJ*, **452**, 704.
- Della Valle, M. and Livio, M. (1996). On the Frequency of Occurrence of Recurrent Novae and Their Role as Type IA Supernova Progenitors. *ApJ*, **473**, 240.
- Della Valle, M. and Livio, M. (1998). The Spectroscopic Differences between Disk and Thick-Disk/Bulge Novae. *ApJ*, **506**(2), 818–823.
- Della Valle, M. and Rosino, L. (1987). Semiregular Variations of Nova V841 Oph (1848) at Minimum. *Information Bulletin on Variable Stars*, **2995**, 1.
- Della Valle, M., Bianchini, A., Livio, M., and Orlo, M. (1992). On the possible existence of two classes of progenitors for classical novae. *A&A*, **266**, 232–236.
- Della Valle, M., Rosino, L., Bianchini, A., and Livio, M. (1994). The nova rate in galaxies of different Hubble types. *A&A*, **287**, 403–409.
- Della Valle, M., Gilmozzi, R., Bianchini, A., and Esenoglu, H. (1997). Study of nova shells. II. FH Serentis 1970 and QU Vulpeculae 1984, nebular expansion, parallax and luminosity. *A&A*, **325**, 1151–1156.
- Della Valle, M., Pasquini, L., Daou, D., and Williams, R. E. (2002). The evolution of Nova V382 Velorum 1999. *A&A*, **390**, 155–166.
- Downes, R. A. and Duerbeck, H. W. (2000). Optical Imaging of Nova Shells and the Maximum Magnitude-Rate of Decline Relationship. *AJ*, **120**(4), 2007–2037.
- Doyle, T. F., Shara, M. M., Lessing, A. M., and Zurek, D. (2019). A Hubble Space Telescope Survey for Novae in the Globular Clusters of M87. *ApJ*, **874**(1), 65.
- Duerbeck, H. W. (1981). Light curve types, absolute magnitudes, and physical properties of galactic novae. *PASP*, **93**, 165–175.
- Duerbeck, H. W. (1984). Constraints for Cataclysmic Binary Evolution as Derived from Space Distributions. *ApJSS*, **99**(80), 363.
- Duerbeck, H. W. (1987). A Reference Catalogue and Atlas of Galactic Novae. *Sp. Sci. Rev.*,

- 45(1-2), 1–14.
- Duerbeck, H. W. (1990). *Galactic Distribution and Outburst Frequency of Classical Novae*, volume 369, page 34.
- Duerbeck, H. W. (1992). The final decline of novae and the hibernation hypothesis. *MNRAS*, **258**, 629–638.
- Duncan, J. C. (1918). A Ninth Nova in the Andromeda Nebula. *PASP*, **30**(176), 255.
- Dunkley, J., Komatsu, E., Nolte, M. R., Spergel, D. N., Larson, D., Hinshaw, G., Page, L., Bennett, C. L., Gold, B., Jarosik, N., Weiland, J. L., Halpern, M., Hill, R. S., Kogut, A., Limon, M., Meyer, S. S., Tucker, G. S., Wollack, E., and Wright, E. L. (2009). Five-Year Wilkinson Microwave Anisotropy Probe Observations: Likelihoods and Parameters from the WMAP Data. *ApJS*, **180**(2), 306–329.
- Ederoclite, A., Mason, E., Della Valle, M., Gilmozzi, R., Williams, R. E., Germany, L., Saviane, I., Matteucci, F., Schaefer, B. E., Walter, F., Rudy, R. J., Lynch, D., Mazuk, S., Venturini, C. C., Puetter, R. C., Perry, R. B., Liller, W., and Rotter, A. (2006). Early spectral evolution of Nova Sagittarii 2004 (V5114 Sagittarii). *A&A*, **459**(3), 875–883.
- Eggleton, P. P. (1983). Approximations to the radii of Roche lobes. *ApJ*, **268**, 368–369.
- Ennis, D., Becklin, E. E., Beckwith, S., Elias, J., Gatley, I., Matthews, K., Neugebauer, G., and Willner, S. P. (1977). Infrared observations of Nova Cygni 1975. *ApJ*, **214**, 478–487.
- Eyres, S. P. S., Davis, R. J., Kenny, H. T., Bode, M. F., Dougherty, S. M., and Llyod, H. M. (1996). *MERLIN Observations of Symbiotic Novae*, volume 93 of *Astronomical Society of the Pacific Conference Series*, page 194.
- Ferrarese, L., Livio, M., Freedman, W., Saha, A., Stetson, P. B., Ford, H. C., Hill, R. J., and Madore, B. F. (1996). Discovery of a Nova in the Virgo Galaxy M100. *ApJ*, **468**, L95.
- Ferrarese, L., Côté, P., and Jordán, A. (2003). Hubble Space Telescope Observations of Novae in M49. *ApJ*, **599**(2), 1302–1319.
- Fields, B. D. (2011). The Primordial Lithium Problem. *Annual Review of Nuclear and Particle Science*, **61**, 47–68.
- Fields, B. D., Molaro, P., and Sarkar, S. (2014). Big-Bang Nucleosynthesis. *arXiv e-prints*, page arXiv:1412.1408.
- Finzell, T., Chomiuk, L., Metzger, B. D., Walter, F. M., Linford, J. D., Mukai, K., Nelson, T., Weston, J. H. S., Zheng, Y., Sokoloski, J. L., Mioduszewski, A., Rupen, M. P., Dong, S., Starrfield, S., Cheung, C. C., Woodward, C. E., Taylor, G. B., Bohlsen, T., Buil, C., Prieto, J., Wagner, R. M., Bensby, T., Bond, I. A., Sumi, T., Bennett, D. P., Abe, F., Koshimoto, N., Suzuki, D., Tristram, P. J., Christie, G. W., Natusch, T., McCormick, J., Yee, J., and Gould, A. (2018). A Detailed Observational Analysis of V1324 Sco, the Most Gamma-Ray-luminous Classical Nova to Date. *ApJ*, **852**(2), 108.
- Ford, H. C. and Ciardullo, R. (1988). Novae as distance indicators. Technical report.
- Franck, J. R., Shafter, A. W., Hornoch, K., and Misselt, K. A. (2012). The Nova Rate in NGC 2403. *ApJ*, **760**(1), 13.
- Franckowiak, A., Jean, P., Wood, M., Cheung, C. C., and Buson, S. (2018). Search for gamma-ray emission from Galactic novae with the Fermi -LAT. *A&A*, **609**, A120.
- Friedjung, M. (1979). Upper limit of the Li/Na ratio in novae. *A&A*, **77**, 357–358.
- Fujimoto, M. Y. (1982). A Theory of Hydrogen Shell Flashes on Accreting White Dwarfs - Part Two - the Stable Shell Burning and the Recurrence Period of Shell Flashes. *ApJ*, **257**, 767.
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., Prusti, T., de Bruijne, J. H. J., Babusaux, C., Bailer-Jones, C. A. L., Biermann, M., Evans, D. W., Eyer, L., Jansen, F., Jordi, C., Klioner, S. A., Lammers, U., Lindegren, L., Luri, X., Mignard, F., Panem, C., Pourbaix, D., Randich, S., Sartoretti, P., Siddiqui, H. I., Soubiran, C., van Leeuwen, F., Walton, N. A., Arenou, F., Bastian, U., Cropper, M., Drimmel, R., Katz, D., Lattanzi, M. G., Bakker, J., Cacciari, C., Castañeda, J., Chaoul, L., Cheek, N., De Angeli, F., Fabricius, C., Guerra, R., Holl, B., Masana, E., Messineo, R., Mowlavi, N., Nienartowicz, K., Panuzzo, P., Portell, J., Riello, M., Seabroke, G. M., Tanga, P., Thévenin, F., Gracia-Abril, G., Comoretto, G., García-Reinaldos, M., Teyssier, D., Altmann, M., Andrae, R., Audard, M., Bellas-Velidis, I., Benson, K., Berthier, J., Blomme, R., Burgess, P., Busso, G., Carry, B., Cellino, A., Clementini, G., Clotet, M., Creevey, O., Davidson, M., De Ridder, J., Delchambre, L., Dell’Oro, A., Ducourant, C., Fernández-Hernández, J., Fouesneau, M., Frémat, Y., Galluccio, L., García-Torres, M., González-Núñez, J., González-Vidal, J. J., Gosset, E., Guy, L. P., Halbwachs, J. L., Hambly, N. C., Harrison, D. L., Hernández, J., Hestroffer, D., Hodgkin, S. T., Hutton, A., Jasiewicz, G., Jean-Antoine-Piccolo, A., Jordan, S., Korn, A. J., Krone-Martins, A.,

- Lanzafame, A. C., Lebzelter, T., Löffler, W., Manteiga, M., Marrese, P. M., Martín-Fleitas, J. M., Moitinho, A., Mora, A., Muinonen, K., Osinde, J., Pancino, E., Pauwels, T., Petit, J. M., Recio-Blanco, A., Richards, P. J., Rimoldini, L., Robin, A. C., Sarro, L. M., Siopis, C., Smith, M., Sozzetti, A., Süveges, M., Torra, J., van Reeven, W., Abbas, U., Abreu Aramburu, A., Accart, S., Aerts, C., Altavilla, G., Alvarez, M. A., Alvarez, R., Alves, J., Anderson, R. I., Andrei, A. H., Anglada Varela, E., Antiche, E., Antoja, T., Arcay, B., Astraatmadja, T. L., Bach, N., Baker, S. G., Balaguer-Núñez, L., Balm, P., Barache, C., Barata, C., Barbato, D., Barblan, F., Barklem, P. S., Barrado, D., Barros, M., Barstow, M. A., Bartholomé Muñoz, S., Bassilana, J. L., Becciani, U., Bellazzini, M., Berihuete, A., Bertone, S., Bianchi, L., Bienaymé, O., Blanco-Cuaresma, S., Boch, T., Boeche, C., Bombrun, A., Borrachero, R., Bossini, D., Bouquillon, S., Bourda, G., Bragaglia, A., Bramante, L., Breddels, M. A., Bressan, A., Brouillet, N., Brüsmeister, T., Brugaletta, E., Bucciarelli, B., Burlacu, A., Busonero, D., Butkevich, A. G., Buzzì, R., Caffau, E., Cancelliere, R., Cannizzaro, G., Cantat-Gaudin, T., Carballo, R., Carlucci, T., Carrasco, J. M., Casamiquela, L., Castellani, M., Castro-Ginard, A., Charlot, P., Chemin, L., Chiavassa, A., Cocozza, G., Costigan, G., Cowell, S., Crifo, F., Crosta, M., Crowley, C., Cuypers, J., Dafonte, C., Damerdjì, Y., Dapergolas, A., David, P., David, M., de Laverny, P., De Luise, F., De March, R., de Martino, D., de Souza, R., de Torres, A., Debosscher, J., del Pozo, E., Delbo, M., Delgado, A., Delgado, H. E., Di Matteo, P., Diakite, S., Diener, C., Distefano, E., Dolding, C., Drazinos, P., Durán, J., Edvardsson, B., Enke, H., Eriksson, K., Esquej, P., Eynard Bontemps, G., Fabre, C., Fabrizio, M., Faigler, S., Falcão, A. J., Farràs Casas, M., Federici, L., Fedorets, G., Fernique, P., Figueras, F., Filippi, F., Findeisen, K., Fonti, A., Fraile, E., Fraser, M., Frézouls, B., Gai, M., Galleti, S., Garabato, D., García-Sedano, F., Garofalo, A., Garralda, N., Gavel, A., Gavras, P., Gerssen, J., Geyer, R., Giacobbe, P., Gilmore, G., Girona, S., Giuffrida, G., Glass, F., Gomes, M., Granvik, M., Gueguen, A., Guerrier, A., Guiraud, J., Gutiérrez-Sánchez, R., Haigron, R., Hatzidimitriou, D., Hauser, M., Haywood, M., Heiter, U., Helmi, A., Heu, J., Hilger, T., Hobbs, D., Hofmann, W., Holland, G., Huckle, H. E., Hypki, A., Icardi, V., Janßen, K., Jevardat de Fombelle, G., Jonker, P. G., Juhász, Á. L., Julbe, F., Karampelas, A., Kewley, A., Klar, J., Kochoska, A., Kohley, R., Kolenberg, K., Kontizas, M., Kontizas, E., Koposov, S. E., Kordopatis, G., Kostrzewa-Rutkowska, Z., Koubsky, P., Lambert, S., Lanza, A. F., Lasne, Y., Lavigne, J. B., Le Fustec, Y., Le Poncin-Lafitte, C., Lebreton, Y., Leccia, S., Leclerc, N., Lecoœur-Taïbi, I., Lenhardt, H., Leroux, F., Liao, S., Licata, E., Lindstrøm, H. E. P., Lister, T. A., Livanou, E., Lobel, A., López, M., Managau, S., Mann, R. G., Mantelet, G., Marchal, O., Marchant, J. M., Marconi, M., Marinoni, S., Marschall, G., Marshall, D. J., Martino, M., Marton, G., Mary, N., Massari, D., Matijević, G., Mazeh, T., McMillan, P. J., Messina, S., Michalik, D., Millar, N. R., Molina, D., Molinaro, R., Molnár, L., Montegriffo, P., Mor, R., Morbidelli, R., Morel, T., Morris, D., Mulone, A. F., Muraveva, T., Musella, I., Nelemans, G., Nicastro, L., Noval, L., O'Mullane, W., Ordénovic, C., Ordóñez-Blanco, D., Osborne, P., Pagani, C., Pagano, I., Pailler, F., Palacin, H., Palaversa, L., Panahi, A., Pawlak, M., Piersimoni, A. M., Pineau, F. X., Plachy, E., Plum, G., Poggio, E., Poujoulet, E., Prša, A., Pulone, L., Racero, E., Ragaini, S., Rambaux, N., Ramos-Lerate, M., Regibo, S., Reylé, C., Riclet, F., Ripepi, V., Riva, A., Rivard, A., Rixon, G., Roegiers, T., Roelens, M., Romero-Gómez, M., Rowell, N., Royer, F., Ruiz-Dern, L., Sadowski, G., Sagristà Sellés, T., Sahlmann, J., Salgado, J., Salguero, E., Sanna, N., Santana-Ros, T., Sarasso, M., Savietto, H., Schultheis, M., Sciacca, E., Segol, M., Segovia, J. C., Ségransan, D., Shih, I. C., Siltala, L., Silva, A. F., Smart, R. L., Smith, K. W., Solano, E., Solitro, F., Sordo, R., Soria Nieto, S., Souchay, J., Spagna, A., Spoto, F., Stampa, U., Steele, I. A., Steidelmüller, H., Stephenson, C. A., Stoev, H., Suess, F. F., Surdej, J., Szabados, L., Szegedi-Elek, E., Tapiador, D., Taris, F., Tauran, G., Taylor, M. B., Teixeira, R., Terrett, D., Teyssandier, P., Thuillot, W., Titarenko, A., Torra Clotet, F., Turon, C., Ulla, A., Utrilla, E., Uzzi, S., Vaillant, M., Valentini, G., Valette, V., van Elteren, A., Van Hemelryck, E., van Leeuwen, M., Vaschetto, M., Vecchiato, A., Veljanoski, J., Viala, Y., Vicente, D., Vogt, S., von Essen, C., Voss, H., Votruba, V., Voutsinas, S., Walmsley, G., Weiler, M., Wertz, O., Wevers, T., Wyrzykowski, L., Yoldas, A., Žerjal, M., Ziaeeppour, H., Zorec, J., Zschocke, S., Zucker, S., Zurbach, C., and Zwitter, T. (2018). Gaia Data Release 2. Summary of the contents and survey properties. *A&A*, **616**, A1.
- Gal-Yam, A. (2012). Luminous Supernovae. *Science*, **337**(6097), 927.
- Gallagher, J. S. and Starrfield, S. (1978). Theory and observations of classical novae. *ARA&A*, **16**, 171–214.
- Gavazzi, G. and Scodreggio, M. (1996). The mass dependence of the star formation history of

- disk galaxies. *A&A*, **312**, L29–L32.
- Gehrels, N., Chincarini, G., Giommi, P., Mason, K. O., Nousek, J. A., Wells, A. A., White, N. E., Barthelmy, S. D., Burrows, D. N., Cominsky, L. R., Hurley, K. C., Marshall, F. E., Mészáros, P., Roming, P. W. A., Angelini, L., Barbier, L. M., Belloni, T., Campana, S., Caraveo, P. A., Chester, M. M., Citterio, O., Cline, T. L., Cropper, M. S., Cummings, J. R., Dean, A. J., Feigelson, E. D., Fenimore, E. E., Frail, D. A., Fruchter, A. S., Garmire, G. P., Gendreau, K., Ghisellini, G., Greiner, J., Hill, J. E., Hunsberger, S. D., Krimm, H. A., Kulkarni, S. R., Kumar, P., Lebrun, F., Lloyd-Ronning, N. M., Markwardt, C. B., Mattson, B. J., Mushotzky, R. F., Norris, J. P., Osborne, J., Paczynski, B., Palmer, D. M., Park, H. S., Parsons, A. M., Paul, J., Rees, M. J., Reynolds, C. S., Rhoads, J. E., Sasseen, T. P., Schaefer, B. E., Short, A. T., Smale, A. P., Smith, I. A., Stella, L., Tagliaferri, G., Takahashi, T., Tashiro, M., Townsley, L. K., Tueller, J., Turner, M. J. L., Vietri, M., Voges, W., Ward, M. J., Willingale, R., Zerbi, F. M., and Zhang, W. W. (2004). The Swift Gamma-Ray Burst Mission. *ApJ*, **611**(2), 1005–1020.
- Gehrz, R. D. (1990). *New Infrared Results for Classical Novae*, volume 369, page 138.
- Gehrz, R. D., Grasdalén, G. L., Hackwell, J. A., and Ney, E. P. (1980a). The evolution of the dust shell of nova SER 1978. *ApJ*, **237**, 855–865.
- Gehrz, R. D., Hackwell, J. A., Grasdalén, G. L., Ney, E. P., Neugebauer, G., and Sellgren, K. (1980b). The optically thin dust shell of nova CYG 1978. *ApJ*, **239**, 570–580.
- Gehrz, R. D., Greenhouse, M. A., Hayward, T. L., Houck, J. R., Mason, C. G., and Woodward, C. E. (1995a). The Infrared Spectrum of the Optically Thin Dust Shell of V705 Cassiopeiae (Nova Cassiopeiae 1993). *ApJ*, **448**, L119.
- Gehrz, R. D., Jones, T. J., Matthews, K., Neugebauer, G., Woodward, C. E., Hayward, T. L., and Greenhouse, M. A. (1995b). The Neon Nova. III. The Infrared Light Curves of Nova QU Vulpeculae (Nova VUL 1984#2). *AJ*, **110**, 325.
- Gehrz, R. D., Truran, J. W., Williams, R. E., and Starrfield, S. (1998). Nucleosynthesis in Classical Novae and Its Contribution to the Interstellar Medium. *PASP*, **110**(743), 3–26.
- Gehrz, R. D., Woodward, C. E., Helton, L. A., Polomski, E. F., Hayward, T. L., Houck, J. R., Evans, A., Krautter, J., Shore, S. N., Starrfield, S., Truran, J., Schwarz, G. J., and Wagner, R. M. (2008). The Neon Abundance in the Ejecta of QU Vulpeculae from Late-Epoch Infrared Spectra. *ApJ*, **672**(2), 1167–1173.
- Gehrz, R. D., Evans, A., Helton, L. A., and Woodward, C. E. (2015). Infrared Observations of Novae in the SOFIA Era. In *EAS Publications Series*, volume 71-72 of *EAS Publications Series*, pages 143–146.
- Gehrz, R. D., Evans, A., Woodward, C. E., Helton, L. A., Banerjee, D. P. K., Srivastava, M. K., Ashok, N. M., Joshi, V., Eyres, S. P. S., Krautter, J., Kuin, N. P. M., Page, K. L., Osborne, J. P., Schwarz, G. J., Shenoy, D. P., Shore, S. N., Starrfield, S. G., and Wagner, R. M. (2018). The Temporal Development of Dust Formation and Destruction in Nova Sagittarii 2015#2 (V5668 SGR): A Panchromatic Study. *ApJ*, **858**(2), 78.
- Geisel, S. L., Kleinmann, D. E., and Low, F. J. (1970). Infrared Emission of Novae. *ApJ*, **161**, L101.
- Gill, C. D. and O’Brien, T. J. (2000). Hubble Space Telescope imaging and ground-based spectroscopy of old nova shells - I. FH Ser, V533 Her, BT Mon, DK Lac and V476 Cyg. *MNRAS*, **314**(1), 175–182.
- Glicenstein, J.-F. (1999a). Nova in Small Magellanic Cloud 1999 No. 3. *IAU Circ.*, **7286**.
- Glicenstein, J.-F. (1999b). Nova in the Small Magellanic Cloud. *IAU Circ.*, **7239**.
- Graham, J. A. (1979). The premaximum spectrum of a Magellanic Cloud nova. *PASP*, **91**, 79–82.
- Greggio, L. (2010). The rates of SNIa: a theoretical viewpoint. In *Progenitors and Environments of Stellar Explosions*, page 65.
- Güth, T., Shafter, A. W., and Misselt, K. A. (2010). The Nova Rate in M94 (NGC 4736). *ApJ*, **720**(2), 1155–1160.
- Hachisu, I. and Kato, M. (2006). A Universal Decline Law of Classical Novae. *ApJS*, **167**(1), 59–80.
- Hachisu, I. and Kato, M. (2007). A Universal Decline Law of Classical Novae. II. GK Persei 1901 and Novae in 2005. *ApJ*, **662**(1), 552–563.
- Hachisu, I. and Kato, M. (2009). Optical and Supersoft X-Ray Light-Curve Models of Classical Nova V2491 Cygni: A New Clue to the Secondary Maximum. *ApJ*, **694**(2), L103–L106.
- Hachisu, I. and Kato, M. (2010). A Prediction Formula of Supersoft X-ray Phase of Classical Novae. *ApJ*, **709**(2), 680–714.

- Hachisu, I. and Kato, M. (2012). A Extremely Massive White Dwarf of the Symbiotic Classical Nova V407 Cyg as Suggested by the RS Oph and U Sco Models. *Baltic Astronomy*, **21**, 68–75.
- Hachisu, I. and Kato, M. (2014). The UBV Color Evolution of Classical Novae. I. Nova-giant Sequence in the Color-Color Diagram. *ApJ*, **785**(2), 97.
- Hachisu, I. and Kato, M. (2015). A Light Curve Analysis of Classical Novae: Free-free Emission versus Photospheric Emission. *ApJ*, **798**(2), 76.
- Hachisu, I. and Kato, M. (2018). A Light Curve Analysis of Recurrent and Very Fast Novae in Our Galaxy, Magellanic Clouds, and M31. *ApJS*, **237**(1), 4.
- Hachisu, I. and Kato, M. (2019). A Light-curve Analysis of 32 Recent Galactic Novae: Distances and White Dwarf Masses. *ApJS*, **242**(2), 18.
- Hachisu, I., Kato, M., and Nomoto, K. (1999). A Wide Symbiotic Channel to Type IA Supernovae. *ApJ*, **522**(1), 487–503.
- Hachisu, I., Kato, M., and Cassatella, A. (2008). A Universal Decline Law of Classical Novae. III. GQ Muscae 1983. *ApJ*, **687**(2), 1236–1252.
- Hartwick, F. D. A. and Hutchings, J. B. (1978). Classical novae: a time-dependent optically thick wind model for the postmaximum phase. *ApJ*, **226**, 203–209.
- Hatano, K., Branch, D., Fisher, A., and Starrfield, S. (1997). On the spatial distribution and occurrence rate of Galactic classical novae. *MNRAS*, **290**(1), 113–118.
- Henze, M., Pietsch, W., Haberl, F., Hernanz, M., Sala, G., Della Valle, M., Hatzidimitriou, D., Rau, A., Hartmann, D. H., Greiner, J., Burwitz, V., and Fliri, J. (2010). X-ray monitoring of classical novae in the central region of M 31. I. June 2006–March 2007. *A&A*, **523**, A89.
- Henze, M., Pietsch, W., Haberl, F., Hernanz, M., Sala, G., Hatzidimitriou, D., Della Valle, M., Rau, A., Hartmann, D. H., and Burwitz, V. (2011). X-ray monitoring of classical novae in the central region of M 31. II. Autumn and winter 2007/2008 and 2008/2009. *A&A*, **533**, A52.
- Henze, M., Pietsch, W., Haberl, F., Hernanz, M., Sala, G., Della Valle, M., and Stiele, H. (2012). M31N 2008-05d: a M 31 disk nova with a dipping supersoft X-ray light curve. *A&A*, **544**, A44.
- Henze, M., Pietsch, W., Haberl, F., Della Valle, M., Riffeser, A., Sala, G., Hatzidimitriou, D., Hofmann, F., Hartmann, D. H., Koppenhoefer, J., Seitz, S., Williams, G. G., Hornoch, K., Itagaki, K., Kabashima, F., Nishiyama, K., Xing, G., Lee, C. H., Magnier, E., and Chambers, K. (2013). Supersoft X-rays reveal a classical nova in the M 31 globular cluster Bol 126. *A&A*, **549**, A120.
- Henze, M., Pietsch, W., Haberl, F., Della Valle, M., Sala, G., Hatzidimitriou, D., Hofmann, F., Hernanz, M., Hartmann, D. H., and Greiner, J. (2014). X-ray monitoring of classical novae in the central region of M 31 III. Autumn and winter 2009/10, 2010/11, and 2011/12. *A&A*, **563**, A2.
- Henze, M., Darnley, M. J., Williams, S. C., Kato, M., Hachisu, I., Anupama, G. C., Arai, A., Boyd, D., Burke, D., Ciardullo, R., Chinetti, K., Cook, L. M., Cook, M. J., Erdman, P., Gao, X., Harris, B., Hartmann, D. H., Hornoch, K., Horst, J. C., Hounsell, R., Husar, D., Itagaki, K., Kabashima, F., Kafka, S., Kaur, A., Kiyota, S., Kojiguchi, N., Kučáková, H., Kuramoto, K., Maehara, H., Mantero, A., Masci, F. J., Matsumoto, K., Naito, H., Ness, J. U., Nishiyama, K., Oksanen, A., Osborne, J. P., Page, K. L., Paunzen, E., Pavana, M., Pickard, R., Prieto-Arranz, J., Rodríguez-Gil, P., Sala, G., Sano, Y., Shafter, A. W., Sugiura, Y., Tan, H., Tordai, T., Vrástl, J., Wagner, R. M., Watanabe, F., Williams, B. F., Bode, M. F., Bruno, A., Buchheim, B., Crawford, T., Goff, B., Hernanz, M., Igarashi, A. S., José, J., Motta, M., O’Brien, T. J., Oswalt, T., Poyner, G., Ribeiro, V. A. R. M., Sabo, R., Shara, M. M., Shears, J., Starkey, D., Starrfield, S., and Woodward, C. E. (2018). Breaking the Habit: The Peculiar 2016 Eruption of the Unique Recurrent Nova M31N 2008-12a. *ApJ*, **857**(1), 68.
- Hernandez, S., Larsen, S., Trager, S., Kaper, L., and Groot, P. (2018). Metallicities of young massive clusters in NGC 5236 (M83). *MNRAS*, **473**(1), 826–837.
- Hernanz, M. (2012). Novae in γ -rays. *Bulletin of the Astronomical Society of India*, **40**, 377.
- Hernanz, M. (2015). Astrophysics: A lithium-rich stellar explosion. *Nature*, **518**(7539), 307–308.
- Hernanz, M. and Sala, G. (2010). X-ray observations of classical novae: Theoretical implications. *Astronomische Nachrichten*, **331**(2), 169.
- Hillebrandt, W. and Thielemann, F. K. (1982). Nucleosynthesis in novae - A source of Ne-E and Al-26. *ApJ*, **255**, 617–623.

- Hillman, Y., Shara, M. M., Prialnik, D., & Kovetz, A. 2020, *Nature Astronomy*
- Hjellming, R. M. (1995). *Radio Images and Light Curves for Nova V1974 Cyg 1992*, volume 205 of *Astrophysics and Space Science Library*, page 139.
- Hjellming, R. M. and Wade, C. M. (1970). Radio Novae. *ApJ*, **162**, L1.
- Hjellming, R. M., Wade, C. M., Vandenberg, N. R., and Newell, R. T. (1979). Radio emission from nova shells. *AJ*, **84**, 1619–1631.
- Hubble, E. and Sandage, A. (1953). The Brightest Variable Stars in Extragalactic Nebulae. I. M31 and M33. *ApJ*, **118**, 353.
- Hubble, E. P. (1929). A spiral nebula as a stellar system, Messier 31. *ApJ*, **69**, 103–158.
- Hyland, A. R. and Neugebauer, G. (1970). Infrared Observations of Nova Serpentis 1970. *ApJ*, **160**, L177.
- Iben, I., J. and Tutukov, A. V. (1984). Supernovae of type I as end products of the evolution of binaries with components of moderate initial mass. *ApJS*, **54**, 335–372.
- Iben, I., J. and Tutukov, A. V. (1985). On the evolution of close binaries with components of initial mass between 3 M and 12 M. *ApJS*, **58**, 661–710.
- Iben, Icko, J. and Livio, M. (1993). Common Envelopes in Binary Star Evolution. *PASP*, **105**, 1373.
- Iben, Icko, J. and Tutukov, A. V. (1992). Rare Thermonuclear Explosions in Short-Period Cataclysmic Variables, with Possible Application to the Nova-like Red Variable in the Galaxy M31. *ApJ*, **389**, 369.
- Iijima, T. (2012). Spectroscopic observations of the recurrent nova CI Aquilae during the 2000 outburst. *A&A*, **544**, A26.
- Ikiewicz, K., Mikołajewska, J., Belczyński, K., Wiktorowicz, G., and Karczmarek, P. (2019). Wind Roche lobe overflow as a way to make Type Ia supernovae from the widest symbiotic systems. *MNRAS*, **485**(4), 5468–5473.
- Iwanowska, W. and Burnicki, A. (1961). *The population assignment of novae*, page 311.
- Izzo, L., Ederoclitte, A., Della Valle, M., Mason, E., Williams, R. E., Altamore, T., Cassatella, A., Gilmozzi, R., Patat, F., Schmidtobreick, L., Selvelli, P., Tappert, C., Thater, S., Covone, G., Dall’Ora, M., and Paolillo, M. (2012). Optical and near infrared multi-site follow up of the recurrent nova T Pyx. *Mem. Soc. Astr. It.*, **83**, 830.
- Izzo, L., Della Valle, M., Mason, E., Matteucci, F., Romano, D., Pasquini, L., Vanzì, L., Jordan, A., Fernandez, J. M., Bluhm, P., Brahm, R., Espinoza, N., and Williams, R. (2015). Early Optical Spectra of Nova V1369 Cen Show the Presence of Lithium. *ApJ*, **808**(1), L14.
- Izzo, L., Molaro, P., Bonifacio, P., Della Valle, M., Cano, Z., de Ugarte Postigo, A., Prieto, J. L., Thöne, C., Vanzì, L., Zapata, A., and Fernandez, D. (2018). Beryllium detection in the very fast nova ASASSN-16kt (V407 Lupi). *MNRAS*, **478**(2), 1601–1610.
- Jacoby, G. H., Branch, D., Ciardullo, R., Davies, R. L., Harris, W. E., Pierce, M. J., Pritchett, C. J., Tonry, J. L., and Welch, D. L. (1992). A Critical Review of Selected Techniques for Measuring Extragalactic Distances. *PASP*, **104**, 599.
- Jordi, K., Grebel, E. K., and Ammon, K. (2006). Empirical color transformations between SDSS photometry and other photometric systems. *A&A*, **460**(1), 339–347.
- José, J. (2012). Classical nova explosions – hydrodynamics and nucleosynthesis. *Bulletin of the Astronomical Society of India*, **40**, 443.
- José, J. and Hernanz, M. (1998). Nucleosynthesis in Classical Novae: CO versus ONe White Dwarfs. *ApJ*, **494**(2), 680–690.
- Kantharia, N. G. (2017). Novae: I. The maximum magnitude relation with decline time (MMRD) and distance. *arXiv e-prints*, page arXiv:1703.04087.
- Kasliwal, M. M., Cenko, S. B., Kulkarni, S. R., Ofek, E. O., Quimby, R., and Rau, A. (2011). Discovery of a New Photometric Sub-class of Faint and Fast Classical Novae. *ApJ*, **735**(2), 94.
- Kato, M. and Hachisu, I. (1989). Mass Loss during Nova Outbursts on Various White Dwarf Masses. *ApJ*, **346**, 424.
- Kato, M. and Hachisu, I. (1994). Optically Thick Winds in Nova Outbursts. *ApJ*, **437**, 802.
- Kato, M. and Hachisu, I. (2012). Recurrent novae as progenitors of Type Ia supernovae. *Bulletin of the Astronomical Society of India*, **40**, 393.
- Kato, M., Hachisu, I., and Cassatella, A. (2009). A Universal Decline Law of Classical Novae. IV. V838 Her (1991): A Very Massive White Dwarf. *ApJ*, **704**(2), 1676–1688.
- Kato, M., Hachisu, I., and Henze, M. (2013). Novae in Globular Clusters. *ApJ*, **779**(1), 19.
- Kato, M., Saio, H., Hachisu, I., and Nomoto, K. (2014). Shortest Recurrence Periods of Novae. *ApJ*, **793**(2), 136.

- Kato, M., Saio, H., and Hachisu, I. (2015). Multi-wavelength Light Curve Model of the One-year Recurrence Period Nova M31N 2008-12A. *ApJ*, **808**(1), 52.
- Kaur, A. (2016). *The puzzling bulge to disk nova ratio in the Andromeda galaxy (M31)*. Ph.D. thesis, Clemson University.
- Kodaira, K. (1970). Energy Distribution around the Light Maximum of Nova Serpentis 1970. *PASJ*, **22**, 447.
- Kolb, U. and Politano, M. (1997). The contribution of O-Ne-Mg novae to the ^{26}Al production in the Galaxy. *A&A*, **319**, 909–922.
- Kopylov, I. M. (1952). *Comm. Crimean Ap. Obs.*, **9**, 116.
- Kopylov, I. M. (1955). *Izv. Krymskoi Astrofiz. Obs.*, **13**, 23.
- Kovetz, A. and Prialnik, D. (1985). CNO abundances resulting from diffusion in accreting nova progenitors. *ApJ*, **291**, 812–821.
- Kraft, R. P. (1964). Binary Stars among Cataclysmic Variables. III. Ten Old Novae. *ApJ*, **139**, 457.
- Kraft, R. P. (1965). Binary Stars among Cataclysmic Variables. VII. on the Kinematics and Space Distribution of W Ursae Majoris and U Geminorum-Type Stars. *ApJ*, **142**, 1588.
- Krautter, J., Oegelman, H., Starrfield, S., Wichmann, R., and Pfeffermann, E. (1996). ROSAT X-Ray Observations of Nova V1974 Cygni: The Rise and Fall of the Brightest Supersoft X-Ray Source. *ApJ*, **456**, 788.
- Kukarkin, B. V. (1949). *The study of the structure and evolution of stellar systems*.
- Lambert, D. L. and Reddy, B. E. (2004). Lithium abundances of the local thin disc stars. *MNRAS*, **349**(2), 757–767.
- Li, K.-L., Metzger, B. D., Chomiuk, L., Vurm, I., Strader, J., Finzell, T., Beloborodov, A. M., Nelson, T., Shappee, B. J., Kochanek, C. S., Prieto, J. L., Kafka, S., Holoien, T. W. S., Thompson, T. A., Luckas, P. J., and Itoh, H. (2017). A nova outburst powered by shocks. *Nature Astronomy*, **1**, 697–702.
- Li, W., Bloom, J. S., Podsiadlowski, P., Miller, A. A., Cenko, S. B., Jha, S. W., Sullivan, M., Howell, D. A., Nugent, P. E., Butler, N. R., Ofek, E. O., Kasliwal, M. M., Richards, J. W., Stockton, A., Shih, H.-Y., Bildsten, L., Shara, M. M., Bibby, J., Filippenko, A. V., Ganeshalingam, M., Silverman, J. M., Kulkarni, S. R., Law, N. M., Poznanski, D., Quimby, R. M., McCully, C., Patel, B., Maguire, K., and Shen, K. J. (2011). Exclusion of a luminous red giant as a companion star to the progenitor of supernova SN 2011fe. *Nature*, **480**(7377), 348–350.
- Liller, W. and Mayer, B. (1987). The rate of nova production in the galaxy. *PASP*, **99**, 606–609.
- Linford, J. D., Chomiuk, L., and Rupen, M. P. (2018). *ngVLA Studies of Classical Novae*, volume 517 of *Astronomical Society of the Pacific Conference Series*, page 271.
- Livio, M. (1992). Classical Novae and the Extragalactic Distance Scale. *ApJ*, **393**, 516.
- Livio, M. and Mazzali, P. (2018). On the progenitors of Type Ia supernovae. *Phys. Rept.*, **736**, 1–23.
- Livio, M. and Truran, J. W. (1994). On the Interpretation and Implications of Nova Abundances: an Abundance of Riches or an Overabundance of Enrichments. *ApJ*, **425**, 797.
- Livio, M., Riess, A., and Sparks, W. (2002). Will Jets Identify the Progenitors of Type Ia Supernovae? *ApJ*, **571**(2), L99–L102.
- Lloyd, H. M., O’Brien, T. J., and Bode, M. F. (1996). *Models for the Radio Emission from Classical Novae*, volume 93 of *Astronomical Society of the Pacific Conference Series*, page 200.
- Lodders, K., Palme, H., and Gail, H.-P. (2009). Abundances of the Elements in the Solar System. *Landolt Börnstein*.
- Lundmark, K. (1922). The Absolute Magnitudes of Novæ. *PASP*, **34**(200), 207.
- Lundmark, K. (1923). Some Facts and Suggestions Concerning Novæ. *PASP*, **35**(204), 95.
- Lundmark, K. (1927). Studies of Anagalactic Nebulae - First Paper. *Nova Acta Regiae Soc. Sci. Upsaliensis Ser. V*, pages 1–127.
- Lundmark, K. (1935). On the novae and their classification among the variable stars. *Meddelanden fran Lunds Astronomiska Observatorium Serie II*, **74**, 1–20.
- Madrid, J. P., Sparks, W. B., Ferguson, H. C., Livio, M., and Macchetto, D. (2007). Discovery of 13 Nova Candidates in M87. *ApJ*, **654**(1), L41–L44.
- Mannucci, F., Della Valle, M., Panagia, N., Cappellaro, E., Cresci, G., Maiolino, R., Petrosian, A., and Turatto, M. (2005). The supernova rate per unit mass. *A&A*, **433**(3), 807–814.
- Mannucci, F., Della Valle, M., and Panagia, N. (2006). Two populations of progenitors for

- Type Ia supernovae? *MNRAS*, **370**(2), 773–783.
- Maoz, D., Mannucci, F., and Nelemans, G. (2014). Observational Clues to the Progenitors of Type Ia Supernovae *ARA&A*, **52**, 107–170.
- Martin, P. and Dubus, G. (2013). Particle acceleration and non-thermal emission during the V407 Cygni nova outburst. *A&A*, **551**, A37.
- Martin, P., Dubus, G., Jean, P., Tatischeff, V., and Dosne, C. (2018). Gamma-ray emission from internal shocks in novae. *A&A*, **612**, A38.
- Martini, P., Wagner, R. M., Tomaney, A., Rich, R. M., Della Valle, M., and Hauschildt, P. H. (1999). Nova Sagittarii 1994 1 (V4332 Sagittarii): The Discovery and Evolution of an Unusual Luminous Red Variable Star. *AJ*, **118**(2), 1034–1042.
- Mason, C. G., Gehrz, R. D., Woodward, C. E., Smilowitz, J. B., Greenhouse, M. A., Hayward, T. L., and Houck, J. R. (1996). Infrared Observations of Dust Formation and Coronal Emission in Nova Aquilae 1995. *ApJ*, **470**, 577.
- Mason, C. G., Gehrz, R. D., Woodward, C. E., Smilowitz, J. B., Hayward, T. L., and Houck, J. R. (1998). The Infrared Development of V705 Cassiopeiae. *ApJ*, **494**(2), 783–791.
- Mason, E., Della Valle, M., Gilmozzi, R., Lo Curto, G., and Williams, R. E. (2005). Early decline spectra of nova SMC 2001 and nova LMC 2002. *A&A*, **435**(3), 1031–1042.
- Mason, E., Shore, S. N., De Gennaro Aquino, I., Izzo, L., Page, K., and Schwarz, G. J. (2018). V1369 Cen High-resolution Panchromatic Late Nebular Spectra in the Context of a Unified Picture for Nova Ejecta. *ApJ*, **853**(1), 27.
- Matteucci, F., D’Antona, F., and Timmes, F. X. (1995). News (ν s) on the Galactic evolution of lithium. *A&A*, **303**, 460.
- Matteucci, F., Renda, A., Pipino, A., and Della Valle, M. (2003). Modelling the nova rate in galaxies. *A&A*, **405**, 23–30.
- Mazzali, P. A., Nomoto, K., Cappellaro, E., et al. 2001, *ApJ*, 547, 988
- McLaughlin, D. B. (1939). The forms of light curves of novae. In *Publications of the American Astronomical Society*, volume 9 of *Publications of the American Astronomical Society*, page 157.
- McLaughlin, D. B. (1940). The Relation Between Absorption Velocity and Rate of Decline for Galactic Novae. *ApJ*, **91**, 369.
- McLaughlin, D. B. (1941). The Distances and Luminosities of Three Novae. *ApJ*, **93**, 417.
- McLaughlin, D. B. (1942). The Luminosities of Novae. *Popular Astronomy*, **50**, 233.
- McLaughlin, D. B. (1943). On the spectra of novae. *Publications of Michigan Observatory*, **8**(12), 149–194.
- McLaughlin, D. B. (1945). A note on the absorption of light in the galaxy. *AJ*, **51**, 97.
- McLaughlin, D. B. (1945). The Relation between Light-Curves and Luminosities of Novae. *PASP*, **57**(335), 69.
- McLaughlin, D. B. (1946). The classification of nova spectra. *AJ*, **52**, 46.
- McLaughlin, D. B. (1960). *The Spectra of Novae*, page 585.
- Metzger, B. D., Finzell, T., Vurm, I., Hascoët, R., Beloborodov, A. M., and Chomiuk, L. (2015). Gamma-ray novae as probes of relativistic particle acceleration at non-relativistic shocks. *MNRAS*, **450**(3), 2739–2748.
- Minkowski, R. (1948). Novae and Planetary Nebulae. *ApJ*, **107**, 106.
- Molaro, P., Izzo, L., Bonifacio, E., Bonifacio, P., Hernanz, M., Selvelli P., and Della Valle, M. (2020). Search for ^7Be in the outbursts of four recent novae. *MNRAS*, **492**(4), 4975–4985.
- Molaro, P., Izzo, L., Mason, E., Bonifacio, P., and Della Valle, M. (2016). Highly enriched ^7Be in the ejecta of Nova Sagittarii 2015 No. 2 (V5668 Sgr) and the Galactic ^7Li origin. *MNRAS*, **463**(1), L117–L121.
- Monck, W. H. S. (1885). New stars and shooting-stars. *The Observatory*, **8**, 335–336.
- Moraes, M. and Diaz, M. (2009). HR Del Remnant Anatomy Using Two-Dimensional Spectral Data and Three-Dimensional Photoionization Shell Models. *AJ*, **138**(6), 1541–1556.
- Morris, P. J., Cotter, G., Brown, A. M., and Chadwick, P. M. (2017). Gamma-ray novae: rare or nearby? *MNRAS*, **465**(1), 1218–1226.
- Mould, J., Cohen, J., Graham, J. R., Hamilton, D., Matthews, K., Picard, A., Reid, N., Schmidt, M., Soifer, T., Wilson, C., Rich, R. M., and Gunn, J. (1990). A Nova-like Red Variable in M31. *ApJ*, **353**, L35.
- Mróz, P., Udalski, A., Poleski, R., Soszyński, I., Szymański, M. K., Pietrzyński, G., Wyrzykowski, L., Ulaczyk, K., Kozłowski, S., Pietrukowicz, P., and Skowron, J. (2015). OGLE Atlas of Classical Novae. I. Galactic Bulge Objects. *ApJS*, **219**(2), 26.
- Mróz, P., Udalski, A., Poleski, R., Soszyński, I., Szymański, M. K., Pietrzyński, G.,

- Wyrzykowski, L., Ulaczyk, K., Kozłowski, S., Pietrukowicz, P., and Skowron, J. (2016). OGLE Atlas of Classical Novae. II. Magellanic Clouds. *ApJS*, **222**(1), 9.
- Mukai, K. and Ishida, M. (2001). The Early X-Ray Emission from V382 Velorum (Nova Velorum 1999): An Internal Shock Model. *ApJ*, **551**(2), 1024–1030.
- Mukai, K. and Sokoloski, J. L. (2019). The new science of novae. *Physics Today*, **72**(11), 38–44.
- Mukai, K., Orio, M., and Della Valle, M. (2008). Novae as a Class of Transient X-Ray Sources. *ApJ*, **677**(2), 1248–1252.
- Munari, U. and Renzini, A. (1992). Are Symbiotic Stars the Precursors of Type IA Supernovae? *ApJ*, **397**, L87.
- Munari, U. and Zwitter, T. (1997). Equivalent width of NA I and K I lines and reddening. *A&A*, **318**, 269–274.
- Munari, U., Mason, E., and Valisa, P. (2014). The narrow and moving HeII lines in nova KT Eridani. *A&A*, **564**, A76.
- Munari, U., Hambsch, F. J., and Frigo, A. (2017). Photometric evolution of seven recent novae and the double-component characterizing the light curve of those emitting in gamma rays. *MNRAS*, **469**(4), 4341–4358.
- Muraveva, T., Delgado, H. E., Clementini, G., Sarro, L. M., and Garofalo, A. (2018). RR Lyrae stars as standard candles in the Gaia Data Release 2 Era. *MNRAS*, **481**(1), 1195–1211.
- Nariai, K., Nomoto, K., and Sugimoto, D. (1980). Nova Explosion of Mass-Accreting White Dwarfs. *PASJ*, **32**, 473.
- Naylor, T., Charles, P. A., Mukai, K., and Evans, A. (1992). An observational case against nova hibernation. *MNRAS*, **258**, 449–456.
- Neill, J. D. and Shara, M. M. (2004). The H α Light Curves and Spatial Distribution of Novae in M81. *AJ*, **127**(2), 816–831.
- Neill, J. D. and Shara, M. M. (2005). A Possible High Nova Rate for Two Local Group Dwarf Galaxies: M32 and NGC 205. *AJ*, **129**(4), 1873–1885.
- Nelson, T., Chomiuk, L., Roy, N., Sokoloski, J. L., Mukai, K., Krauss, M. I., Mioduszewski, A. J., Rupen, M. P., and Weston, J. (2014). The 2011 Outburst of Recurrent Nova T Pyx: Radio Observations Reveal the Ejecta Mass and Hint at Complex Mass Loss. *ApJ*, **785**(1), 78.
- Ness, J. U., Schwarz, G. J., Retter, A., Starrfield, S., Schmitt, J. H. M. M., Gehrels, N., Burrows, D., and Osborne, J. P. (2007). Swift X-Ray Observations of Classical Novae. *ApJ*, **663**(1), 505–515.
- Ness, J. U., Drake, J. J., Starrfield, S., Bode, M. F., O’Brien, T. J., Evans, A., Eyres, S. P. S., Helton, L. A., Osborne, J. P., Page, K. L., Schneider, C., and Woodward, C. E. (2009). High-Resolution X-Ray Spectroscopy of the Evolving Shock in the 2006 Outburst of RS Ophiuchi. *AJ*, **137**(2), 3414–3436.
- Ness, J. U., Beardmore, A. P., Osborne, J. P., Kuulkers, E., Henze, M., Piro, A. L., Drake, J. J., Dobrotka, A., Schwarz, G., Starrfield, S., Kretschmar, P., Hirsch, M., and Wilms, J. (2015). Short-period X-ray oscillations in super-soft novae and persistent super-soft sources. *A&A*, **578**, A39.
- Newton, I. (1729). *The Mathematical Principles of Natural Philosophy*.
- Nomoto, K. (1982a). Accreting white dwarf models for type I supernovae. I - Presupernova evolution and triggering mechanisms. *ApJ*, **253**, 798–810.
- Nomoto, K. (1982b). Accreting white dwarf models for type I supernovae. II. Off-center detonation supernovae. *ApJ*, **257**, 780–792.
- Nomoto, K. and Kondo, Y. (1991). Conditions for Accretion-induced Collapse of White Dwarfs. *ApJ*, **367**, L19.
- O’Brien, T., Rupen, M., Chomiuk, L., Ribeiro, V., Bode, M., Sokoloski, J., and Woudt, P. A. (2015). Thermal radio emission from novae & symbiotics with the Square Kilometre Array. In *Advancing Astrophysics with the Square Kilometre Array (AASKA14)*, page 62.
- Oegelman, H., Beuermann, K., and Krautter, J. (1984). The detection of X-rays from nova MUSCAE 1983 with the EXOSAT satellite. *ApJ*, **287**, L31–L34.
- Oegelman, H., Orio, M., Krautter, J., and Starrfield, S. (1993). Detection of supersoft X-ray emission from GQ Muscae nine years after a nova outburst. *Nature*, **361**(6410), 331–333.
- Oort, J. H. (1926). Asymmetry in the distribution of stellar velocities. *The Observatory*, **49**, 302–304.
- Orio, M. (2006). A Close Look at the Population of Supersoft and Quasi-soft X-Ray Sources Observed in M31 with XMM-Newton. *ApJ*, **643**(2), 844–862.

- Orio, M., Covington, J., and Ögelman, H. (2001). X-ray emission from classical and recurrent novae observed with ROSAT. *A&A*, **373**, 542–554.
- Orio, M., Tepedelenlioglu, E., Starrfield, S., Woodward, C. E., and Della Valle, M. (2005). Chandra Observations of the Recurrent Nova IM Normae. *ApJ*, **620**(2), 938–942.
- Orio, M., Nelson, T., Bianchini, A., Di Mille, F., and Harbeck, D. (2010). A Census of the Supersoft X-ray Sources in M31. *ApJ*, **717**(2), 739–765.
- Orio, M., Ness, J. U., Dobrotka, A., Gatuzz, E., Ospina, N., Aydi, E., Behar, E., Buckley, D. A. H., Ciroi, S., Della Valle, M., Hernanz, M., Henze, M., Osborne, J. P., Page, K. L., Rauch, T., Sala, G., Starrfield, S., Williams, R. E., Woodward, C. E., and Zemko, P. (2018). What We Learn from the X-Ray Grating Spectra of Nova SMC 2016. *ApJ*, **862**(2), 164.
- Ortolani, S. and D’Odorico, S. (1980). A discussion on the nature of the Herbig-Haro object No 1 from its far UV spectrum. *A&A*, **83**, L8–L9.
- Osborne, J. P. (2015). Getting to know classical novae with Swift. *Journal of High Energy Astrophysics*, **7**, 117–125.
- Osborne, J. P., Page, K. L., Beardmore, A. P., Bode, M. F., Goad, M. R., O’Brien, T. J., Starrfield, S., Rauch, T., Ness, J. U., Krautter, J., Schwarz, G., Burrows, D. N., Gehrels, N., Drake, J. J., Evans, A., and Eyres, S. P. S. (2011). The Supersoft X-ray Phase of Nova RS Ophiuchi 2006. *ApJ*, **727**(2), 124.
- Osterbrock, D. E. & Ferland, G. J. 2006, *Astrophysics of gaseous nebulae and active galactic nuclei*
- Özdönmez, A., Ege, E., Güver, T., and Ak, T. (2018). A new catalogue of Galactic novae: investigation of the MMRD relation and spatial distribution. *MNRAS*, **476**(3), 4162–4186.
- Paczynski, B. (1965). Cataclysmic Variables Among Binary Stars. II. Physical Parameters for Novae. *Acta Astr.*, **15**, 197.
- Page, K. L., Osborne, J. P., Kuin, N. P. M., Henze, M., Walter, F. M., Beardmore, A. P., Bode, M. F., Darnley, M. J., Delgado, L., Drake, J. J., Hernanz, M., Mukai, K., Nelson, T., Ness, J. U., Schwarz, G. J., Shore, S. N., Starrfield, S., and Woodward, C. E. (2015). Swift detection of the super-swift switch-on of the super-soft phase in nova V745 Sco (2014). *MNRAS*, **454**(3), 3108–3120.
- Pagnotta, A., Walker, E. S., and Schaefer, B. E. (2014). The Diffuse Source at the Center of LMC SNR 0509-67.5 is a Background Galaxy at $z = 0.031$. *ApJ*, **788**(2), 173.
- Pagnotta, A., Schaefer, B. E., Clem, J. L., et al. 2015, *ApJ*, 811, 32
- Panagia, N., Van Dyk, S. D., Weiler, K. W., Sramek, R. A., Stockdale, C. J., and Murata, K. P. (2006). A Search for Radio Emission from Type Ia Supernovae. *ApJ*, **646**(1), 369–377.
- Pastorello, A., Mason, E., Taubenberger, S., Fraser, M., Cortini, G., Tomasella, L., Botticella, M. T., Elias-Rosa, N., Kotak, R., Smartt, S. J., Benetti, S., Cappellaro, E., Turatto, M., Tartaglia, L., Djorgovski, S. G., Drake, A. J., Berton, M., Briganti, F., Brimacombe, J., Bufano, F., Cai, Y. Z., Chen, S., Christensen, E. J., Ciabattari, F., Congiu, E., Dimai, A., Inserra, C., Kankare, E., Magill, L., Maguire, K., Martinelli, F., Morales-Garoffolo, A., Ochner, P., Pignata, G., Reguitti, A., Sollerman, J., Spiro, S., Terreran, G., and Wright, D. E. (2019). Luminous red novae: Stellar mergers or giant eruptions? *A&A*, **630**, A75.
- Patat, F. and Hussain, G. A. J. (2013). *Selecting and Scheduling Observing Programmes at ESO*, volume 2, pages 231–256.
- Patat, F., Chandra, P., Chevalier, R., Justham, S., Podsiadlowski, P., Wolf, C., Gal-Yam, A., Pasquini, L., Crawford, I. A., Mazzali, P. A., Pauldrach, A. W. A., Nomoto, K., Benetti, S., Cappellaro, E., Elias-Rosa, N., Hillebrandt, W., Leonard, D. C., Pastorello, A., Renzini, A., Sabbadin, F., Simon, J. D., and Turatto, M. (2007a). Detection of Circumstellar Material in a Normal Type Ia Supernova. *Science*, **317**(5840), 924.
- Patat, F., Benetti, S., Justham, S., Mazzali, P. A., Pasquini, L., Cappellaro, E., Della Valle, M., Podsiadlowski, P., Turatto, M., Gal-Yam, A., and Simon, J. D. (2007b). Upper limit for circumstellar gas around the type Ia SN 2000cx. *A&A*, **474**(3), 931–936.
- Patat, F., Kerzendorf, W., Bordelon, D., Van de Ven, G., and Pritchard, T. (2019). The Distributed Peer Review Experiment. *The Messenger*, **177**, 3–13.
- Patterson, J. (1984). The evolution of cataclysmic and low-mass X-ray binaries. *ApJS*, **54**, 443–493.
- Patterson, J. (1998). Late Evolution of Cataclysmic Variables. *PASP*, **110**(752), 1132–1147.
- Payne-Gaposchkin, C. (1957). *The galactic novae*.
- Pease, F. G. (1917). A Suspected Nova in the Nebula N. G. C. 2841. *PASP*, **29**(171), 213.
- Peimbert, M. and Sarmiento, A. (1984). Novae and galactic chemical evolution. *Astronomy Express*, **1**, 97–103.

- Perlmutter, S., Aldering, G., Della Valle, M., Deustua, S., Ellis, R. S., Fabbro, S., Fruchter, A., Goldhaber, G., Groom, D. E., Hook, I. M., Kim, A. G., Kim, M. Y., Knop, R. A., Lidman, C., McMahon, R. G., Nugent, P., Pain, R., Panagia, N., Pennypacker, C. R., Ruiz-Lapuente, P., Schaefer, B., and Walton, N. (1998). Discovery of a supernova explosion at half the age of the Universe. *Nature*, **391**(6662), 51–54.
- Pfau, W. (1976). Recalibration of the absolute magnitudes of novae and application to nova Cygni 1975. *A&A*, **50**, 113–115.
- Phillips, M. M. (1993). The Absolute Magnitudes of Type IA Supernovae. *ApJ*, **413**, L105.
- Pietrzyński, G., Graczyk, D., Gallenne, A., Gieren, W., Thompson, I. B., Pilecki, B., Karczmarek, P., Górski, M., Suchomska, K., Taormina, M., Zgierski, B., Wielgórski, P., Kolaczowski, Z., Konorski, P., Villanova, S., Nardetto, N., Kervella, P., Bresolin, F., Kudritzki, R. P., Storm, J., Smolec, R., and Narloch, W. (2019). A distance to the Large Magellanic Cloud that is precise to one per cent. *Nature*, **567**(7747), 200–203.
- Pinto, C., Ness, J. U., Verbunt, F., Kaastra, J. S., Costantini, E., and Detmers, R. G. (2012). A phenomenological model for the X-ray spectrum of nova V2491 Cygni. *A&A*, **543**, A134.
- Plaut, L. (1965). *Distribution of Novae in the Galaxy*, page 311.
- Poggiani, R. 2018a, in *Accretion Processes in Cosmic Sources - II*, 57
- Poggiani, R. 2018b, arXiv e-prints, arXiv:1803.11529
- Politano, M., Livio, M., Truran, J. W., and Webbink, R. F. (1990). *The Theoretical Frequency of Classical Nova Outbursts as a Function of White Dwarf Mass*, volume 369, page 386.
- Prantzos, N. (2012). Production and evolution of Li, Be, and B isotopes in the Galaxy. *A&A*, **542**, A67.
- Pretorius, M. L. and Knigge, C. (2007). Discovery of a new cataclysmic variable through optical variability and X-ray emission. *A&A*, **461**(3), 1103–1106.
- Prialnik, D. and Kovetz, A. (1995). An Extended Grid of Multicycle Nova Evolution Models. *ApJ*, **445**, 789.
- Prialnik, D., Shara, M. M., and Shaviv, G. (1978). The evolution of a slow nova model with a $Z = 0.03$ envelope from pre-explosion to extinction. *A&A*, **62**(3), 339–348.
- Prialnik, D., Livio, M., Shaviv, G., and Kovetz, A. (1982). On the role of the accretion rate in nova outbursts. *ApJ*, **257**, 312–317.
- Pritchett, C. J. and van den Bergh, S. (1987). Observations of Novae in the Virgo Cluster. *ApJ*, **318**, 507.
- Rafanelli, P. & Rosino, L. 1978, *A&AS*, **31**, 337
- Ramírez, I., Fish, J. R., Lambert, D. L., and Allende Prieto, C. (2012). Lithium Abundances in nearby FGK Dwarf and Subgiant Stars: Internal Destruction, Galactic Chemical Evolution, and Exoplanets. *ApJ*, **756**(1), 46.
- Reeves, H., Richer, J., Sato, K., and Terasawa, N. (1990). On the Origin of Lithium. *ApJ*, **355**, 18.
- Ribeiro, V. A. R. M., Bode, M. F., Darnley, M. J., Harman, D. J., Newsam, A. M., O’Brien, T. J., Bohigas, J., Echevarría, J. M., Bond, H. E., Chavushyan, V. H., Costero, R., Coziol, R., Evans, A., Eyres, S. P. S., León-Tavares, J., Richer, M. G., Tovmassian, G., Starrfield, S., and Zharikov, S. V. (2009). The Expanding Nebular Remnant of the Recurrent Nova RS Ophiuchi (2006). II. Modeling of Combined Hubble Space Telescope Imaging and Ground-based Spectroscopy. *ApJ*, **703**(2), 1955–1963.
- Ribeiro, V. A. R. M., Bode, M. F., Darnley, M. J., Barnsley, R. M., Munari, U., and Harman, D. J. (2013a). Morpho-kinematical modelling of Nova Eridani 2009 (KT Eri). *MNRAS*, **433**(3), 1991–1996.
- Ribeiro, V. A. R. M., Munari, U., and Valisa, P. (2013b). Optical Morphology, Inclination, and Expansion Velocity of the Ejected Shell of Nova Monocerotis 2012. *ApJ*, **768**(1), 49.
- Rich, R. M., Mould, J., Picard, A., Frogel, J. A., and Davies, R. (1989). Luminous M Giants in the Bulge of M31. *ApJ*, **341**, L51.
- Riess, A. G., Filippenko, A. V., Challis, P., Clocchiatti, A., Diercks, A., Garnavich, P. M., Gilliland, R. L., Hogan, C. J., Jha, S., Kirshner, R. P., Leibundgut, B., Phillips, M. M., Reiss, D., Schmidt, B. P., Schommer, R. A., Smith, R. C., Spyromilio, J., Stubbs, C., Suntzeff, N. B., and Tonry, J. (1998). Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant. *AJ*, **116**(3), 1009–1038.
- Riess, A. G., Fliri, J., and Valls-Gabaud, D. (2012). Cepheid Period-Luminosity Relations in the Near-infrared and the Distance to M31 from the Hubble Space Telescope Wide Field Camera 3. *ApJ*, **745**(2), 156.
- Riess, A. G., Casertano, S., Yuan, W., Macri, L. M., and Scolnic, D. (2019). Large Magellanic

- Cloud Cepheid Standards Provide a 1% Foundation for the Determination of the Hubble Constant and Stronger Evidence for Physics beyond Λ CDM. *ApJ*, **876**(1), 85.
- Ritchey, G. W. (1917a). Another Faint Nova in the Andromeda Nebula. *PASP*, **29**(172), 257.
- Ritchey, G. W. (1917b). Novae in Spiral Nebulae. *PASP*, **29**(171), 210.
- Ritchey, G. W. (1918). Three Additional Novae in the Andromeda Nebula. *PASP*, **30**(174), 162.
- Ritter, H., Politano, M., Livio, M., and Webbink, R. F. (1991). The White Dwarf Mass Distribution in Classical Nova Systems. *ApJ*, **376**, 177.
- Robbins, R. R. (1968a). The Helium Triplet Spectrum in Expanding Nebulae. I. The Capture-Cascade Intensities. *ApJ*, **151**, 497.
- Robbins, R. R. (1968b). The Helium Triplet Spectrum in Expanding Nebulae, II. Self-Absorption. *ApJ*, **151**, 511.
- Robinson, E. L. (1975). Preeruption light curves of novae2. *AJ*, **80**, 515.
- Romano, D. and Matteucci, F. (2003). Nova nucleosynthesis and Galactic evolution of the CNO isotopes. *MNRAS*, **342**(1), 185–198.
- Romano, D., Matteucci, F., Ventura, P., and D’Antona, F. (2001). The stellar origin of ^7Li . Do AGB stars contribute a substantial fraction of the local Galactic lithium abundance? *A&A*, **374**, 646–655.
- Rosino, L. (1956). Nuove ricerche sulle stelle variabili della nebulosa di Orione. *Mem. Soc. Astr. It.*, **27**, 335.
- Rosino, L. (1964). Novae in Messier 31 discovered and observed at Asiago from 1955 to 1963. *Annales d’Astrophysique*, **27**, 498–505.
- Rosino, L. (1973). Novae in M 31 discovered and observed at Asiago from 1963 to 1970. *A&AS*, **9**, 347.
- Rosino, L. and Bianchini, A. (1973). Discovery of Five Novae in Messier 33 and a Supernova in a Field Galaxy. *A&A*, **22**, 461.
- Rosino, L., Ciatti, F., and Della Valle, M. (1986). Researches on novae and related objects. I. The spectral evolution of nova FH SER (1970). *A&A*, **158**, 34–44.
- Rosino, L., Capaccioli, M., D’Onofrio, M., and Della Valle, M. (1989). Fifty-Two Novae in M31 Discovered and Observed at Asiago from 1971 to 1986. *AJ*, **97**, 83.
- Roy, N., Chomiuk, L., Sokoloski, J. L., et al. 2012a, Bulletin of the Astronomical Society of India, 40, 293
- Roy, N., Kantharia, N. G., Eyres, S. P. S., et al. 2012b, MNRAS, 427, L55
- Rupen, M. P., Mioduszewski, A. J., and Sokoloski, J. L. (2008). An Expanding Shell and Synchrotron Jet in RS Ophiuchi. *ApJ*, **688**(1), 559–567.
- Sackmann, I. J. and Boothroyd, A. I. (1992). The Creation of Superrich Lithium Giants. *ApJ*, **392**, L71.
- Sackmann, I. J. and Boothroyd, A. I. (1999). Creation of ^7Li and Destruction of ^3He , ^9Be , ^{10}B , and ^{11}B in Low-Mass Red Giants, Due to Deep Circulation. *ApJ*, **510**(1), 217–231.
- Sadakane, K., Tajitsu, A., Mizoguchi, S., Arai, A., and Naito, H. (2010). Discovery of Multiple High-Velocity Narrow Circumstellar NaI D Lines in Nova V1280 Sco. *PASJ*, **62**, L5.
- Sanford, R. F. (1918). Two Novae in the Andromeda Nebula. *PASP*, **30**(178), 341.
- Sanford, R. F. (1919). Two New Novae in the Andromeda Nebula. *PASP*, **31**(180), 109.
- Sawyer, H. B. (1938). The Bright Nova of 1860 in the Globular Cluster Messier 80. and its Relation to Supernovae. *J. Royal. Astr. Soc. Canada*, **32**, 69.
- Sawyer Hogg, H. and Wehlau, A. (1964). Probable nova in the globular cluster M 14. *AJ*, **69**, 141.
- Sbordone, L., Bonifacio, P., Caffau, E., Ludwig, H.-G., Behara, N. T., González Hernández, J. I., Steffen, M., Cayrel, R., Freytag, B., van’t Veer, C., Molaro, P., Plez, B., Sivarani, T., Spite, M., Spite, F., Beers, T. C., Christlieb, N., François, P., and Hill, V. (2010). The metal-poor end of the Spite plateau. I. Stellar parameters, metallicities, and lithium abundances. *A&A*, **522**, A26.
- Schaefer, G. H., Brummelaar, T. T., Gies, D. R., et al. 2014, Nature, 515, 234
- Schaefer, B. E. (2018). The distances to Novae as seen by Gaia. *MNRAS*, **481**(3), 3033–3051.
- Schatzman, E. (1951). Remarques sur le phénomène de Nova: IV. L’onde de détonation due à l’isotope ^3He . *Annales d’Astrophysique*, **14**, 294.
- Schmidt, B. P., Suntzeff, N. B., Phillips, M. M., Schommer, R. A., Clocchiatti, A., Kirshner, R. P., Garnavich, P., Challis, P., Leibundgut, B., Spyromilio, J., Riess, A. G., Filippenko, A. V., Hamuy, M., Smith, R. C., Hogan, C., Stubbs, C., Diercks, A., Reiss, D., Gilliland, R., Tonry, J., Maza, J., Dressler, A., Walsh, J., and Ciardullo, R. (1998). The High-Z Supernova

- Search: Measuring Cosmic Deceleration and Global Curvature of the Universe Using Type IA Supernovae. *ApJ*, **507**(1), 46–63.
- Schmidt, T. (1957). Die Lichtkurven-Leuchtkraft-Beziehung Neuer Sterne. Mit 8 Textabbildungen. *ZAP*, **41**, 182.
- Schwarz, G. J. (2002). A New Abundance Analysis of the ONeMg Nova QU Vulpeculae. *ApJ*, **577**(2), 940–950.
- Schwarz, G. J., Shore, S. N., Starrfield, S., Hauschildt, P. H., Della Valle, M., and Baron, E. (2001). Multiwavelength analyses of the extraordinary nova LMC 1991*. *MNRAS*, **320**(1), 103–123.
- Schwarz, G. J., Shore, S. N., Starrfield, S., and Vanlandingham, K. M. (2007). Abundance Analysis of the Extremely Fast ONeMg Novae V838 Herculis and V4160 Sagittarii. *ApJ*, **657**(1), 453–464.
- Schwarz, G. J., Ness, J.-U., Osborne, J. P., Page, K. L., Evans, P. A., Beardmore, A. P., Walter, F. M., Helton, L. A., Woodward, C. E., Bode, M., Starrfield, S., and Drake, J. J. (2011). Swift X-Ray Observations of Classical Novae. II. The Super Soft Source Sample. *ApJS*, **197**(2), 31.
- Schwöpe, A. D. (2018). Exploring the space density of X-ray selected cataclysmic variables. *A&A*, **619**, A62.
- Schwöpe, A. D., Brunner, H., Buckley, D., Greiner, J., Heyden, K. v. d., Neizvestny, S., Potter, S., and Schwarz, R. (2002). The census of cataclysmic variables in the ROSAT Bright Survey. *A&A*, **396**, 895–910.
- Seaquist, E. R. and Palimaka, J. (1977). Thick inhomogeneous shell models for the radio emission from Nova Serpentis 1970. *ApJ*, **217**, 781–787.
- Selvelli, P. and Gilmozzi, R. (2013). An archive study of 18 old novae. I. The UV spectra. *A&A*, **560**, A49.
- Selvelli, P. and Gilmozzi, R. (2019). A UV and optical study of 18 old novae with Gaia DR2 distances: mass accretion rates, physical parameters, and MMRD. *A&A*, **622**, A186.
- Selvelli, P., Molaro, P., and Izzo, L. (2018). Absorption and emission features of ${}^7\text{Be}$ II in the outburst spectra of V838 Her (Nova Her 1991). *MNRAS*, **481**(2), 2261–2272.
- Shafter, A. W. (1997). On the Nova Rate in the Galaxy. *ApJ*, **487**(1), 226–236.
- Shafter, A. W. (2013). Photometric and Spectroscopic Properties of Novae in the Large Magellanic Cloud. *AJ*, **145**(5), 117.
- Shafter, A. W. (2017). The Galactic Nova Rate Revisited. *ApJ*, **834**(2), 196.
- Shafter, A. W. and Irby, B. K. (2001). On the Spatial Distribution, Stellar Population, and Rate of Novae in M31. *ApJ*, **563**(2), 749–767.
- Shafter, A. W., Ciardullo, R., and Pritchett, C. J. (2000). Novae in External Galaxies: M51, M87, and M101. *ApJ*, **530**(1), 193–206.
- Shafter, A. W., Darnley, M. J., Hornoch, K., Filippenko, A. V., Bode, M. F., Ciardullo, R., Misselt, K. A., Hounsell, R. A., Chornock, R., and Matheson, T. (2011). A Spectroscopic and Photometric Survey of Novae in M31. *ApJ*, **734**(1), 12.
- Shafter, A. W., Darnley, M. J., Bode, M. F., and Ciardullo, R. (2012). On the Spectroscopic Classes of Novae in M33. *ApJ*, **752**(2), 156.
- Shafter, A. W., Henze, M., Rector, T. A., et al. 2015, *ApJS*, 216, 34
- Shafter, A. W. 2019, Extragalactic Novae; A historical perspective
- Shapley, H. (1917). A Faint Nova in the Nebula of Andromeda. *PASP*, **29**(171), 213–217.
- Shara, M. M. (1981). A theoretical explanation of the absolute magnitude-decline time $/M_B - t_{\text{sub } 3/}$ relationship for classical novae. *ApJ*, **243**, 926–934.
- Shara, M. M., Livio, M., Moffat, A. F. J., and Orio, M. (1986). Do Novae Hibernate during Most of the Millennia between Eruptions? Links between Dwarf and Classical Novae, and Implications for the Space Densities and Evolution of Cataclysmic Binaries. *ApJ*, **311**, 163.
- Shara, M. M., Moffat, A. F. J., and Potter, M. (1990). Spectroscopic Authentication of Very Old Nova Candidates. *AJ*, **100**, 540.
- Shara, M. M., Sandage, A., and Zurek, D. R. (1999). The Early Palomar Program (1950–1955) for the Discovery of Classical Novae in M81: Analysis of the Spatial Distribution, Magnitude Distribution, and Distance Suggestion. *PASP*, **111**(765), 1367–1381.
- Shara, M. M., Zurek, D. R., Baltz, E. A., Lauer, T. R., and Silk, J. (2004). An Erupting Classical Nova in a Globular Cluster of M87. *ApJ*, **605**(2), L117–L120.
- Shara, M. M., Martin, C. D., Seibert, M., Rich, R. M., Salim, S., Reitzel, D., Schiminovich, D., Deliyannis, C. P., Sarrazine, A. R., Kulkarni, S. R., Ofek, E. O., Brosch, N., Lépine, S., Zurek, D., De Marco, O., and Jacoby, G. (2007). An ancient nova shell around the dwarf

- nova Z Camelopardalis. *Nature*, **446**(7132), 159–162.
- Shara, M. M., Mizusawa, T., Wehinger, P., Zurek, D., Martin, C. D., Neill, J. D., Forster, K., and Seibert, M. (2012). AT Cnc: A Second Dwarf Nova with a Classical Nova Shell. *ApJ*, **758**(2), 121.
- Shara, M. M., Zurek, D., Schaefer, B. E., Bond, H. E., Godon, P., Mac Low, M.-M., Pagnotta, A., Prialnik, D., Sion, E. M., Toraskar, J., and Williams, R. E. (2015). HST Images Flash Ionization of Old Ejecta by the 2011 Eruption of Recurrent Nova T Pyxidis. *ApJ*, **805**(2), 148.
- Shara, M. M., Doyle, T. F., Lauer, T. R., Zurek, D., Neill, J. D., Madrid, J. P., Mikołajewska, J., Welch, D. L., and Baltz, E. A. (2016). A Hubble Space Telescope Survey for Novae in M87. I. Light and Color Curves, Spatial Distributions, and the Nova Rate. *ApJS*, **227**(1), 1.
- Shara, M. M., Doyle, T., Lauer, T. R., Zurek, D., Baltz, E. A., Kovetz, A., Madrid, J. P., Mikołajewska, J., Neill, J. D., Prialnik, D., Welch, D. L., and Yaron, O. (2017a). A Hubble Space Telescope Survey for Novae in M87. II. Snuffing out the Maximum Magnitude-Rate of Decline Relation for Novae as a Non-standard Candle, and a Prediction of the Existence of Ultrafast Novae. *ApJ*, **839**(2), 109.
- Shara, M. M., Hkiewicz, K., Mikołajewska, J., Pagnotta, A., Bode, M. F., Crause, L. A., Drozd, K., Faherty, J., Fuentes-Morales, I., and Grindlay, J. E. (2017b). Proper-motion age dating of the progeny of Nova Scorpii AD 1437. *Nature*, **548**(7669), 558–560.
- Shara, M. M., Doyle, T. F., Pagnotta, A., Garland, J. T., Lauer, T. R., Zurek, D., Baltz, E. A., Goerl, A., Kovetz, A., Machac, T., Madrid, J. P., Mikołajewska, J., Neill, J. D., Prialnik, D., Welch, D. L., and Yaron, O. (2018). A Hubble Space Telescope survey for novae in M87 - III. Are novae good standard candles 15 d after maximum brightness? *MNRAS*, **474**(2), 1746–1751.
- Sharov, A. S. (1972). Estimate for the Frequency of Novae in the Andromeda Nebula and our Galaxy. *Sov. Astr.*, **16**, 41.
- Sharov, A. S. (1979). The maximum luminosity-rate of decline relation and the problem of distance determination of novae. *Peremennye Zvezdy*, **21**, 141–147.
- Sharov, A. S. (1993). The rate of explosions of novae in galaxies of the Local Group. *Astronomy Letters*, **19**, 147.
- Shore, S. N. (2012). Spectroscopy of novae – a user’s manual. *Bulletin of the Astronomical Society of India*, **40**, 185.
- Shore, S. N. (2013). A unified model for the spectrophotometric development of classical and recurrent novae. The role of asphericity of the ejecta. *A&A*, **559**, L7.
- Shore, S. N. (2014). *The Spectroscopic Development of a Nova*, volume 490 of *Astronomical Society of the Pacific Conference Series*, page 145.
- Shore, S. N., Sonneborn, G., Starrfield, S., Riestra-Gonzalez, R., and Ake, T. B. (1993). The Early Ultraviolet Spectral Evolution of Nova Cygni 1992. *AJ*, **106**, 2408.
- Shore, S. N., Starrfield, S., and Sonneborn, G. (1996). The Ultraviolet and X-Ray View of the Demise of Nova V1974 Cygni. *ApJ*, **463**, L21.
- Shore, S. N., Schwarz, G., Bond, H. E., Downes, R. A., Starrfield, S., Evans, A., Gehrz, R. D., Hauschildt, P. H., Krautter, J., and Woodward, C. E. (2003). The Early Ultraviolet Evolution of the ONeMg Nova V382 Velorum 1999. *AJ*, **125**(3), 1507–1518.
- Shore, S. N., Augusteijn, T., Ederoclite, A., and Uthas, H. (2011). The spectroscopic evolution of the recurrent nova T Pyxidis during its 2011 outburst. I. The optically thick phase and the origin of moving lines in novae. *A&A*, **533**, L8.
- Shore, S. N., Kuin, N. P., Mason, E., and De Gennaro Aquino, I. (2018). Spectroscopic diagnostics of dust formation and evolution in classical nova ejecta. *A&A*, **619**, A104.
- Siebert, T., Coc, A., Delgado, L., Diehl, R., Greiner, J., Hernanz, M., Jean, P., José, J., Molaro, P., Pleintinger, M. M. M., Savchenko, V., Starrfield, S., Tatischeff, V., and Weinberger, C. (2018). Gamma-ray observations of Nova Sgr 2015 No. 2 with INTEGRAL. *A&A*, **615**, A107.
- Slavin, A. J., O’Brien, T. J., and Dunlop, J. S. (1995). A deep optical imaging study of the nebular remnants of classical novae. *MNRAS*, **276**(2), 353–371.
- Smith, D. A., Dhillon, V. S., and Marsh, T. R. (1998). The mass of the white dwarf in the old nova BT MON. *MNRAS*, **296**(3), 465–482.
- Sokoloski, J. L., Rupen, M. P., and Mioduszewski, A. J. (2008). Uncovering the Nature of Nova Jets: A Radio Image of Highly Collimated Outflows from RS Ophiuchi. *ApJ*, **685**(2), L137.

- Soraisam, M. D. (2016). *Novae a theoretical and observational study*. Ph.D. thesis, Ludwig-Maximilians-Universitat Munchen.
- Sparks, W. M. (1969). Dynamical Models of Novae. *ApJ*, **156**, 569.
- Sparks, W. M., Starrfield, S., and Truran, J. W. (1977). *a Review of the Thermonuclear Runaway Model of a Nova Outburst*, volume 65 of *Astrophysics and Space Science Library*, page 189.
- Sparks, W. M., Starrfield, S., and Truran, J. W. (1978). A hydrodynamic study of a slow nova outburst. *ApJ*, **220**, 1063–1075.
- Spite, F. (1990). Lithium in population II stars. *Mem. Soc. Astr. It.*, **61**, 663–675.
- Spite, F. and Spite, M. (1982). Abundance of lithium in unevolved halo stars and old disk stars - Interpretation and consequences. *A&A*, **115**, 357–366.
- Starrfield, S. (1999). Recent advances in studies of the nova outburst. *Phys. Rept.*, **311**, 371–381.
- Starrfield, S., Truran, J. W., Sparks, W. M., and Arnould, M. (1978). On ${}^7\text{Li}$ production in nova explosions. *ApJ*, **222**, 600–603.
- Starrfield, S., Schwarz, G. J., Truran, J. W., and Sparks, W. M. (1998). Hydrodynamic Studies of the Evolution of the Extraordinary Nova LMC 1991. In *American Astronomical Society Meeting Abstracts*, volume 193 of *American Astronomical Society Meeting Abstracts*, page 101.06.
- Starrfield, S., Sparks, W. M., Truran, J. W., and Wiescher, M. C. (2000). The Effects of New Nuclear Reaction Rates and Opacities on Hydrodynamic Simulations of the Nova Outburst. *ApJS*, **127**(2), 485–495.
- Starrfield, S., Iliadis, C., Hix, W. R., Timmes, F. X., and Sparks, W. M. (2009). The Effects of the pep Nuclear Reaction and Other Improvements in the Nuclear Reaction Rate Library on Simulations of the Classical Nova Outburst. *ApJ*, **692**(2), 1532–1542.
- Starrfield, S., Bose, M., Iliadis, C., Hix, W. R., Woodward, C. E., and Wagner, R. M. (2019). Carbon-Oxygen Classical Novae are Galactic ${}^7\text{Li}$ Producers as well as Potential Supernova Ia Progenitors. *arXiv e-prints*, page arXiv:1910.00575.
- Starrfield, S. G., Sparks, W. M., and Truran, J. W. (1975). Accretion and the nova outburst. *Memoires of the Societe Royale des Sciences de Liege*, **8**, 413–424.
- Stickland, D. J., Penn, C. J., Seaton, M. J., Snijders, M. A. J., and Storey, P. J. (1981). Nova Cygni 1978 - I. The nebular phase. *MNRAS*, **197**, 107–138.
- Subramaniam, A. and Anupama, G. C. (2002). The local stellar population of nova regions in the Large Magellanic Cloud. *A&A*, **390**, 449–471.
- Tajitsu, A., Sadakane, K., Naito, H., Arai, A., and Aoki, W. (2015). Explosive lithium production in the classical nova V339 Del (Nova Delphini 2013). *Nature*, **518**(7539), 381–384.
- Tajitsu, A., Sadakane, K., Naito, H., Arai, A., Kawakita, H., and Aoki, W. (2016). The ${}^7\text{Be}$ II Resonance Lines in Two Classical Novae V5668 Sgr and V2944 Oph. *ApJ*, **818**(2), 191.
- Takei, D., Ness, J.-U., Tsujimoto, M., Kitamoto, S., Drake, J. J., Osborne, J. P., Takahashi, H., and Kinugasa, K. (2011). X-Ray Study of Rekindled Accretion in the Classical Nova V2491 Cygni. *PASJ*, **63**, S729–S738.
- Tappert, C., Vogt, N., Della Valle, M., Schmidtobreick, L., and Ederoclite, A. (2014). Life after eruption - IV. Spectroscopy of 13 old novae. *MNRAS*, **442**(1), 565–576.
- Taylor, A. R., Hjellming, R. M., Seaquist, E. R., and Gehrz, R. D. (1988). Radio images of the expanding ejecta of nova QU Vulpeculae 1984. *Nature*, **335**(6187), 235–238.
- Tomaney, A. B. and Shafter, A. W. (1992). The Spectroscopic and Photometric Evolution of Novae in the Bulge of M31. *ApJS*, **81**, 683.
- Tornambe, A. and Matteucci, F. (1986). Type I SNe from double degenerate CO dwarfs and their rate in the solar neighbourhood. *MNRAS*, **223**, 69–77.
- Townsley, D. M. and Bildsten, L. (2004). Physical Interpretation of Dwarf Nova Primary Effective Temperatures. In G. Tovmassian and E. Sion, editors, *Revista Mexicana de Astronomia y Astrofisica Conference Series*, volume 20 of *Revista Mexicana de Astronomia y Astrofisica Conference Series*, pages 192–193.
- Travaglio, C., Randich, S., Galli, D., Lattanzio, J., Elliott, L. M., Forestini, M., and Ferrini, F. (2001). Galactic Chemical Evolution of Lithium: Interplay between Stellar Sources. *ApJ*, **559**(2), 909–924.
- Truran, J. W. and Livio, M. (1986). On the Frequency of Occurrence of Oxygen-Neon-Magnesium White Dwarfs in Classical Nova Systems. *ApJ*, **308**, 721.
- Vainu Bappu, M. K. and Menzel, D. H. (1954). Emission-Line Profiles from Expanding Envelopes. *ApJ*, **119**, 508.

- van den Bergh, S. (1988). Novae, Supernovae, and the Island Universe Hypothesis. *PASP*, **100**, 8.
- van den Bergh, S. (1991). The Stellar Populations of M31. *PASP*, **103**, 1053.
- van den Bergh, S. (1996). The Extragalactic Distance Scale. *PASP*, **108**, 1091–1096.
- van den Bergh, S. and Younger, P. F. (1987). UVB photometry of novae. *A&AS*, **70**, 125–140.
- Vanlandingham, K. M., Starrfield, S., Wagner, R. M., Shore, S. N., and Sonneborn, G. (1996). Optical and ultraviolet spectrophotometry of the ONeMg Nova V838 Herculis 1991. *MNRAS*, **282**(2), 563–579.
- Vanlandingham, K. M., Starrfield, S., and Shore, S. N. (1997). Elemental abundances for Nova V693 Coronae Austrinae 1981. *MNRAS*, **290**(1), 87–98.
- Vanlandingham, K. M., Schwarz, G. J., Shore, S. N., and Starrfield, S. (2001). Nuclear Turnoff Times for ONEMG Novae Determined Using Ultraviolet Spectral Evolution. *AJ*, **121**(2), 1126–1135.
- Vlasov, A., Vurm, I., and Metzger, B. D. (2016). Shocks in nova outflows - II. Synchrotron radio emission. *MNRAS*, **463**(1), 394–412.
- Vogt, N. (1990). Evidence for Hibernation from a Statistical Analysis of Nova Eruption Amplitudes. *ApJ*, **356**, 609.
- Vorontsov-Velyaminov, B. (1947). The blue-white sequence. *The Observatory*, **67**, 224–226.
- Vurm, I. and Metzger, B. D. (2018). High-energy Emission from Nonrelativistic Radiative Shocks: Application to Gamma-Ray Novae. *ApJ*, **852**(1), 62.
- Wade, C. M. and Hjellming, R. M. (1971). Further Radio Observations of Novae. *ApJ*, **163**, L65.
- Wade, R. A., Harlow, J. J. B., and Ciardullo, R. (2000). Biases in Expansion Distances of Novae Arising from the Prolate Geometry of Nova Shells. *PASP*, **112**(771), 614–624.
- Wagner, R. M., Woodward, C. E., Starrfield, S., Ilyin, I., and Strassmeier, K. (2018). High Resolution Optical Spectroscopy of the Classical Nova V5668 Sgr Showing the Presence of Lithium. In *American Astronomical Society Meeting Abstracts #231*, volume 231 of *American Astronomical Society Meeting Abstracts*, page 358.10.
- Walterbos, R. A. M. (1986). The Relation Between Dust and Gas in the Andromeda Galaxy. In *Bull. Am. Astr. Soc.*, volume 18, page 915.
- Warner, B. (2001). *Cataclysmic Variables: A 'SWOT' Analysis*, volume 246 of *Astronomical Society of the Pacific Conference Series*, page 159.
- Warner, B. (2003). *Cataclysmic Variable Stars*.
- Webbink, R. F. (1984). Double white dwarfs as progenitors of R Coronae Borealis stars and type I supernovae. *ApJ*, **277**, 355–360.
- Welch, D. L. and MACHO Collaboration (1999a). Nova in the Small Magellanic Cloud. *IAU Circ.*, **7121**.
- Welch, D. L. and MACHO Collaboration (1999b). Nova in the Small Magellanic Cloud 1999 No. 4. *IAU Circ.*, **7308**.
- Wendeln, C., Chomiuk, L., Finzell, T., Linford, J. D., and Strader, J. (2017). A Radio Emission Analysis of Classical Nova V351 Pup (1991). *ApJ*, **840**(2), 110.
- Wenzel, W. and Meinunger, I. (1978). Note on the Population Type of Novae. *Astronomische Nachrichten*, **299**, 239.
- Weston, J. H. S., Sokoloski, J. L., Metzger, B. D., Zheng, Y., Chomiuk, L., Krauss, M. I., Linford, J. D., Nelson, T., Mioduszewski, A. J., Rupen, M. P., Finzell, T., and Mukai, K. (2016a). Non-thermal radio emission from colliding flows in classical nova V1723 Aql. *MNRAS*, **457**(1), 887–901.
- Weston, J. H. S., Sokoloski, J. L., Chomiuk, L., Linford, J. D., Nelson, T., Mukai, K., Finzell, T., Mioduszewski, A., Rupen, M. P., and Walter, F. M. (2016b). Shock-powered radio emission from V5589 Sagittarii (Nova Sgr 2012 #1). *MNRAS*, **460**(3), 2687–2697.
- Whelan, J. and Iben, Icko, J. (1973). Binaries and Supernovae of Type I. *ApJ*, **186**, 1007–1014.
- Wickramasinghe, D. T., Tout, C. A., and Ferrario, L. (2014). The most magnetic stars. *MNRAS*, **437**(1), 675–681.
- Williams, R. (2012). Origin of the “He/N” and “Fe II” Spectral Classes of Novae. *AJ*, **144**(4), 98.
- Williams, R. (2016). Transient classification and novae ejecta. In *Journal of Physics Conference Series*, volume 728 of *Journal of Physics Conference Series*, page 042001.
- Williams, R., Mason, E., Della Valle, M., and Ederoclite, A. (2008). Transient Heavy Element Absorption Systems in Novae: Episodic Mass Ejection from the Secondary Star. *ApJ*, **685**(1), 451–462.

- Williams, R. E. (1977). Element Abundance Analyses of Novae. In R. Kippenhahn, J. Rahe, and W. Strohmeier, editors, *IAU Colloq. 42: The Interaction of Variable Stars with their Environment*, page 242.
- Williams, R. E. (1985). CNO abundances in novae ejecta. In *European Southern Observatory Conference and Workshop Proceedings*, volume 21 of *European Southern Observatory Conference and Workshop Proceedings*, pages 225–232.
- Williams, R. E. (1992). The Formation of Novae Spectra. *AJ*, **104**, 725.
- Williams, R. E. (1994). Extinction, Ejecta Masses, and Radial Velocities of Novae. *ApJ*, **426**, 279.
- Williams, R. E., Hamuy, M., Phillips, M. M., Heathcote, S. R., Wells, L., and Navarrete, M. (1991). The Evolution and Classification of Postoutburst Novae Spectra. *ApJ*, **376**, 721.
- Williams, R. E., Phillips, M. M., and Hamuy, M. (1994). The Tololo Nova Survey: Spectra of Recent Novae. *ApJS*, **90**, 297.
- Williams, S. J. and Shafter, A. W. (2004). On the Nova Rate in M33. *ApJ*, **612**(2), 867–876.
- Woodward, C. E. and Starrfield, S. (2011). Recent observational and theoretical studies of the classical nova outburst. *Canadian Journal of Physics*, **89**(4), 333–343.
- Woodward, C. E., Gehrz, R. D., Jones, T. J., and Lawrence, G. F. (1992). The Peculiar, Fast Nova Herculis 1991. *ApJ*, **384**, L41.
- Woodward, C. E., Greenhouse, M. A., Gehrz, R. D., Pendleton, Y. J., Joyce, R. R., van Buren, D., Fischer, J., Jennerjohn, N. J., and Kaminski, C. D. (1995). The Temporal Evolution of the 1–5 Micron Spectrum of V1974 Cygni (Nova Cygni 1992). *ApJ*, **438**, 921.
- Woodward, C. E., Gehrz, R. D., Jones, T. J., Lawrence, G. F., and Skrutskie, M. F. (1997). The Temporal Evolution of the Near-Infrared Light Curves of V1974 Cygni (Nova Cygni 1992). *ApJ*, **477**(2), 817–824.
- Woudt, P. A., Warner, B., and Pretorius, M. L. (2004). High-speed photometry of faint cataclysmic variables - IV. V356 Aql, Aqr1, FIRST J1023+0038, H α 0242-2802, GI Mon, AO Oct, V972 Oph, SDSS 0155+00, SDSS 0233+00, SDSS 1240-01, SDSS 1556-00, SDSS 2050-05, FH Ser. *MNRAS*, **351**(3), 1015–1025.
- Wygoda, N., Elbaz, Y., & Katz, B. 2019, *MNRAS*, 484, 3941
- Yaron, O., Prialnik, D., Shara, M. M., and Kovetz, A. (2005). An Extended Grid of Nova Models. II. The Parameter Space of Nova Outbursts. *ApJ*, **623**(1), 398–410.
- Yungelson, L. and Livio, M. (1998). Type IA Supernovae: an Examination of Potential Progenitors and the Redshift Distribution. *ApJ*, **497**(1), 168–177.
- Yungelson, L., Livio, M., and Tutukov, A. (1997). On The Rate of Novae in Galaxies of Different Types. *ApJ*, **481**(1), 127–131.
- Zwicky, F. (1936). Life-Luminosity Relation for Novae. *PASP*, **48**(284), 191.

A Tables

Nova	Distance (<i>kpc</i>)	V_{max} (<i>mag</i>)	V_{min} (<i>mag</i>)	t_2 (<i>days</i>)	t_3 (<i>days</i>)	l (<i>deg</i>)	b (<i>deg</i>)
V1494 Aql	1.239±0.422	4.1	17.1	8	16	40.97	-4.742
V705 Cas	2.157±0.799	5.7	16.4	33	67	113.7	-4.096
V842 Cen	1.379±0.12	4.9	15.8	43	48	316.6	2.452
V476 Cyg	0.665±0.107	1.9	16.2	6	16	87.37	12.42
V1974 Cyg	1.631±0.261	4.3	21.0	19	43	89.13	7.819
T Pyx	3.185±0.607	6.4	18.5	32	62	257.2	9.707
V732 Sgr	1.795±0.458	6.4	16.0	65	75	2.53	-1.188
FH Ser	1.06±0.112	4.5	16.8	49	62	32.91	5.786
V382 Vel	1.8±0.243	2.8	16.6	6	13	284.2	5.771
NQ Vul	1.08±0.169	6.2	17.2	21	50	55.36	1.29
PW Vul	2.42±1.337	6.4	16.9	44	116	61.1	5.197
V603 Aql	0.311±0.007	-1.4	11.7	5	9	33.16	0.829
T Aur	0.857±0.038	4.2	15.2	80	100	177.14	-1.7
Q Cyg	1.32±0.043	3.0	14.9	5	11	89.93	-7.552
HR Del	0.932±0.031	3.8	12.1	160	230	63.43	-13.97
DN Gem	1.318±0.146	3.5	15.5	17	37	184.02	14.71
DQ Her	0.494±0.006	1.35	14.3	67	94	73.15	26.44
V446 Her	1.308±0.13	3.0	16.9	6	15	45.409	4.708
V533 Her	1.165±0.044	3.0	14.5	25	44	69.19	24.273
CP Lac	1.129±0.054	2.1	15.5	5	10	102.14	-0.837
DI Lac	1.569±0.051	4.6	14.7	20	40	103.107	-4.855
DK Lac	2.296±0.391	5.7	16.8	40	60	105.237	-5.352
HR Lyr	4.493±0.684	6.5	15.5	45	80	59.584	12.47
GI Mon	2.849±0.46	5.4	15.8	20	30	222.93	4.749
V841 Oph	0.805±0.018	4.2	13.5	54	130	7.621	17.779
GK Per	0.437±0.008	0.2	13.4	6	13	150.956	-10.104
RR Pic	0.504±0.008	2.0	12.0	125	250	272.355	-25.672
CP Pup	0.795±0.013	0.4	15.0	5	8	252.926	-0.835
V1500 Cyg	1.287±0.309	1.9	20.5	2	5	89.823	-0.073

Table 9 The list of novae (Col. 1), their distance as measured from GAIA DR2 (Col. 2), the V magnitude at maximum (Col. 3) and at minimum (Col 4), the corresponding t_2 and t_3 values (Cols. 5 and 6) and the Galactic longitude l and latitude b (Cols. 7 and 8). Data from S18 and SG19

Nova	A_V (mag)	$E(B - V)^*$ (mag)
CI Aql	2.6	1.0
V1494 Aql	1.9	-
V705 Cas	1.3	0.45
V1330 Cyg	2.1	-
DN Gem	0.5	-
V446 Her	1.1	0.40
T Pyx	0.8	-
V732 Sgr	2.5	-
V382 Vel	0.4	0.25
Q Cyg	0.81	-
DI Lac	0.81	-
HR Lyr	0.56	-
GI Mon	0.31	-
V841 Oph	1.36	-
CP Pup	0.79	-
CT Ser	0.02	-
HR Del	0.47	-
NQ Vul	2.1	-
PW Vul	1.73	0.57
QU Vul	1.89	0.55
V842 Cen	1.73	-
QV Vul	1.26	-
V1974 Cyg	1.12	0.30
T Aur	0.64	-
GK Per	0.95	-
V603 Aql	0.22	-
V476 Cyg	0.82	-
RR Pic	0.16	-
DQ Her	0.32	-
CP Lac	0.64	-
DK Lac	1.4	-
V446 Her	1.12	0.40
V533 Her	0.0	0.038
V1229 Aql	1.58	-
FH Ser	2.02	-

Table 10 The list of absorptions A_V used in this work. Data from Downes & Duerbeck (2000), S18 and SG19. * The third column shows the extinction as provided by Hachisu and Kato (2019).

Nova	t_2 (days)	FWHM (km/s)	Type	Nova	t_2 (days)	FWHM (km/s)	Type
V356 Aql	127	1140	Fe II	V533 Her	26	1600	Fe II
V500 Aql	20	2800	He/N	V838 Her	2	3300	He/N
V603 Aql	4	4000	He/N	CP Lac	5	4200	He/N
V1229 Aql	18	2200	Fe II	DK Lac	19	2500	Fe II
v1425 Aql	27	1340	Fe II	V382 Nor	12	1850	Fe II
V1494 Aql	8	2470	Fe II	V2264 Oph	18	2300	Fe II
V1663 Aql	16	1900	Fe II	V2615 Oph	26.5	800	Fe II
V1974 Cyg	23	2300	Fe II	V2672 Oph	2.3	8000	He/N
T Aur	80	1200	Fe II	GK Per	6	4000	He/N
V723 Cas	263	600	Fe II	RR Pic	80	1600	Fe II
V842 Cen	43	760	Fe II	CP Pup	5	2000	He/N
V868 Cen	18	2300	Fe II	V351 Pup	10	2500	Fe II
V1065 Cen	11	2700	Fe II	V745 Sco	7	2000	He/N
V407 Cyg	5.9	2300	He/N	V1187 Sco	10	3000	Fe II
V476 Cyg	7	2500.0	Fe II	V1280 Sco	21.3	640	Fe II
V1500 Cyg	2.4	5000	He/N	V1324 Sco	25	2000	Fe II
V1668 Cyg	11	2400	Fe II	V496 Sct	59	1200	Fe II
V1819 Cyg	95	2742	Fe II	FH Ser	42	2100	Fe II
V2467 Cyg	8	950	Fe II	V3890 Sgr	10	2500	He/N
V2468 Cyg	9	2300	Fe II-b	V5558 Sgr	281	1000	Fe II-b
V2491 Cyg	4	4860	He/N	V382 Vel	6	2400	He/N
HR Del	150	1700	Fe II	LV Vul	21	2700	Fe II
DQ Her	67	1000	Fe II	QU Vul	22	1700	Fe II
V446 Her	5	2400	He/N	QV Vul	37	920	Fe II

Table 11 The average expansion velocity (Cols. 3 & 7) for the novae shown in Fig. 28. In the table are also reported the corresponding t_2 value (Col. 2 & 6) and the spectral type (Col. 4 & 8) for each nova considered. Data from Ozdonmez et al. (2018) and Della Valle & Livio (1998).

Nova	Nebular distance (kpc)	GAIA DR2 distance (kpc)
CP Pup	1.70	0.81
CT Ser	1.40	2.77
HR Del	0.76	0.96
NQ Vul	1.16	1.08
PW Vul	1.80	2.42
QU Vul	1.75	1.79
V842 Cen	1.15	1.38
QV Vul	2.70	3.62
V1974 Cyg	2.00	1.63
T Aur	0.96	0.88
GK Per	0.45	0.44
V603 Aql	0.33	0.31
V476 Cyg	1.62	0.66
RR Pic	0.58	0.51
DQ Her	0.48	0.50
CP Lac	1.35	1.17
DK Lac	3.90	2.52
V446 Her	1.35	1.36
V533 Her	1.25	1.20
V1229 Aql	2.10	2.78
FH Ser	0.95	1.06

Table 12 The distance measured from the nebular parallax method (Col. 2) by Downes & Duerbeck (2000) compared with the recent GAIA DR2 measurements (Col. 3) as given in S18 and SG19, for a sample of novae (Col. 1).

Nova	t_2 (days)	m_{pg} (mag)	σm_{pg} (mag)	Nova	t_2 (days)	m_{pg} (mag)	σm_{pg} (mag)
A01	2.0	15.73	0.2	R24	10.0	16.1	0.3
A02	2.0	15.67	0.2	R27	28.6	16.9	0.3
A03	5.3	15.9	0.3	R28	8.0	14.85	0.1
A04	10.0	18.2	0.1	R29	6.7	16.0	0.3
A05	11.8	15.9	0.2	R30	11.8	16.2	0.1
A06	8.7	16.0	0.1	R32	4.0	15.75	0.3
A07	13.3	15.9	0.1	R33	8.0	16.2	0.2
A08	8.7	16.0	0.2	R34	16.7	17.0	0.3
A09	11.5	16.0	0.3	R36	80.0	16.9	0.3
A10	6.9	16.0	0.2	R37	18.2	16.8	0.2
A12	11.1	16.1	0.1	R38	18.2	16.9	0.1
A13	26.0	17.0	0.1	R41	100.0	17.9	0.2
A14	12.3	16.2	0.2	R42	28.6	16.85	0.2
A15	15.9	16.4	0.2	R43	9.1	16.3	0.1
A16	12.8	16.7	0.2	R46	16.0	16.6	0.2
A17	29.0	17.2	0.2	R48	11.0	17.8	0.1
A18	34.5	17.5	0.2	R49	22.2	16.35	0.3
A19	28.6	17.6	0.1	R52	16.0	16.35	0.1
A20	33.3	17.2	0.1	R53	50.0	16.95	0.1
A21	26.7	17.4	0.1	R55	29.9	17.0	0.3
A22	29.9	17.6	0.2	R57	9.1	14.95	0.1
A23	43.5	17.4	0.2	R58	16.0	16.5	0.3
A24	33.9	17.8	0.1	R59	20.0	17.3	0.3
A25	32.8	17.6	0.1	R60	15.4	16.4	0.3
A26	46.5	18.0	0.1	R66	1.5	16.4	0.2
A27	83.3	17.8	0.3	R67	15.4	16.2	0.1
A28	105.3	17.8	0.2	R76	14.3	16.1	0.2
A29	117.6	18.0	0.1	R77	9.1	15.9	0.1
A30	117.6	18.1	0.2	R78	25.0	16.9	0.3
R04	15.4	16.4	0.2	R80	58.8	17.4	0.1
R05	3.3	15.6	0.2	R85	16.7	15.7	0.1
R06	14.3	16.5	0.1	R86	33.3	17.3	0.3
R07	5.0	16.2	0.2	R89	20.0	16.8	0.3
R09	13.3	16.8	0.3	R95	6.7	16.2	0.3
R12	40.0	17.6	0.1	R98	9.5	16.0	0.3
R13	16.7	16.8	0.3	R100	10.0	16.0	0.3
R14	25.0	17.3	0.3	E104	50.0	18.4	0.3
R15	18.2	16.3	0.3	E105	15.4	16.9	0.3
R16	12.5	16.25	0.3	E106	44.4	17.6	0.3
R17	11.8	16.7	0.3	E109	10.0	16.3	0.2
R18	22.2	17.1	0.1	E112	20.0	17.3	0.2
R19	80.0	17.9	0.2	E113	5.0	16.0	0.3
R20	22.2	16.9	0.1	E117	100.0	17.9	0.3
R21	22.2	16.7	0.3	E118	28.6	16.9	0.2
R23	13.3	16.85	0.3	E120	40.0	17.4	0.2
-	-	-	-	E129	22.2	16.8	0.2

Table 13 The list of M31 novae used in this work. Data from Arp (1956), Rosino (1964), Rosino (1973) and Rosino et al. (1989) . In this table novae are listed according to the original papers. t_2 values are reported in Cols. 2 & 6. The photographic magnitudes m_{pg} at peak and errors are reported in Cols. 3 & 7 and 4 & 8, respectively.

Nova	t_2 (<i>days</i>)	V_{obs} (<i>mag</i>)	σV_{obs} (<i>mag</i>)
LMC1926	117.6	12.0	0.3
LMC1935	13.8	11.0	0.3
LMC1936	17.7	10.5	0.3
LMC1937	10.7	10.6	0.3
LMC1948	58.8	12.82	0.3
LMC1951	2.6	11.9	0.0
LMC1968	2.0	10.4	0.4
LMC1970B	8.0	11.0	0.5
LMC1971A	15.7	11.77	0.1
LMC1977B	11.2	10.7	0.1
LMC1978A	3.5	9.75	0.6
LMC1987	2.0	9.6	0.2
LMC1988A	22.2	10.85	0.1
LMC1988B	4.7	9.9	0.2
LMC1990A	3.4	9.5	0.3
LMC1990B	3.4	10.2	0.1
LMC1991	5.7	9.0	0.1
LMC1992	8.0	10.2	0.1
LMC1995	11.8	10.4	0.3
LMC2000	10.0	10.7	0.1
LMC2002	13.3	10.1	0.2
LMC2003	8.0	11.0	0.1
LMC2004	11.8	10.9	0.1
LMC2005	62.5	11.5	0.1
LMC2009a	5.4	10.6	0.1
LMC2009b	54.1	12.1	0.1
LMC2012b	10.0	11.5	0.1

Table 14 The list of LMC novae used in this work. Data from Capaccioli et al. (1990), Shafter (2013). t_2 values, magnitudes at maximum and errors are reported in Col. 2, Col. 3 and 4, respectively.

Nova	t_2 (days)	$m_{V,peak}$ (mag)
M31-2007-10a	8.6	17.5
M31-2008-07b	17.0	19.0
M31-2008-8c	26.3	17.1
M31-2008-09a	25.0	17.8
M31-2008-09c	16.6	16.6
M31-2008-10b	35.0	18.1
M31-2008-11a	7.5	17.0
M31-2008-12b	25.0	17.0
M31-2009-08b	45.0	17.86
M31-2009-09a	79.0	17.36
M31-2009-10a	20.0	17.05
M31-2009-10b	8.0	14.94
M31-2009-11a	22.0	17.33
M31-2009-11b	90.0	18.35
M31-2009-11d	25.0	17.82
M31-2009-12a	10.0	16.98
M31-2010-06a	40.0	16.98
M31-2010-06b	8.0	17.72
M31-2010-06c	19.0	17.57
M31-2010-07a	15.0	15.75
M31-2010-09a	40.0	17.52
M31-2010-09b	10.0	16.46
M31-2010-10b	57.0	17.14
M31-2010-10a	15.0	17.13
M31-2010-10c	60.0	17.54

Table 15 The list of M31 novae taken from the analysis of Kasliwal *et al.* (2011) and Cao *et al.* (2012). t_2 values and m_V magnitudes are reported in Col. 2 and Col. 3, respectively.

Nova	t_2 (days)	$m_{pg,peak}$ (mag)
M31-Hubble-nova-1	100.0	16.7
M31-Hubble-nova-2	22.2	15.8
M31-Hubble-nova-3	16.7	15.8
M31-Hubble-nova-4	80.0	17.1
M31-Hubble-nova-5	33.3	16.2
M31-Hubble-nova-6	66.7	16.9
M31-Hubble-nova-7	15.4	16.2
M31-Hubble-nova-8	100.0	17.5
M31-Hubble-nova-9	28.6	16.8
M31-Hubble-nova-10	20.0	16.3
M31-Hubble-nova-11	25.0	16.4
M31-Hubble-nova-12	11.1	16.3
M31-Hubble-nova-13	36.4	17.3
M31-Hubble-nova-14	28.6	17.2
M31-Hubble-nova-15	15.4	15.3
M31-Hubble-nova-16	33.3	17.4
M31-Hubble-nova-17	133.3	17.5

Table 16 The list of M31 novae taken from the analysis of Hubble (1929). t_2 values and m_{pg} magnitudes are reported in Col. 2 and Col. 3, respectively.

Nova	t_2 (<i>days</i>)	V_{peak} (<i>mag</i>)	σV_{peak} (<i>mag</i>)
NGC4365	18.2	23.97	0.30
M49-1W	20.0	22.51	0.11
M49-2W	12.6	21.01	0.57
M49-3W	54.0	24.13	0.21
M49-5W	10.5	23.81	0.12
M49-2E	33.2	23.11	0.10
M49-4	52.0	23.33	0.13
M49-5	7.5	23.10	0.52
M49-6	19.3	23.33	0.28
M49-7	11.7	22.72	0.36
M49-9	37.2	23.74	0.22
M87-1	15.2	21.84	-
M87-2	11.2	22.25	-
M87-3	2.0	22.64	-
M87-4	7.7	22.74	-
M87-5	17.1	22.99	-
M87-6	11.2	23.18	-
M87-7	9.3	23.21	-
M87-8	3.7	23.37	-
M87-10	22.3	23.51	-
M87-11	28.9	23.57	-
M87-12	33.1	23.58	-
M87-13	6.7	23.67	-
M87-14	31.5	23.74	-
M87-16	32.6	23.77	-
M87-17	30.4	23.81	-
M87-19	9.0	23.90	-
M87-20	36.3	23.94	-
M87-21	29.9	24.07	-
M87-23	8.0	24.16	-

Table 17 The list of VIRGO novae used in this work. Data are from Pritchett & van den Bergh (1987), Ferrarese et al. (2003) and Shara et al. (2016). t_2 values, magnitudes at maximum and errors are reported in Col. 2, Col. 3 and 4, respectively. B_{peak} mags have been transformed into V_{peak} mags by applying $\langle B - V \rangle = 0.23 \pm 0.06$ mag (van den Bergh & Younger 1987).