

# Planar Graphs Have Bounded Queue-Number

Vida Dujmović<sup>1</sup>   Gwenaël Joret<sup>2</sup>   Piotr Micek<sup>3</sup>   Pat Morin<sup>4</sup>  
Torsten Ueckerdt<sup>5</sup>   David R. Wood<sup>6</sup>

<sup>1</sup>University of Ottawa

<sup>2</sup>Université Libre de Bruxelles

<sup>3</sup>Jagiellonian University


<sup>4</sup>Carleton University

<sup>5</sup>Karlsruhe Institute of Technology

<sup>6</sup>Monash University


Geometry: Combinatorics and Algorithms Seminar  
Shengzhe Wang  
April 21, 2023

# Queue Layout




A path graph with 6 vertices


# Queue Layout



A path graph with 6 vertices





# Queue Layout





Not Valid Queue-Layout


# Queue Layout




# Queue Layout



# Queue Layout



# Queue Layout






# Queue Layout

## Definition: Queue

Let  $G = (V, E)$ , consider a linear ordering  $\preceq$  of  $V$ , a queue of  $G$  is a set of edges  $E' \subseteq E$  such that any disjoint edges  $vw, xy \in E'$ , w.l.o.g,  $v \prec w, x \prec y$  and  $v \prec x$ , we have  $w \prec y$ .



## Definition: K-Queue Layout

Let  $G = (V, E)$ , consider a linear ordering  $\preceq$  of  $V$ , for an integer  $k \geq 0$  a  $k$ -queue layout of  $G$  is a partition of  $E$  into  $E_1, E_2, \dots, E_k$  such that each  $E_i$  is a queue of  $G$ .

# Queue-Number

## Definition: K-Queue Layout


Let  $G = (V, E)$ , consider a linear ordering  $\preceq$  of  $V$ , for an integer  $k \geq 0$  a  $k$ -queue layout of  $G$  is a partition of  $E$  into  $E_1, E_2, \dots, E_k$  such that each  $E_i$  is a queue of  $G$ .

## Definition: Queue-Number

The queue-number of  $G$ , denoted by  $qn(G)$ , is the minimum integer  $k$  such that  $G$  has a  $k$ -queue layout.


# Queue-Number: Tree

What is the queue-number of tree?




# Queue-Number: Tree

What is the queue-number of tree?





# Queue-Number: Tree





What is the queue-number of tree?



# Queue Number: Cycle








# Queue Number: Cycle











# Queue Number: Cycle







# Queue Number: $K_4$



# Queue Number: $K_4$



# Queue Number: $K_4$



Theorem (Heath, Rosenberg, 1992)

*The complete graph  $K_n$  has queue number  $\lfloor \frac{n}{2} \rfloor$ .*

Do we have some tools to help bound the queue-number?

Do we have some tools to help bound the queue-number?

Theorem (Wiechert, 2017)


*Every graph with treewidth  $k$  has queue-number at most  $2^k - 1$ .*

# Tree-decomposition

## Definition: Tree-decomposition

A tree-decomposition of a graph  $G$  is a pair  $(B, T)$ .  $T$  is a tree and  $B = \{B_x | x \in V(T)\}$  where each  $B_x$  is a subset of  $V(G)$  for every vertex  $x$  in  $V(T)$  such that

- $\forall \{v, w\} \in E(G)$ , there exists  $x \in V(T)$  with  $v, w \in B_x$
- $\forall v \in V(G)$ , the set  $\{x | x \in V(T) \wedge v \in B_x\}$  induces a non-empty connected subtree of  $T$ .




# Tree-decomposition


## Definition: Tree-decomposition

A tree-decomposition of a graph  $G$  is a pair  $(B, T)$ .  $T$  is a tree and  $B = \{B_x | x \in V(T)\}$  where each  $B_x$  is a subset of  $V(G)$  for every vertex  $x$  in  $V(T)$  such that

- $\forall \{v, w\} \in E(G)$ , there exists  $x \in V(T)$  with  $v, w \in B_x$
- $\forall v \in V(G)$ , the set  $\{x | x \in V(T) \wedge v \in B_x\}$  induces a non-empty connected subtree of  $T$ .



$G$



$T$




# Tree-decomposition

## Definition: Tree-decomposition

A tree-decomposition of a graph  $G$  is a pair  $(B, T)$ .  $T$  is a tree and  $B = \{B_x | x \in V(T)\}$  where each  $B_x$  is a subset of  $V(G)$  for every vertex  $x$  in  $V(T)$  such that

- $\forall \{v, w\} \in E(G)$ , there exists  $x \in V(T)$  with  $v, w \in B_x$
- $\forall v \in V(G)$ , the set  $\{x | x \in V(T) \wedge v \in B_x\}$  induces a non-empty connected subtree of  $T$ .




# Tree-decomposition

## Definition: Tree-decomposition

A tree-decomposition of a graph  $G$  is a pair  $(B, T)$ .  $T$  is a tree and  $B = \{B_x | x \in V(T)\}$  where each  $B_x$  is a subset of  $V(G)$  for every vertex  $x$  in  $V(T)$  such that

- $\forall \{v, w\} \in E(G)$ , there exists  $x \in V(T)$  with  $v, w \in B_x$
- $\forall v \in V(G)$ , the set  $\{x | x \in V(T) \wedge v \in B_x\}$  induces a non-empty connected subtree of  $T$ .




# Tree-decomposition

## Definition: Tree-decomposition

A tree-decomposition of a graph  $G$  is a pair  $(B, T)$ .  $T$  is a tree and  $B = \{B_x | x \in V(T)\}$  where each  $B_x$  is a subset of  $V(G)$  for every vertex  $x$  in  $V(T)$  such that

- $\forall \{v, w\} \in E(G)$ , there exists  $x \in V(T)$  with  $v, w \in B_x$
- $\forall v \in V(G)$ , the set  $\{x | x \in V(T) \wedge v \in B_x\}$  induces a non-empty connected subtree of  $T$ .




# Tree-decomposition

## Definition: Tree-decomposition

A tree-decomposition of a graph  $G$  is a pair  $(B, T)$ .  $T$  is a tree and  $B = \{B_x | x \in V(T)\}$  where each  $B_x$  is a subset of  $V(G)$  for every vertex  $x$  in  $V(T)$  such that

- $\forall \{v, w\} \in E(G)$ , there exists  $x \in V(T)$  with  $v, w \in B_x$
- $\forall v \in V(G)$ , the set  $\{x | x \in V(T) \wedge v \in B_x\}$  induces a non-empty connected subtree of  $T$ .




# Tree-decomposition

## Definition: Tree-decomposition

A tree-decomposition of a graph  $G$  is a pair  $(B, T)$ .  $T$  is a tree and  $B = \{B_x | x \in V(T)\}$  where each  $B_x$  is a subset of  $V(G)$  for every vertex  $x$  in  $V(T)$  such that

- $\forall \{v, w\} \in E(G)$ , there exists  $x \in V(T)$  with  $v, w \in B_x$
- $\forall v \in V(G)$ , the set  $\{x | x \in V(T) \wedge v \in B_x\}$  induces a non-empty connected subtree of  $T$ .




# Tree-decomposition

## Definition: Tree-decomposition

A tree-decomposition of a graph  $G$  is a pair  $(B, T)$ .  $T$  is a tree and  $B = \{B_x | x \in V(T)\}$  where each  $B_x$  is a subset of  $V(G)$  for every vertex  $x$  in  $V(T)$  such that

- $\forall \{v, w\} \in E(G)$ , there exists  $x \in V(T)$  with  $v, w \in B_x$
- $\forall v \in V(G)$ , the set  $\{x | x \in V(T) \wedge v \in B_x\}$  induces a non-empty connected subtree of  $T$ .



## Definition: Tree-decomposition

A tree-decomposition of a graph  $G$  is a pair  $(B, T)$ .  $T$  is a tree and  $B = \{B_x | x \in V(T)\}$  where each  $B_x$  is a subset of  $V(G)$  for every vertex  $x$  in  $V(T)$  such that

- $\forall \{v, w\} \in E(G)$ , there exists  $x \in V(T)$  with  $v, w \in B_x$
- $\forall v \in V(G)$ , the set  $\{x | x \in V(T) \wedge v \in B_x\}$  induces a non-empty connected subtree of  $T$ .

## Definition: Width of Tree-decomposition

The width of a tree-decomposition of  $G$  is  $\max_{x \in V(T)} |B_x| - 1$

# Treewidth

## Definition: Tree-decomposition

A tree-decomposition of a graph  $G$  is a pair  $(B, T)$ .  $T$  is a tree and  $B = \{B_x | x \in V(T)\}$  where each  $B_x$  is a subset of  $V(G)$  for every vertex  $x$  in  $V(T)$  such that

- $\forall \{v, w\} \in E(G)$ , there exists  $x \in V(T)$  with  $v, w \in B_x$
- $\forall v \in V(G)$ , the set  $\{x | x \in V(T) \wedge v \in B_x\}$  induces a non-empty connected subtree of  $T$ .

## Definition: Width of Tree-decomposition

The width of a tree-decomposition of  $G$  is  $\max_{x \in V(T)} |B_x| - 1$

## Definition: Treewidth

The treewidth of a graph  $G$  is the minimum width of all tree-decompositions of  $G$ .



## Treewidth: Fixed-Parameter Tractability

- Computing a maximum independent set in a graph  $G$  is NP-hard.


## Treewidth: Fixed-Parameter Tractability

- Computing a maximum independent set in a graph  $G$  is NP-hard.
- If  $G$  has treewidth  $\leq k$ , then a maximum independent set in  $G$  can be computed in time  $O(k^2 4^k |V|)$ .


# Treewidth: Fixed-Parameter Tractability

- Computing a maximum independent set in a graph  $G$  is NP-hard.
- If  $G$  has treewidth  $\leq k$ , then a maximum independent set in  $G$  can be computed in time  $O(k^2 4^k |V|)$ .
- Dynamic Programming on trees is relatively fast.



# Treewidth: Tree





# Treewidth: Tree





# Treewidth: Tree



# Treewidth: Tree



# Treewidth: Tree



Tree has treewidth 1



# Treewidth: Planar graph

## Theorem (Wiechert, 2017)

*Every graph with treewidth  $k$  has queue-number at most  $2^k - 1$ .*

If planar graph has bounded treewidth, then planar graph has bounded queue-number.

# Treewidth: Planar graph

## Theorem (Wiechert, 2017)

*Every graph with treewidth  $k$  has queue-number at most  $2^k - 1$ .*

If planar graph has bounded treewidth, then planar graph has bounded queue-number.

## Theorem (Alon, Seymour, Thomas, 1990)

*Planar graph on  $n$  vertices has treewidth  $O(\sqrt{n})$ .*

# Treewidth: Planar graph

## Theorem (Wiechert, 2017)

*Every graph with treewidth  $k$  has queue-number at most  $2^k - 1$ .*


If planar graph has bounded treewidth, then planar graph has bounded queue-number.

## Theorem (Alon, Seymour, Thomas, 1990)

*Planar graph on  $n$  vertices has treewidth  $O(\sqrt{n})$ .*

## Theorem (Robertson, Seymour)

*A grid graph with size  $n \times n$  has treewidth  $n$ .*



# Partitions

Maybe we need more structure

Maybe we need more structure

**Theorem (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)**

*For a graph  $G$ , if  $G$  has an  $H$ -partition of layered width  $\ell$  and  $H$  has treewidth  $k$ , then*

$$qn(G) \leq 3l(2^k - 1) + \lfloor \frac{3}{2}\ell \rfloor$$

## Definition: Partition and Quotient

A partition of  $G$  is a set  $\mathcal{P} = \{P_1, \dots, P_n\}$  of non-empty subsets of  $V(G)$  and each vertex of  $G$  is in exactly one element (part) of  $\mathcal{P}$ .


The quotient of  $\mathcal{P}$  is a graph, denoted by  $G/\mathcal{P}$ , with each vertex  $v_i$  corresponds  $P_i$ . For any two vertices  $v_i, v_j$  in  $G/\mathcal{P}$ , they are connected if and only if some vertex in  $P_i$  is connected to some vertex in  $P_j$  in graph  $G$ .

# Partitions

## Definition: Partition and Quotient

A partition of  $G$  is a set  $\mathcal{P} = \{P_1, \dots, P_n\}$  of non-empty subsets of  $V(G)$  and each vertex of  $G$  is in exactly one element (part) of  $\mathcal{P}$ .

The quotient of  $\mathcal{P}$  is a graph, denoted by  $G/\mathcal{P}$ , with each vertex  $v_i$  corresponds  $P_i$ . For any two vertices  $v_i, v_j$  in  $G/\mathcal{P}$ , they are connected if and only if some vertex in  $P_i$  is connected to some vertex in  $P_j$  in graph  $G$ .




# Partitions

## Definition: Partition and Quotient

A partition of  $G$  is a set  $\mathcal{P} = \{P_1, \dots, P_n\}$  of non-empty subsets of  $V(G)$  and each vertex of  $G$  is in exactly one element (part) of  $\mathcal{P}$ .

The quotient of  $\mathcal{P}$  is a graph, denoted by  $G/\mathcal{P}$ , with each vertex  $v_i$  corresponds  $P_i$ . For any two vertices  $v_i, v_j$  in  $G/\mathcal{P}$ , they are connected if and only if some vertex in  $P_i$  is connected to some vertex in  $P_j$  in graph  $G$ .






# Partitions

## Definition: Partition and Quotient

A partition of  $G$  is a set  $\mathcal{P} = \{P_1, \dots, P_n\}$  of non-empty subsets of  $V(G)$  and each vertex of  $G$  is in exactly one element (part) of  $\mathcal{P}$ .

The quotient of  $\mathcal{P}$  is a graph, denoted by  $G/\mathcal{P}$ , with each vertex  $v_i$  corresponds  $P_i$ . For any two vertices  $v_i, v_j$  in  $G/\mathcal{P}$ , they are connected if and only if some vertex in  $P_i$  is connected to some vertex in  $P_j$  in graph  $G$ .



# Partitions


## Definition: Partition and Quotient

A partition of  $G$  is a set  $\mathcal{P} = \{P_1, \dots, P_n\}$  of non-empty subsets of  $V(G)$  and each vertex of  $G$  is in exactly one element (part) of  $\mathcal{P}$ .

The quotient of  $\mathcal{P}$  is a graph, denoted by  $G/\mathcal{P}$ , with each vertex  $v_i$  corresponds  $P_i$ . For any two vertices  $v_i, v_j$  in  $G/\mathcal{P}$ , they are connected if and only if some vertex in  $P_i$  is connected to some vertex in  $P_j$  in graph  $G$ .

## Definition: $H$ -partition

A  $H$ -partition of a graph  $G$  is a pair  $(A, H)$ .  $A = \{A_x | x \in V(H)\}$  is a partition of  $V(G)$  and  $H$  is a graph isomorphic to the quotient  $G/A$ .



# Partitions


## Definition: Partition and Quotient

A partition of  $G$  is a set  $\mathcal{P} = \{P_1, \dots, P_n\}$  of non-empty subsets of  $V(G)$  and each vertex of  $G$  is in exactly one element (part) of  $\mathcal{P}$ .

The quotient of  $\mathcal{P}$  is a graph, denoted by  $G/\mathcal{P}$ , with each vertex  $v_i$  corresponds  $P_i$ . For any two vertices  $v_i, v_j$  in  $G/\mathcal{P}$ , they are connected if and only if some vertex in  $P_i$  is connected to some vertex in  $P_j$  in graph  $G$ .

## Definition: $H$ -partition

A  $H$ -partition of a graph  $G$  is a pair  $(A, H)$ .  $A = \{A_x | x \in V(H)\}$  is a partition of  $V(G)$  and  $H$  is a graph isomorphic to the quotient  $G/A$ .



# Partitions


## Definition: Partition and Quotient

A partition of  $G$  is a set  $\mathcal{P} = \{P_1, \dots, P_n\}$  of non-empty subsets of  $V(G)$  and each vertex of  $G$  is in exactly one element (part) of  $\mathcal{P}$ .


The quotient of  $\mathcal{P}$  is a graph, denoted by  $G/\mathcal{P}$ , with each vertex  $v_i$  corresponds  $P_i$ . For any two vertices  $v_i, v_j$  in  $G/\mathcal{P}$ , they are connected if and only if some vertex in  $P_i$  is connected to some vertex in  $P_j$  in graph  $G$ .

## Definition: $H$ -partition


A  $H$ -partition of a graph  $G$  is a pair  $(A, H)$ .  $A = \{A_x | x \in V(H)\}$  is a partition of  $V(G)$  and  $H$  is a graph isomorphic to the quotient  $G/A$ .



# Layering



# Layering



# Layered width


## Definition: Layered width

The layered width of a partition  $\mathcal{P}$  of a graph  $G$  is the minimum integer  $\ell$  such that there exists a path-partition (layering) of  $G$ , each element in  $\mathcal{P}$  has at most  $\ell$  vertices in each element of path-partition.

# Layered width

## Definition: Layered width

The layered width of a partition  $\mathcal{P}$  of a graph  $G$  is the minimum integer  $\ell$  such that there exists a path-partition (layering) of  $G$ , each element in  $\mathcal{P}$  has at most  $\ell$  vertices in each element of path-partition.






# Layered width

## Definition: Layered width


The layered width of a partition  $\mathcal{P}$  of a graph  $G$  is the minimum integer  $\ell$  such that there exists a path-partition (layering) of  $G$ , each element in  $\mathcal{P}$  has at most  $\ell$  vertices in each element of path-partition.



# Layered width

## Definition: Layered width

The layered width of a partition  $\mathcal{P}$  of a graph  $G$  is the minimum integer  $\ell$  such that there exists a path-partition (layering) of  $G$ , each element in  $\mathcal{P}$  has at most  $\ell$  vertices in each element of path-partition.




### Definition: Vertical Path

Let  $T$  be a tree rooted at a vertex  $r$ , a non-empty path  $(x_1, \dots, x_p)$  in  $T$  is vertical if for some  $d \geq 0$  and for all  $1 \leq i \leq p$  we have  $\text{dist}_T(x_i, r) = d + i$ .

# Layered width: Tree

## Definition: Vertical Path


Let  $T$  be a tree rooted at a vertex  $r$ , a non-empty path  $(x_1, \dots, x_p)$  in  $T$  is vertical if for some  $d \geq 0$  and for all  $1 \leq i \leq p$  we have  $\text{dist}_T(x_i, r) = d + i$ .



# Layered width: Tree

## Definition: Vertical Path


Let  $T$  be a tree rooted at a vertex  $r$ , a non-empty path  $(x_1, \dots, x_p)$  in  $T$  is vertical if for some  $d \geq 0$  and for all  $1 \leq i \leq p$  we have  $\text{dist}_T(x_i, r) = d + i$ .



# Layered width: Tree

## Definition: Vertical Path


Let  $T$  be a tree rooted at a vertex  $r$ , a non-empty path  $(x_1, \dots, x_p)$  in  $T$  is vertical if for some  $d \geq 0$  and for all  $1 \leq i \leq p$  we have  $\text{dist}_T(x_i, r) = d + i$ .



# Layered width: Tree

## Definition: Vertical Path


Let  $T$  be a tree rooted at a vertex  $r$ , a non-empty path  $(x_1, \dots, x_p)$  in  $T$  is vertical if for some  $d \geq 0$  and for all  $1 \leq i \leq p$  we have  $\text{dist}_T(x_i, r) = d + i$ .



# Layered width: Tree

## Definition: Vertical Path

Let  $T$  be a tree rooted at a vertex  $r$ , a non-empty path  $(x_1, \dots, x_p)$  in  $T$  is vertical if for some  $d \geq 0$  and for all  $1 \leq i \leq p$  we have  $\text{dist}_T(x_i, r) = d + i$ .






# Layered width: Tree

## Definition: Vertical Path


Let  $T$  be a tree rooted at a vertex  $r$ , a non-empty path  $(x_1, \dots, x_p)$  in  $T$  is vertical if for some  $d \geq 0$  and for all  $1 \leq i \leq p$  we have  $\text{dist}_T(x_i, r) = d + i$ .



# Layered width: Tree

## Definition: Vertical Path

Let  $T$  be a tree rooted at a vertex  $r$ , a non-empty path  $(x_1, \dots, x_p)$  in  $T$  is vertical if for some  $d \geq 0$  and for all  $1 \leq i \leq p$  we have  $\text{dist}_T(x_i, r) = d + i$ .



A partition of the tree where each part is a vertical path has layered width 1.

# Planar Graph Decomposition

Theorem (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

*For a graph  $G$ , if  $G$  has an  $H$ -partition of layered width  $\ell$  and  $H$  has treewidth  $k$ , then*

$$qn(G) \leq 3l(2^k - 1) + \lfloor \frac{3}{2}\ell \rfloor$$

# Planar Graph Decomposition

Theorem (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

*For a graph  $G$ , if  $G$  has an  $H$ -partition of layered width  $\ell$  and  $H$  has treewidth  $k$ , then*

$$qn(G) \leq 3l(2^k - 1) + \lfloor \frac{3}{2}\ell \rfloor$$

Theorem (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

*Every planar graph  $G$  has a connected partition  $\mathcal{P}$  with layered width 1 such that  $H = G/\mathcal{P}$  has treewidth at most 8.*

# Planar Graph Decomposition

Theorem (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

*For a graph  $G$ , if  $G$  has an  $H$ -partition of layered width  $\ell$  and  $H$  has treewidth  $k$ , then*

$$qn(G) \leq 3l(2^k - 1) + \lfloor \frac{3}{2}\ell \rfloor$$

Theorem (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

*Every planar graph  $G$  has a connected partition  $\mathcal{P}$  with layered width 1 such that  $H = G/\mathcal{P}$  has treewidth at most 8.*

Theorem (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

*Every planar graph  $G$  has queue-number at most*

$$3(2^8 - 1) + \lfloor \frac{3}{2} \rfloor = 766$$

# Planar Graph Decomposition

## Lemma (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

Let  $G^+$  be a maximal planar graph, let  $T$  be a spanning tree of  $G^+$  rooted at vertex  $r$  on outerface of  $G^+$ . For any cycle  $F$  in  $G^+$ , which can be partitioned into at most 6 pairwise disjoint vertical paths of  $T$ , with  $F = [P_1, \dots, P_k]$  and  $1 \leq k \leq 6$ . Let  $G$  be the internally triangulated subgraph of  $G^+$  which consists of all edges and vertices of  $G^+$  contained in  $F$  and the interior of  $F$ , then  $G$  has a partition  $\mathcal{P}$  into vertical paths of  $T$ , and  $P_1, \dots, P_k \in \mathcal{P}$ , and the quotient graph  $H = G/\mathcal{P}$  has a tree-decomposition  $(B, T)$  that


- $|B_x| \leq 9$  for any  $x \in V(T)$
- $\exists x \in V(T)$ , all vertices correspond to  $P_1, \dots, P_k$  in  $H$  is in  $B_x$

# Planar Graph Decomposition

## Lemma (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

Let  $G^+$  be a maximal planar graph, let  $T$  be a spanning tree of  $G^+$  rooted at vertex  $r$  on outerface of  $G^+$ . For any cycle  $F$  in  $G^+$ , which can be partitioned into at most 6 pairwise disjoint vertical paths of  $T$ , with  $F = [P_1, \dots, P_k]$  and  $1 \leq k \leq 6$ . Let  $G$  be the internally triangulated subgraph of  $G^+$  which consists of all edges and vertices of  $G^+$  contained in  $F$  and the interior of  $F$ , then  $G$  has a partition  $\mathcal{P}$  into vertical paths of  $T$ , and  $P_1, \dots, P_k \in \mathcal{P}$ , and the quotient graph  $H = G/\mathcal{P}$  has a tree-decomposition  $(B, T)$  that

- $|B_x| \leq 9$  for any  $x \in V(T)$
- $\exists x \in V(T)$ , all vertices correspond to  $P_1, \dots, P_k$  in  $H$  is in  $B_x$




# Planar Graph Decomposition

Lemma (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

Let  $G^+$  be a maximal planar graph, let  $T$  be a spanning tree of  $G^+$  rooted at vertex  $r$  on outerface of  $G^+$ . For any cycle  $F$  in  $G^+$ , which can be partitioned into at most 6 pairwise disjoint vertical paths of  $T$ , with  $F = [P_1, \dots, P_k]$  and  $1 \leq k \leq 6$ . Let  $G$  be the internally triangulated subgraph of  $G^+$  which consists of all edges and vertices of  $G^+$  contained in  $F$  and the interior of  $F$ , then  $G$  has a partition  $\mathcal{P}$  into vertical paths of  $T$ , and  $P_1, \dots, P_k \in \mathcal{P}$ , and the quotient graph  $H = G/\mathcal{P}$  has a tree-decomposition  $(B, T)$  that

- $|B_x| \leq 9$  for any  $x \in V(T)$
- $\exists x \in V(T)$ , all vertices correspond to  $P_1, \dots, P_k$  in  $H$  is in  $B_x$






# Planar Graph Decomposition

Lemma (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

Let  $G^+$  be a maximal planar graph, let  $T$  be a spanning tree of  $G^+$  rooted at vertex  $r$  on outerface of  $G^+$ . For any cycle  $F$  in  $G^+$ , which can be partitioned into at most 6 pairwise disjoint vertical paths of  $T$ , with  $F = [P_1, \dots, P_k]$  and  $1 \leq k \leq 6$ . Let  $G$  be the internally triangulated subgraph of  $G^+$  which consists of all edges and vertices of  $G^+$  contained in  $F$  and the interior of  $F$ , then  $G$  has a partition  $\mathcal{P}$  into vertical paths of  $T$ , and  $P_1, \dots, P_k \in \mathcal{P}$ , and the quotient graph  $H = G/\mathcal{P}$  has a tree-decomposition  $(B, T)$  that

- $|B_x| \leq 9$  for any  $x \in V(T)$
- $\exists x \in V(T)$ , all vertices correspond to  $P_1, \dots, P_k$  in  $H$  is in  $B_x$




# Planar Graph Decomposition

Lemma (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

Let  $G^+$  be a maximal planar graph, let  $T$  be a spanning tree of  $G^+$  rooted at vertex  $r$  on outerface of  $G^+$ . For any cycle  $F$  in  $G^+$ , which can be partitioned into at most 6 pairwise disjoint vertical paths of  $T$ , with  $F = [P_1, \dots, P_k]$  and  $1 \leq k \leq 6$ . Let  $G$  be the internally triangulated subgraph of  $G^+$  which consists of all edges and vertices of  $G^+$  contained in  $F$  and the interior of  $F$ , then  $G$  has a partition  $\mathcal{P}$  into vertical paths of  $T$ , and  $P_1, \dots, P_k \in \mathcal{P}$ , and the quotient graph  $H = G/\mathcal{P}$  has a tree-decomposition  $(B, T)$  that

- $|B_x| \leq 9$  for any  $x \in V(T)$
- $\exists x \in V(T)$ , all vertices correspond to  $P_1, \dots, P_k$  in  $H$  is in  $B_x$




# Planar Graph Decomposition

Lemma (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

Let  $G^+$  be a maximal planar graph, let  $T$  be a spanning tree of  $G^+$  rooted at vertex  $r$  on outerface of  $G^+$ . For any cycle  $F$  in  $G^+$ , which can be partitioned into at most 6 pairwise disjoint vertical paths of  $T$ , with  $F = [P_1, \dots, P_k]$  and  $1 \leq k \leq 6$ . Let  $G$  be the internally triangulated subgraph of  $G^+$  which consists of all edges and vertices of  $G^+$  contained in  $F$  and the interior of  $F$ , then  $G$  has a partition  $\mathcal{P}$  into vertical paths of  $T$ , and  $P_1, \dots, P_k \in \mathcal{P}$ , and the quotient graph  $H = G/\mathcal{P}$  has a tree-decomposition  $(B, T)$  that

- $|B_x| \leq 9$  for any  $x \in V(T)$
- $\exists x \in V(T)$ , all vertices correspond to  $P_1, \dots, P_k$  in  $H$  is in  $B_x$




# Planar Graph Decomposition

Lemma (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

Let  $G^+$  be a maximal planar graph, let  $T$  be a spanning tree of  $G^+$  rooted at vertex  $r$  on outerface of  $G^+$ . For any cycle  $F$  in  $G^+$ , which can be partitioned into at most 6 pairwise disjoint vertical paths of  $T$ , with  $F = [P_1, \dots, P_k]$  and  $1 \leq k \leq 6$ . Let  $G$  be the internally triangulated subgraph of  $G^+$  which consists of all edges and vertices of  $G^+$  contained in  $F$  and the interior of  $F$ , then  $G$  has a partition  $\mathcal{P}$  into vertical paths of  $T$ , and  $P_1, \dots, P_k \in \mathcal{P}$ , and the quotient graph  $H = G/\mathcal{P}$  has a tree-decomposition  $(B, T)$  that

- $|B_x| \leq 9$  for any  $x \in V(T)$
- $\exists x \in V(T)$ , all vertices correspond to  $P_1, \dots, P_k$  in  $H$  is in  $B_x$




# Planar Graph Decomposition

Lemma (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

Let  $G^+$  be a maximal planar graph, let  $T$  be a spanning tree of  $G^+$  rooted at vertex  $r$  on outerface of  $G^+$ . For any cycle  $F$  in  $G^+$ , which can be partitioned into at most 6 pairwise disjoint vertical paths of  $T$ , with  $F = [P_1, \dots, P_k]$  and  $1 \leq k \leq 6$ . Let  $G$  be the internally triangulated subgraph of  $G^+$  which consists of all edges and vertices of  $G^+$  contained in  $F$  and the interior of  $F$ , then  $G$  has a partition  $\mathcal{P}$  into vertical paths of  $T$ , and  $P_1, \dots, P_k \in \mathcal{P}$ , and the quotient graph  $H = G/\mathcal{P}$  has a tree-decomposition  $(B, T)$  that

- $|B_x| \leq 9$  for any  $x \in V(T)$
- $\exists x \in V(T)$ , all vertices correspond to  $P_1, \dots, P_k$  in  $H$  is in  $B_x$




# Planar Graph Decomposition

Lemma (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

Let  $G^+$  be a maximal planar graph, let  $T$  be a spanning tree of  $G^+$  rooted at vertex  $r$  on outerface of  $G^+$ . For any cycle  $F$  in  $G^+$ , which can be partitioned into at most 6 pairwise disjoint vertical paths of  $T$ , with  $F = [P_1, \dots, P_k]$  and  $1 \leq k \leq 6$ . Let  $G$  be the internally triangulated subgraph of  $G^+$  which consists of all edges and vertices of  $G^+$  contained in  $F$  and the interior of  $F$ , then  $G$  has a partition  $\mathcal{P}$  into vertical paths of  $T$ , and  $P_1, \dots, P_k \in \mathcal{P}$ , and the quotient graph  $H = G/\mathcal{P}$  has a tree-decomposition  $(B, T)$  that

- $|B_x| \leq 9$  for any  $x \in V(T)$
- $\exists x \in V(T)$ , all vertices correspond to  $P_1, \dots, P_k$  in  $H$  is in  $B_x$




# Planar Graph Decomposition

## Lemma (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

Let  $G^+$  be a maximal planar graph, let  $T$  be a spanning tree of  $G^+$  rooted at vertex  $r$  on outerface of  $G^+$ . For any cycle  $F$  in  $G^+$ , which can be partitioned into at most 6 pairwise disjoint vertical paths of  $T$ , with  $F = [P_1, \dots, P_k]$  and  $1 \leq k \leq 6$ . Let  $G$  be the internally triangulated subgraph of  $G^+$  which consists of all edges and vertices of  $G^+$  contained in  $F$  and the interior of  $F$ , then  $G$  has a partition  $\mathcal{P}$  into vertical paths of  $T$ , and  $P_1, \dots, P_k \in \mathcal{P}$ , and the quotient graph  $H = G/\mathcal{P}$  has a tree-decomposition  $(B, T)$  that

- $|B_x| \leq 9$  for any  $x \in V(T)$
- $\exists x \in V(T)$ , all vertices correspond to  $P_1, \dots, P_k$  in  $H$  is in  $B_x$




# Planar Graph Decomposition

## Lemma (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)


Let  $G^+$  be a maximal planar graph, let  $T$  be a spanning tree of  $G^+$  rooted at vertex  $r$  on outerface of  $G^+$ . For any cycle  $F$  in  $G^+$ , which can be partitioned into at most 6 pairwise disjoint vertical paths of  $T$ , with  $F = [P_1, \dots, P_k]$  and  $1 \leq k \leq 6$ . Let  $G$  be the internally triangulated subgraph of  $G^+$  which consists of all edges and vertices of  $G^+$  contained in  $F$  and the interior of  $F$ , then  $G$  has a partition  $\mathcal{P}$  into vertical paths of  $T$ , and  $P_1, \dots, P_k \in \mathcal{P}$ , and the quotient graph  $H = G/\mathcal{P}$  has a tree-decomposition  $(B, T)$  that

- $|B_x| \leq 9$  for any  $x \in V(T)$
- $\exists x \in V(T)$ , all vertices correspond to  $P_1, \dots, P_k$  in  $H$  is in  $B_x$







# Planar Graph Decomposition



# Planar Graph Decomposition



# Planar Graph Decomposition



# Planar Graph Decomposition


## Lemma (Sperner's Lemma)

Let  $G$  be an internally triangulated graph whose vertices are colored 1,2,3 with the outer-face  $F = [P_1, P_2, P_3]$  where each vertex in  $P_i$  is colored  $i$ . Then  $G$  contains an internal face whose vertices are colored 1,2,3.

# Planar Graph Decomposition

## Lemma (Sperner's Lemma)


Let  $G$  be an internally triangulated graph whose vertices are colored 1,2,3 with the outer-face  $F = [P_1, P_2, P_3]$  where each vertex in  $P_i$  is colored  $i$ . Then  $G$  contains an internal face whose vertices are colored 1,2,3.




# Planar Graph Decomposition

## Lemma (Sperner's Lemma)


Let  $G$  be an internally triangulated graph whose vertices are colored 1,2,3 with the outer-face  $F = [P_1, P_2, P_3]$  where each vertex in  $P_i$  is colored  $i$ . Then  $G$  contains an internal face whose vertices are colored 1,2,3.



# Planar Graph Decomposition




# Planar Graph Decomposition







# Planar Graph Decomposition




# Planar Graph Decomposition





# Planar Graph Decomposition: Construction





# Planar Graph Decomposition: Construction





# Planar Graph Decomposition: Construction





# Planar Graph Decomposition: Construction



# Planar Graph Decomposition: Construction








# Planar Graph Decomposition: Construction










# Planar Graph Decomposition: Construction



# Planar Graph Decomposition: Construction




# Planar Graph Decomposition: Correctness




- $|B_x| \leq 9$  for any  $x \in V(T)$
- $\exists x \in V(T)$ , all vertices correspond to  $P_1, \dots, P_k$  in  $H$  is in  $B_x$

# Planar Graph Decomposition: Correctness



H





T





# Planar Graph Decomposition: Correctness





# Planar Graph Decomposition: Correctness



# Planar Graph Decomposition: Correctness



# Planar Graph Decomposition: Correctness





# Wrap-Up

Theorem (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

*Every planar graph  $G$  has a connected partition  $\mathcal{P}$  with layered width 1 such that  $H = G/\mathcal{P}$  has treewidth at most 8.*

# Wrap-Up

Theorem (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

*Every planar graph  $G$  has a connected partition  $\mathcal{P}$  with layered width 1 such that  $H = G/\mathcal{P}$  has treewidth at most 8.*

Theorem (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

*For a graph  $G$ , if  $G$  has an  $H$ -partition of layered width  $\ell$  and  $H$  has treewidth  $k$ , then*

$$qn(G) \leq 3l(2^k - 1) + \lfloor \frac{3}{2}\ell \rfloor$$

# Wrap-Up

Theorem (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

*Every planar graph  $G$  has a connected partition  $\mathcal{P}$  with layered width 1 such that  $H = G/\mathcal{P}$  has treewidth at most 8.*

Theorem (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

*For a graph  $G$ , if  $G$  has an  $H$ -partition of layered width  $\ell$  and  $H$  has treewidth  $k$ , then*

$$qn(G) \leq 3l(2^k - 1) + \lfloor \frac{3}{2}\ell \rfloor$$

Theorem (Dujmović, Joret, Micek, Morin, Ueckerdt, Wood, 2020)

*Every planar graph  $G$  has queue-number at most*

$$3(2^8 - 1) + \lfloor \frac{3}{2} \rfloor = 766$$

# Open Problems

- Maximum queue-number of planar graphs
  - With a similar decomposition, the constant bound can be lower to 48.
  - With method in such a direction, that is the optimum result.
- Largest class of graphs with bounded queue-number
  - The decomposition method also applies to the proper minor-closed class of graphs.
  - Every proper minor-closed class of graphs has bounded queue-number.
- Relation between treewidth and queue-number
  - For graph with treewidth  $k$
  - Best known lower bound for queue-number is  $k + 1$ .
  - Best known upper bound for queue-number is  $2^k - 1$ .
- ...

# Portal

## 1 Introduction to Queue-Number

- Queue Layout
- Queue-Number

## 2 Introduction to Treewidth

- Treewidth

## 3 Introduction to Partitions

- Partitions
- Layering
- Layered Width
- Vertical Path

## 4 Planar Graph Decomposition

- The Decomposition Lemma
- Induction
- Construction
- Correctness

## 5 Summary

- Wrap-Up
- Open Problems