Combinatorics

September 27, 2022

Contents

1 Basis

The best way to think about combinations and permutations is to think about all permutations of n things:

$$n! = P(n, n) \tag{1}$$

Then we think about discounting orderings that are the same:

$$\frac{n!}{n_1! n_2! \cdots n_m!} = \binom{n}{n_1, n_2, \ldots, n_m}, \quad \text{where} \sum_{k=1}^{m} n_k = n \tag{2}$$

This is referred to as **permutations of multisets**. These represent items in the set that are considered the same.

From this, we can derive all other types of combinations and permutations. For standard **permutations**:

$$P(n, r) = \binom{n}{\underbrace{1, 1, \ldots, 1}_{r \text{ times}}, n - r} \tag{3}$$

In fact, since the sum of the n_k's must be n, there are always a number of implicit 1's to add up to n, e.g.:

$$\binom{n}{n_1, \ldots, n_m} = \binom{n}{n_1, \ldots, n_m, \underbrace{1, \ldots, 1}_{n-k \text{ times}}}, \quad \text{where} \; k = \sum_{i=1}^{m} n_k < n \tag{4}$$

The one exception is when there is only one term on the bottom, then there is an implicit $n - r$ term as well:

$$\binom{n}{r} = \binom{n}{r, n-r} \tag{5}$$

These are standard **combinations**.

2 Alternate Basis

Another way to think about combinations and permutations is to start with combinations:

$$C(n, r) = \binom{n}{r} = \frac{n!}{r!(n-r)!} \tag{6}$$

Permutations can then be thought of as reordering n things:

$$P(n, n) = n! \tag{7}$$

To permute a subgroup, first *choose* the subgroup, then permute:

$$P(n, r) = \binom{n}{r} r! = \frac{n!}{\cancel{r!}(n-r)!} \cancel{r!} = \frac{n!}{(n-r)!}, \quad r \leq n \tag{8}$$

3 Identities

$$\binom{n}{0} = P(n, 0) = 1 \tag{9}$$

$$\binom{0}{r} = P(0, r) = 0 \tag{10}$$

$$\binom{n}{1} = P(n, 1) = n \tag{11}$$

$$\binom{n}{r} = \binom{n}{n-r} \tag{12}$$

$$\binom{n}{r} = \binom{n-1}{r} + \binom{n-1}{r-1} \tag{13}$$

4 Notes

Permutations of multisets (repeats in input):

$$P_M(n, x, y, z) = \binom{n}{x, y, z} = \frac{n!}{x!y!z!} \tag{14}$$

Circular permutations:

$$P_C(n) = (n-1)! \tag{15}$$

4.1 Repeats

k consecutive items:

$$P(n - k + 1, r) \tag{16}$$

Count n minus k consecutive items, plus 1 representing the items as a group.
k items can't be together:

$$P(n - k, r) \cdot P(n - k + 1, k) \tag{17}$$

Count all $n - k$ items without restriction. Then place k items in $n - k$ spaces after of each item, and 1 space before first item. Then the items

Table 1: Combinatorial Operations

Operation	Meaning
$+n$	Add n choices to an event
$-n$	Remove n choices from an event
$\times n$	Add an event with n choices
$\div n$	Remove an event with n choices

4.2 Integer Solutions to Linear Expressions

$$x_1 + x_2 + x_3 + \cdots + x_n = r$$

$$\binom{n + r - 1}{r}$$

$$x_1 + x_2 + x_3 + \cdots + x_n \leq r$$

$$\binom{n + r}{r}$$

$$x_1 + x_2 + x_3 + \cdots + x_n < r$$

$$\binom{n + r - 1}{r - 1}$$