

Relation Between the Mass-Spring System and the Dynamic Speckle

December 28, 2019

1 Introduction

The biospeckle laser analysis has presented as a versatile tool in the analysis of biological activity.

2 System description

The signal y with samples $y(n)$, represent the adquisition of the signal x_M with samples $x_M(n)$, obtained in a mass-spring system of M elements, where each mass is separated of another by a distance of L/M , like can be seen in the Fig. 1.




Figure 1: Data acquisition system setup of the coffee seed.

Thus, in this system the mass are denoted as m_i , the springs as k_i and the displacements of each mass by x_i , for all $1 \leq i \leq M$.

The objective of this work it is to solve the next inverse problem: Known $y(n)$, and assuming M elements with $m_i = m = 1/L$; what values of k_i generate a signal x_M that minimize E , where

$$E = \frac{1}{2} \sum_n (y(n) - x_M(n))^2 \quad (1)$$

3 Mass-spring system

Assuming a mass spring system like seen in the Fig. 1 with $m_i = m$ we can to get the system of Eq. (2).

$$m\ddot{\mathbf{x}} = -\mathbf{P}\mathbf{x}, \quad (2)$$

where

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_{N-1} \\ x_N \end{pmatrix} \quad (3)$$

and

$$\mathbf{P}(\mathbf{k}) \equiv \mathbf{P} = \begin{pmatrix} k_1 + k_2 & -k_2 & 0 & \dots & 0 & 0 \\ -k_2 & k_2 + k_3 & -k_3 & \dots & 0 & 0 \\ 0 & -k_3 & k_3 + k_4 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & x_{N-1} + x_N & -x_N \\ 0 & 0 & 0 & \dots & -x_N & x_N \end{pmatrix}, \quad (4)$$

so that \mathbf{P} is a function of $\mathbf{k} = (k_1 \ k_2 \ k_3 \ \dots \ k_{M-1} \ k_M)^T$.

3.1 Exact solution

Knowing the system shown in the Eq. (2), we can solve It using the Eq. (5),

$$\mathbf{x}(t) = \mathbf{V} (\mathbf{D}_1 \cos(\mathbf{w}t) + \mathbf{D}_2 \sin(\mathbf{w}t)), \quad (5)$$

or Eq. (6)

$$\mathbf{x}(t) = \mathbf{V} (\text{diag}(\cos(\mathbf{w}t))\mathbf{d}_1 + \text{diag}(\sin(\mathbf{w}t))\mathbf{d}_2), \quad (6)$$

where, $\mathbf{V} = (\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_M)$ and $\mathbf{w} = (\sqrt{\lambda_1}, \sqrt{\lambda_2}, \dots, \sqrt{\lambda_M})^T$ are a matrix and a column vector conform using the eigenvectors \mathbf{e}_i and eigenvalues λ_i of \mathbf{P}/m . By other side, \mathbf{D}_1 and \mathbf{D}_2 are two any constant diagonal matrices conform by the elements of column vectors \mathbf{d}_1 and \mathbf{d}_2 respectively. Thus, we now that $\dot{\mathbf{x}}(t)$ and $\ddot{\mathbf{x}}(t)$ are defined by the Eqs. (7) and (8) respectively.

$$\dot{\mathbf{x}}(t) = \mathbf{V} (-\mathbf{D}_1 \mathbf{W} \sin(\mathbf{w}t) + \mathbf{D}_2 \mathbf{W} \cos(\mathbf{w}t)), \quad (7)$$

$$\ddot{\mathbf{x}}(t) = -\mathbf{V} (\mathbf{D}_1 \mathbf{W}^2 \cos(\mathbf{w}t) + \mathbf{D}_2 \mathbf{W}^2 \sin(\mathbf{w}t)), \quad (8)$$

being \mathbf{W} a diagonal matrix conform with the elements of vector \mathbf{w} ; thus, It is fulfill that $\mathbf{W}^2 = (\mathbf{V}\mathbf{D}_1)^{-1}(\mathbf{P}/m)(\mathbf{V}\mathbf{D}_1) = (\mathbf{V}\mathbf{D}_2)^{-1}(\mathbf{P}/m)(\mathbf{V}\mathbf{D}_2)$.

3.1.1 Constant values from two points

Now, to get the constant values in the column vectors \mathbf{d}_1 and \mathbf{d}_2 , we can use the Eq. (9)

$$\begin{pmatrix} \mathbf{V} \text{diag}(\cos(\mathbf{w}t_1)) & \mathbf{V} \text{diag}(\sin(\mathbf{w}t_1)) \\ \mathbf{V} \text{diag}(\cos(\mathbf{w}t_2)) & \mathbf{V} \text{diag}(\sin(\mathbf{w}t_2)) \end{pmatrix}^{-1} \begin{pmatrix} \mathbf{x}(t_1) \\ \mathbf{x}(t_2) \end{pmatrix} = \begin{pmatrix} \mathbf{d}_1 \\ \mathbf{d}_2 \end{pmatrix} \quad (9)$$

3.1.2 Constant values from position and velocity of a point

Now, to get the constant values in the column vectors \mathbf{d}_1 and \mathbf{d}_2 , we can use the Eq. (10)

$$\begin{pmatrix} \mathbf{V} \text{diag}(\cos(\mathbf{w}t_1)) & \mathbf{V} \text{diag}(\sin(\mathbf{w}t_1)) \\ -\mathbf{V}\mathbf{W} \text{diag}(\sin(\mathbf{w}t_1)) & \mathbf{V}\mathbf{W} \text{diag}(\cos(\mathbf{w}t_1)) \end{pmatrix}^{-1} \begin{pmatrix} \mathbf{x}(t_1) \\ \dot{\mathbf{x}}(t_1) \end{pmatrix} = \begin{pmatrix} \mathbf{d}_1 \\ \mathbf{d}_2 \end{pmatrix} \quad (10)$$

3.2 Finite differences: Knowing two consecutive samples

Applying finite differences we known that $\mathbf{x} \equiv \mathbf{x}(n)$ and $\ddot{\mathbf{x}} \equiv (\mathbf{x}(n+1) - 2\mathbf{x}(n) + \mathbf{x}(n-1))/\tau^2$, so that the Eq. (2) can be rewrite as

$$\mathbf{x}(n) = \left(2\mathbf{I} - \mathbf{P}\frac{\tau^2}{m}\right) \mathbf{x}(n-1) - \mathbf{x}(n-2), \quad (11)$$

now deriving $\mathbf{x}(n)$ by the vector $\mathbf{k} = (k_1 \ k_2 \ k_3 \ \dots \ k_{M-1} \ k_M)$, so that $\mathbf{J}(n) \equiv \frac{\partial \mathbf{x}(n)}{\partial \mathbf{k}}$, we get the Eq. (12)

$$\mathbf{J}(n) = -\frac{\tau^2}{m} \bigcup_i \left[\frac{\partial \mathbf{P}}{\partial k_i} \mathbf{x}(n-1) \right] + \left(2\mathbf{I} - \mathbf{P}\frac{\tau^2}{m}\right) \mathbf{J}(n-1) - \mathbf{J}(n-2), \quad (12)$$

where

$$\frac{\partial \mathbf{P}}{\partial k_i} = \begin{pmatrix} 0 & \dots & 0 & 0 & \dots & 0 & 0 \\ \vdots & \dots & \vdots & \vdots & \dots & \vdots & \vdots \\ 0 & \dots & 1 & -1 & \dots & 0 & 0 \\ 0 & \dots & -1 & 1 & \dots & 0 & 0 \\ \vdots & \dots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & \dots & 0 & 0 & \dots & 0 & 0 \\ 0 & \dots & 0 & 0 & \dots & 0 & 0 \end{pmatrix} i-th, \quad (13)$$

3.3 Finite differences: Knowing one sample and velocity

In this case, It is necessary define $\dot{\mathbf{x}} = \mathbf{v}$, so that the Eq. (2) can be rewrite as

$$\begin{pmatrix} \dot{\mathbf{x}} \\ \dot{\mathbf{v}} \end{pmatrix} = \begin{pmatrix} \mathbf{0} & \mathbf{I}_{M \times M} \\ -\mathbf{P}/m & \mathbf{0} \end{pmatrix} \begin{pmatrix} \mathbf{x} \\ \mathbf{v} \end{pmatrix}, \quad (14)$$

so that we got

$$\dot{\mathbf{u}} = A\mathbf{u}, \quad (15)$$

where $\mathbf{u} = (\mathbf{x}; \mathbf{v})$ and $A = (\mathbf{0}, \mathbf{I}_{M \times M}; -\mathbf{P}/m, \mathbf{0})$.

Applying finite differences we known that $\mathbf{u} \equiv \mathbf{u}(n)$ and $\dot{\mathbf{u}} \equiv (\mathbf{u}(n) - \mathbf{u}(n-1))/\tau_2$, so that the Eq. (15) can be rewrite as

$$\mathbf{u}(n) = (\mathbf{I} - A\tau_2)^{-1} \mathbf{u}(n-1), \quad (16)$$

Problems with finite differences: To got a good aproximation it is necessary to choose $\tau_2 \gg \tau$ (where τ is the value used in the section 3.2). Experimentaly was see that $\tau_2 \geq \tau^2$.

Now deriving $\mathbf{u}(n)$ by the vector $\mathbf{k} = (k_1 \ k_2 \ k_3 \ \dots \ k_{M-1} \ k_M)$, so that $\mathbf{Q}(n) \equiv \frac{\partial \mathbf{u}(n)}{\partial \mathbf{k}}$, we get the Eq. (17)

$$\mathbf{Q}(n) = \tau_2 (\mathbf{I} - A\tau_2)^{-1} \bigcup_i \left[\frac{\partial A}{\partial k_i} \mathbf{u}(n) \right] + (\mathbf{I} - A\tau_2)^{-1} \mathbf{Q}(n-1), \quad (17)$$

where

$$\frac{\partial A}{\partial k_i} = \begin{pmatrix} \mathbf{0} & \mathbf{0}_{M \times M} \\ -\frac{1}{m} \frac{\partial \mathbf{P}}{\partial k_i} & \mathbf{0} \end{pmatrix}, \quad (18)$$

4 Minimization problem

The minimization problem seen in the Eq. (1) can be rewrite as

$$E(\mathbf{k}) = \frac{1}{2} \sum_n (y(n) - \mathbf{b}^T \mathbf{x}(n, \mathbf{k}))^2 \quad (19)$$

where $\mathbf{b} = (0 \ 0 \ 0 \ \dots \ 0 \ 1)^T$, $y(n)$ are known values and $\mathbf{x}(n)$ that is a function of $\mathbf{k} = (k_1 \ k_2 \ k_3 \ \dots \ k_{M-1} \ k_M)^T$.

Now, knowing that a minimum of $E(\mathbf{k})$ in \mathbf{k} is found when $\frac{\partial E(\mathbf{k})}{\partial k_i} = 0$ for all integer $1 \leq i \leq M$; we calculate the Eq. (20).

$$\frac{\partial E(\mathbf{k})}{\partial k_i} = \sum_n \left(\mathbf{b}^T \frac{\partial \mathbf{x}(n, \mathbf{k})}{\partial k_i} \right)^T (\mathbf{b}^T \mathbf{x}(n, \mathbf{k}) - y(n)), \quad (20)$$

Now reordering the Eq. (20) using a vectorial differentiation by \mathbf{k} , we get the Eq. (21).

$$\frac{\partial E(\mathbf{k})}{\partial \mathbf{k}} = \sum_n (\mathbf{b}^T \mathbf{J}(n, \mathbf{k}))^T (\mathbf{b}^T \mathbf{x}(n, \mathbf{k}) - y(n)), \quad (21)$$

where $\mathbf{J}(n) = \frac{\partial \mathbf{x}(n)}{\partial \mathbf{k}}$.

4.1 Landweber iterative method

The Landweber iteration method propose that the minimization of a nonlinear function $E(\mathbf{k})$ can be found using the gradient descent method, so that

$$\mathbf{k}_j \leftarrow \mathbf{k}_{j-1} - \alpha \frac{\partial E(\mathbf{k}_{j-1})}{\partial \mathbf{k}} \quad (22)$$

where $0 < \alpha < 2/\|\frac{\partial E(\mathbf{k})}{\partial \mathbf{k}}\|^2$ and $\|\cdot\|$ is the spectral norm. Thus, following the Landweber iteration method and using the Eq. (21) in our minimization problem, It can be solved using the Eq. (23).

$$\mathbf{k}_j \leftarrow \mathbf{k}_{j-1} - \alpha \sum_n (\mathbf{b}^T \mathbf{J}(n, \mathbf{k}_{j-1}))^T (\mathbf{b}^T \mathbf{x}(n, \mathbf{k}_{j-1}) - y(n)) \quad (23)$$

4.2 Tikhonov iterative method

If we assume that the problem of to get \mathbf{k} will be solved iteratively, we can rewrite the Eq. (21) as if was evaluated by $\mathbf{x}_j(n)$ and $\mathbf{J}_{j-1}(n)$, as in the Eq. (24).

$$\sum_n \left\{ (\mathbf{b}^T \mathbf{J}_{j-1}(n))^T (\mathbf{b}^T \mathbf{x}_j(n) - y(n)) \right\} = \mathbf{0}. \quad (24)$$

Where $\mathbf{J}_{j-1}(n) = \mathbf{J}(n, \mathbf{k}_{j-1})$ and $\mathbf{x}_j(n) = \mathbf{x}(n, \mathbf{k}_{j-1})$.

Knowing by the Taylor theorem that $\mathbf{x}_j(n) \approx \mathbf{x}_{j-1}(n) + \mathbf{J}_{j-1}(n) (\mathbf{k}_j - \mathbf{k}_{j-1})$

$$\mathbf{k}_j = \mathbf{k}_{j-1} + \left(\sum_n (\mathbf{b}^T \mathbf{J}_{j-1}(n))^T (\mathbf{b}^T \mathbf{J}_{j-1}(n)) \right)^{-1} \sum_n (\mathbf{b}^T \mathbf{J}_{j-1}(n))^T (y(n) - \mathbf{b}^T \mathbf{x}_{j-1}(n)). \quad (25)$$

joint with the Eqs. (11) and (12) we got:

$$\mathbf{x}(n, \mathbf{k}) = \left(2\mathbf{I} - \mathbf{P}(\mathbf{k}) \frac{\tau^2}{m} \right) \mathbf{x}(n-1, \mathbf{k}) - \mathbf{x}(n-2, \mathbf{k}), \quad (26)$$

$$\mathbf{J}(n, \mathbf{k}) = -\frac{\tau^2}{m} \bigcup_i \left[\frac{\partial(\mathbf{P})}{\partial k_i} \mathbf{x}(n-1, \mathbf{k}) \right] + \left(2\mathbf{I} - \mathbf{P}(\mathbf{k}) \frac{\tau^2}{m} \right) \mathbf{J}(n-1, \mathbf{k}) - \mathbf{J}(n-2, \mathbf{k}), \quad (27)$$

5 Numerical results

oioio

6 Conclusion

In this work were presented

7 Acknowledgment

We wish to acknowledge the partial financial support for this study provided by the *CAPES* scholarship *PNPD* Program, *FAPEMIG* and *CNPQ*.

8 Bibliography

References
