

# Thesis Abstract

## Space-Time Discontinuous Petrov-Galerkin Finite Elements for Transient Computational Fluid Dynamics

Truman E. Ellis

**Committee:** Leszek Demkowicz (co-advisor), Robert Moser (co-advisor),  
Thomas Hughes, Clint Dawson, Jayathi Murthy

Please note that at the time of this application, I am still about a year out from completing my dissertation. As such, I am including a slightly modified summary of my thesis proposal.

## Background and Completed Work

**DPG - a robust adaptive method for CFD.** The discontinuous Petrov-Galerkin method [1] is a novel adaptive finite element framework with exceptional stability properties. Initial mesh design is a time-consuming and expensive part of CFD simulations, as a domain expert has to manually design the mesh to achieve near resolution in all parts of the domain lest the numerical method become unstable. DPG in contrast does not have a pre-asymptotic regime, allowing simulations to start on the coarsest mesh that can adequately represent the domain geometry. *A posteriori* error estimation and adaptivity can also be done very naturally as DPG comes with an error representation function that indicates residual error.

Automatic adaptivity and pre-asymptotic stability produce a powerful synthesis when combined with high performance computing. Human intervention to correct or adapt a failed massively parallel simulation can be a costly endeavor, prompting the desire for a numerical technology that will automatically adapt to changing physical dynamics while avoiding “crashing”

due to under-resolved meshes. An ideal parallel algorithm should be locally compute intensive while maintaining minimal memory transfer requirements [3]. These are all observed properties of the discontinuous Petrov-Galerkin finite element method. Locally computed *optimal test functions* ensure stability on convection-dominated flows as well as resolved viscous flow. Element contributions to the global stiffness matrix can be computed independently due to the use of discontinuous test functions. Furthermore, static condensation allows the global solve to concern only the *trace* degrees of freedom; trace variables are defined on element interfaces. This allows significant reduction of the cost of the global solve. The internal degrees of freedom can then be resolved using a fully parallel post-processing step.

**Importance of local conservation.** Local conservation was an early concern about applying DPG to fluid problems. This property is tied to whether or not constants are represented in a method’s test space. Many of the top numerical methods for CFD are automatically conservative by construction, but DPG carried no such guarantee since the test space is constructed on the fly. I ran a number of test problems and found that indeed, standard DPG is not exactly conservative but converges to near exact local conservation as it resolves the flow features under consideration. Leveraging Lagrange multipliers, I developed a new locally conservative DPG formulation [2] that maintains exact conservation even on coarse meshes. This was found to dramatically improve the coarse mesh quality on several Stokes flow test problems. We also proved stability and robustness estimates for this variant formulation.

**Extending DPG to transient problems.** Most transient CFD simulation tools are usually a combination of some spatial solver coupled with a time stepping algorithm. Often every spatial element is advanced in lock-step – requiring the time step to conform to the strictest element in the whole domain. This approach can be very ineffective for an adaptive simulation where element sizes can vary by several orders of magnitude. Technologies exist to address these inefficiencies (such as asynchronous variational integrators [4]), but bolting on another technology causes us to lose a lot of the desirable DPG properties such as automatic adaptivity and stability.

The goal of this dissertation is to extend the current technology of adaptive steady state DPG to transient problems in space-time. The emphasis

will be on transient incompressible and compressible Navier-Stokes. We have already pursued some preliminary research in this direction with a space-time DPG solver for spatially 1D flows. We’ve applied this solver to a number of test problems for the heat equation, convection-diffusion, Burgers’ equation, and compressible Navier-Stokes shock tube problems. On problems with analytical solutions, we have observed optimal rates of convergence. The application of DPG to massively parallel CFD is the ultimate goal of this research, but this dissertation will be largely developmental.

## Current Work

**Mathematical analysis of space-time DPG.** DPG is a method built on rigorous mathematical theory. While rooted in the standard theory of finite elements, DPG is novel enough that many of the old analysis tools do not directly apply. As a consequence, much of the early DPG literature has been laden with mathematical proofs of stability, convergence, and robustness. While developing the theory for a locally conservative DPG formulation we performed a stability and robustness analysis. Another robustness analysis will be necessary as we explore space-time convection-diffusion since the equation is parabolic in space-time. Past DPG robustness analysis focused on purely elliptic problems.

One significant remaining obstacle to obtaining robust solutions for compressible Navier-Stokes is guaranteeing positivity of the density and energy in the presence of unresolved shocks. Positivity is an auxiliary stability condition that is not guaranteed by traditional stabilization techniques (including DPG), so we are going to need to augment an additional technology to the method in order to guarantee physically realizable densities and energies. Unfortunately, positivity preservation is itself an incredibly difficult problem. We have several ideas we want to try in the context of DPG, but solving the issue may not lie within the scope of this dissertation.

**Scientific code development and computation.** I will support development of the DPG software framework Camellia [5] including running verification on several test problems. This will include the development and verification of spatially 2D space-time simulations. I will also develop a means to perform time stepping via space-time slabs which should reduce the overall run time and memory requirements for longer simulations. I’ve already con-

tributed several auxiliary features to Camellia including mesh readers and solution export, but I also plan on looking into adding HDF5 support along with parallel solution output.

A space-time implementation adds additional dimensionality to the problem under consideration increasing the problem size and memory requirements. This makes parallel simulation an increasingly important aspect of this work. The 16 core *Nozomi* cluster is sufficient for early experimentation, but as we ramp up to larger problems, I plan to run on allocations at TACC and on *Mira* at Argonne National Laboratory.

The major bottleneck to parallel simulations at the moment is that Camellia still relies on a serial direct solver for the global solve. Time permitting, I would like to investigate the implementation of an effective iterative solver for DPG.

**Validation problems to consider.** As we are exploring a method that is still very much in development, I will be running a number of standard test problems. I've already explored the conservation properties of DPG through a couple of simulations of Stokes flow around a cylinder and over a backward facing step.

The steady state DPG formulation failed to converge on the Carter plate problem for Reynolds numbers of  $10^7$ . One hypothesis is that this was due to transient effects on the unresolved mesh which pushed the time step requirements very low. The hope is that by using a space-time formulation with temporal adaptivity, we would be able to resolve these temporal oscillations and they would die away to the correct steady state solution. Another hypothesis is that the non-convergence was caused by Gibbs phenomenon near the unresolved leading edge driving the density negative. Our line search algorithm scales the Newton updates to prevent negative densities, but sometimes the line search factor can be driven so small as to effectively stall convergence. Therefore my first priority will be to investigate which factor is stalling convergence and attempt to solve it either via adaptive space-time refinements or some sort of positivity preserving technology yet to be decided upon. If this proves to be an effective way of handling transient shock problems, I would also like to simulate the Sedov explosion problem and the Noh implosion problem. Time permitting, I would also like to run the triple point shock interaction problem and a Rayleigh-Taylor instability problem. These will be ideal for demonstrating the local adaptivity of DPG.

That said, I believe that the most promising application for space-time DPG will be the incompressible Navier-Stokes equations. Here we don't run into any of the same issues with Gibbs phenomenon and we can use vanilla space-time DPG. The Taylor-Green vortex problem is an obvious choice since it has a exact solution that we can compare to. There are several vortex shedding problems that would be of interest including flow over a triangle, square, and cylinder.

## References

- [1] L.F. Demkowicz and J. Gopalakrishnan. *Recent Developments in Discontinuous Galerkin Finite Element Methods for Partial Differential Equations* (eds. X. Feng, O. Karakashian, Y. Xing), volume 157, chapter An Overview of the DPG Method, pages 149–180. IMA Volumes in Mathematics and its Applications, 2014.
- [2] T.E. Ellis, L.F. Demkowicz, and J.L. Chan. Locally conservative discontinuous Petrov-Galerkin finite elements for fluid problems. *Comp. Math. Appl.*, 2014.
- [3] Tz. Kolev, R. Rieben, V. Dobrev, and R. Anderson. BLAST: Paving the Way for Future Computing Architectures. <https://computation-rnd.llnl.gov/blast/>, 2014. [Online; accessed 6-March-2014].
- [4] A. Lew, J.E. Marsden, M. Ortiz, and M. West. Asynchronous variational integrators. In *Arch. Rational Mech. Anal.*, 2003.
- [5] N.V. Roberts, D. Ridzal, P.B. Bochev, and L.F. Demkowicz. A toolbox for a class of discontinuous Petrov-Galerkin methods using Trilinos. Technical Report SAND2011-6678, Sandia National Laboratories, 2011.