
Constructibility of polygons

With special emphasis on examples

Carl Friedrich Gauss, ID 036782976

Supervisor: Prof. Johann Friedrich Pfaff



A thesis presented for the degree of Master of Science in *name of programme*

School of Mathematical Sciences and possible joint school

Queen Mary University of London

Declaration of original work

This declaration is made on June 26, 2024.

Student's Declaration: I Student Name hereby declare that the work in this thesis is my original work. I have not copied from any other students' work, work of mine submitted elsewhere, or from any other sources except where due reference or acknowledgement is made explicitly in the text. Furthermore, no part of this dissertation has been written for me by another person, by generative artificial intelligence (AI), or by AI-assisted technologies.

Referenced text has been flagged by:

- 1. Using italic fonts, and
- 2. using quotation marks "...", and
- 3. explicitly mentioning the source in the text.

This work is dedicated to my dog Charles Frederick.

Acknowledgements

Abstract

Contents

1	Introduction			6	
	1.1	.1 Motivation for this work			
		1.1.1	The problem of exponential extensions	7	
		1.1.2	The approach of Junderstein	7	
2	Eulerian topological string motives			8	
	2.1	Defini	tions	8	
		2.1.1	Tate's theorem	9	
		2.1.2	Grothendieck topologies	9	
	2.2	Calcu	lation of the invariant cycles	10	
		2.2.1	Fontaine's theorem	10	
3	Conclusions		11		
A	Implementation of the BarrierOptionCVA class			12	
В	3 Shorter running title			13	

Chapter 1

Introduction

This note presents a conjecture stemming from our investigations in the generation of sigmoid tensor categories of Picard numbers of tori in Banach algebras. Nulla ac nisl. Nullam urna nulla, ullamcorper in, interdum sit amet, gravida ut, risus. Aenean ac enim. In luctus. Phasellus eu quam vitae turpis viverra pellentesque. Duis feugiat felis ut enim. Phasellus pharetra, sem id porttitor sodales, magna nunc aliquet nibh, nec blandit nisl mauris at pede. Suspendisse risus risus, lobortis eget, semper at, imperdiet sit amet, quam. Quisque scelerisque dapibus nibh. Nam enim. Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Nunc ut metus. Ut metus justo, auctor at, ultrices eu, sagittis ut, purus. Aliquam aliquam.

1.1 Motivation for this work

In the works of Petri ([Petri]) we find the following statement

Theorem 1.1.1 ([Petri], see also [BlackScholes]). The Gramm matrix for

 E_8 is:

$$\begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 2 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 & 2 \end{pmatrix}.$$

1.1.1 The problem of exponential extensions

Donec et nisl id sapien blandit mattis. Aenean dictum odio sit amet risus. Morbi purus. Nulla a est sit amet purus venenatis iaculis. Vivamus viverra purus vel magna. Donec in justo sed odio malesuada dapibus. Nunc ultrices aliquam nunc. Vivamus facilisis pellentesque velit. Nulla nunc velit, vulputate dapibus, vulputate id, mattis ac, justo. Nam mattis elit dapibus purus. Quisque enim risus, congue non, elementum ut, mattis quis, sem. Quisque elit.

1.1.2 The approach of Junderstein

Chapter 2

Eulerian topological string motives

Donec et nisl id sapien blandit mattis. Aenean dictum odio sit amet risus. Morbi purus. Nulla a est sit amet purus venenatis iaculis. Vivamus viverra purus vel magna. Donec in justo sed odio malesuada dapibus. Nunc ultrices aliquam nunc. Vivamus facilisis pellentesque velit. Nulla nunc velit, vulputate dapibus, vulputate id, mattis ac, justo. Nam mattis elit dapibus purus. Quisque enim risus, congue non, elementum ut, mattis quis, sem. Quisque elit.

2.1 Definitions

2.1.1 Tate's theorem

Preliminary considerations Donec et nisl id sapien blandit mattis. Aenean dictum odio sit amet risus. Morbi purus. Nulla a est sit amet purus venenatis iaculis. Vivamus viverra purus vel magna. Donec in justo sed odio malesuada dapibus. Nunc ultrices aliquam nunc. Vivamus facilisis pellentesque velit. Nulla nunc velit, vulputate dapibus, vulputate id, mattis ac, justo. Nam mattis elit dapibus purus. Quisque enim risus, congue non, elementum ut, mattis quis, sem. Quisque elit.

Motivic financial algebroids Donec et nisl id sapien blandit mattis. Aenean dictum odio sit amet risus. Morbi purus. Nulla a est sit amet purus venenatis iaculis. Vivamus viverra purus vel magna. Donec in justo sed odio malesuada dapibus. Nunc ultrices aliquam nunc. Vivamus facilisis pellentesque velit. Nulla nunc velit, vulputate dapibus, vulputate id, mattis ac, justo. Nam mattis elit dapibus purus. Quisque enim risus, congue non, elementum ut, mattis quis, sem. Quisque elit.

2.1.2 Grothendieck topologies

2.2 Calculation of the invariant cycles

Donec et nisl id sapien blandit mattis. Aenean dictum odio sit amet risus. Morbi purus. Nulla a est sit amet purus venenatis iaculis. Vivamus viverra purus vel magna. Donec in justo sed odio malesuada dapibus. Nunc ultrices aliquam nunc. Vivamus facilisis pellentesque velit. Nulla nunc velit, vulputate dapibus, vulputate id, mattis ac, justo. Nam mattis elit dapibus purus. Quisque enim risus, congue non, elementum ut, mattis quis, sem. Quisque elit.

2.2.1 Fontaine's theorem

Chapter 3

Conclusions

Appendix A

Implementation of the BarrierOptionCVA class

Appendix B

Additional details on the Gundermanian determinant

Bibliography

- [P99] William Petri, Analysis of infinitely generated frog complexes, Rendicoti Ranæ Analysorum, 234 (4), 34–21, 2015
- [Ross] Sheldon Ross, An Elementary Introduction to Mathematical Finance, 3rd Edition, Cambridge University Press, 2011
- [Hull] John C. Hull, *Options, Futures, and Other Derivatives*, 8th Edition, Pearson Education, 2011
- [BS] Fischer Black and Myron Scholes, *The Pricing of Options and Corporate Liabilities*, Journal of Political Economy 81 **3**, 637–654, (1973)