

Chapter 4. Part 4



February 14, 2022

Chapter 4. Part 4



February 14, 2022

Theorem

If $X = B(n, p)$ is a binomial random variable, then

$$Z = \frac{X - np}{\sqrt{np(1-p)}}$$

is approximated standard normal.

We can approximate

$$P(X \leq a) = P(X \leq a + 0.5) \approx P\left(Z \leq \frac{a + 0.5 - np}{\sqrt{np(1-p)}}\right)$$

$$P(X \geq a) = P(X \geq a - 0.5) \approx P\left(Z \geq \frac{a - 0.5 - np}{\sqrt{np(1-p)}}\right)$$

The approximation is good for $np > 5$ and $n(1-p) > 5$.

Theorem

If $X = B(n, p)$ is a binomial random variable, then

$$Z = \frac{X - np}{\sqrt{np(1-p)}}$$

is approximated standard normal.

We can approximate

$$P(X \leq a) = P(X \leq a + 0.5) \approx P\left(Z \leq \frac{a + 0.5 - np}{\sqrt{np(1-p)}}\right)$$

$$P(X \geq a) = P(X \geq a - 0.5) \approx P\left(Z \geq \frac{a - 0.5 - np}{\sqrt{np(1-p)}}\right)$$

The approximation is good for $np > 5$ and $n(1-p) > 5$.

Example

The manufacturing of semiconductor chips produces 3% defective chips. Assume the chips are independent and that a lot contains 800 chips. Approximate the probability that more than 30 chips are defective.

Example

The manufacturing of semiconductor chips produces 3% defective chips. Assume the chips are independent and that a lot contains 800 chips. Approximate the probability that more than 30 chips are defective.

Answer: Let X be the number of defective chips from the lot. Then $X = B(800, 0.03)$. We want to approximate $P(X \geq 31)$ by the normal distribution.

Example

The manufacturing of semiconductor chips produces 3% defective chips. Assume the chips are independent and that a lot contains 800 chips. Approximate the probability that more than 30 chips are defective.

Answer: Let X be the number of defective chips from the lot. Then $X = B(800, 0.03)$. We want to approximate $P(X \geq 31)$ by the normal distribution.

$$\begin{aligned} P(X \geq 31) &= P(X \geq 30.5) \approx P\left(Z \geq \frac{30.5 - 800 * 0.03}{\sqrt{800 * 0.03 * 0.97}}\right) \\ &= P(Z > 1.347) = 1 - \Phi(1.347) = 1 - 0.911 = 0.089. \end{aligned}$$

In fact,

$$P(X \geq 31) = \sum_{x=31}^{800} \binom{800}{x} 0.03^x 0.97^{800-x} = 0.09254$$

Example

The manufacturing of semiconductor chips produces 3% defective chips. Assume the chips are independent and that a lot contains 800 chips. Approximate the probability that more than 30 chips are defective.

Answer: Let X be the number of defective chips from the lot. Then $X = B(800, 0.03)$. We want to approximate $P(X \geq 31)$ by the normal distribution.

$$\begin{aligned} P(X \geq 31) &= P(X \geq 30.5) \approx P\left(Z \geq \frac{30.5 - 800 * 0.03}{\sqrt{800 * 0.03 * 0.97}}\right) \\ &= P(Z > 1.347) = 1 - \Phi(1.347) = 1 - 0.911 = 0.089. \end{aligned}$$

In fact,

$$P(X \geq 31) = \sum_{x=31}^{800} \binom{800}{x} 0.03^x 0.97^{800-x} = 0.09254$$

Example

The manufacturing of semiconductor chips produces 3% defective chips. Assume the chips are independent and that a lot contains 800 chips. Approximate the probability that between 20 and 30 chips are defective.

Example

The manufacturing of semiconductor chips produces 3% defective chips. Assume the chips are independent and that a lot contains 800 chips. Approximate the probability that between 20 and 30 chips are defective.

Answer: Let X be the number of defective chips from the lot. Then $X = B(800, 0.03)$. We want to approximate $P(21 \leq X \leq 29)$ by normal distribution.

Example

The manufacturing of semiconductor chips produces 3% defective chips. Assume the chips are independent and that a lot contains 800 chips. Approximate the probability that between 20 and 30 chips are defective.

Answer: Let X be the number of defective chips from the lot. Then $X = B(800, 0.03)$. We want to approximate $P(21 \leq X \leq 29)$ by normal distribution.

$$\begin{aligned} P(21 \leq X \leq 29) &= P(20.5 \leq X \leq 29.5) \\ &\approx P\left(\frac{20.5 - 800 * 0.03}{\sqrt{800 * 0.03 * 0.97}} \leq Z \leq \frac{29.5 - 800 * 0.03}{\sqrt{800 * 0.03 * 0.97}}\right) \\ &= P(-0.725 \leq Z \leq 1.14) = \Phi(1.14) - \Phi(-0.725) = 0.873 - 0.234 = 0.639 \end{aligned}$$

In fact,

$$P(21 \leq X \leq 29) = \sum_{x=21}^{29} \binom{800}{x} 0.03^x 0.97^{800-x} = 0.632$$

Example

The manufacturing of semiconductor chips produces 3% defective chips. Assume the chips are independent and that a lot contains 800 chips. Approximate the probability that between 20 and 30 chips are defective.

Answer: Let X be the number of defective chips from the lot. Then $X = B(800, 0.03)$. We want to approximate $P(21 \leq X \leq 29)$ by normal distribution.

$$\begin{aligned} P(21 \leq X \leq 29) &= P(20.5 \leq X \leq 29.5) \\ &\approx P\left(\frac{20.5 - 800 * 0.03}{\sqrt{800 * 0.03 * 0.97}} \leq Z \leq \frac{29.5 - 800 * 0.03}{\sqrt{800 * 0.03 * 0.97}}\right) \\ &= P(-0.725 \leq Z \leq 1.14) = \Phi(1.14) - \Phi(-0.725) = 0.873 - 0.234 = 0.639 \end{aligned}$$

In fact,

$$P(21 \leq X \leq 29) = \sum_{x=21}^{29} \binom{800}{x} 0.03^x 0.97^{800-x} = 0.632$$

Theorem

If X is a Poisson random variable with parameter λ , then

$$Z = \frac{X - \lambda}{\sqrt{\lambda}}$$

is approximately a standard normal random variable.

We can approximate:

$$P(X \leq a) = P(X \leq a + 0.5) \approx P\left(Z \leq \frac{a + 0.5 - \lambda}{\sqrt{\lambda}}\right)$$

$$P(X \geq a) = P(X \geq a - 0.5) \approx P\left(Z \geq \frac{a - 0.5 - \lambda}{\sqrt{\lambda}}\right)$$

Theorem

If X is a Poisson random variable with parameter λ , then

$$Z = \frac{X - \lambda}{\sqrt{\lambda}}$$

is approximately a standard normal random variable.

We can approximate:

$$P(X \leq a) = P(X \leq a + 0.5) \approx P\left(Z \leq \frac{a + 0.5 - \lambda}{\sqrt{\lambda}}\right)$$

$$P(X \geq a) = P(X \geq a - 0.5) \approx P\left(Z \geq \frac{a - 0.5 - \lambda}{\sqrt{\lambda}}\right)$$

The approximation is good for $\lambda > 5$.

Theorem

If X is a Poisson random variable with parameter λ , then

$$Z = \frac{X - \lambda}{\sqrt{\lambda}}$$

is approximately a standard normal random variable.

We can approximate:

$$P(X \leq a) = P(X \leq a + 0.5) \approx P\left(Z \leq \frac{a + 0.5 - \lambda}{\sqrt{\lambda}}\right)$$

$$P(X \geq a) = P(X \geq a - 0.5) \approx P\left(Z \geq \frac{a - 0.5 - \lambda}{\sqrt{\lambda}}\right)$$

The approximation is good for $\lambda > 5$.

Example

The number of customers that arrive at a fast-food business during a one-hour period is known to be Poisson distributed with a mean equal to 9.6. What is the probability that more than 10 customers will arrive in a one-hour period?

Example

The number of customers that arrive at a fast-food business during a one-hour period is known to be Poisson distributed with a mean equal to 9.6. What is the probability that more than 10 customers will arrive in a one-hour period?

Answer. Let X be the number of customers that arrive. X has a Poisson distribution with parameter $\lambda = 9.6$. We want to approximate $P(X \geq 11)$.

Example

The number of customers that arrive at a fast-food business during a one-hour period is known to be Poisson distributed with a mean equal to 9.6. What is the probability that more than 10 customers will arrive in a one-hour period?

Answer. Let X be the number of customers that arrive. X has a Poisson distribution with parameter $\lambda = 9.6$. We want to approximate $P(X \geq 11)$.

$$\begin{aligned} P(X \geq 11) &= P(X \geq 10.5) \approx P\left(Z \geq \frac{10.5 - 9.6}{\sqrt{9.6}}\right) \\ &= P(Z \geq 0.29) = 1 - \Phi(0.29) = 1 - 0.614 = 0.386 \end{aligned}$$

In fact,

$$P(X \geq 11) = \sum_{x=11}^{\infty} \frac{e^{-9.6} 9.6^x}{x!} = 0.367$$

Example

The number of customers that arrive at a fast-food business during a one-hour period is known to be Poisson distributed with a mean equal to 9.6. What is the probability that more than 10 customers will arrive in a one-hour period?

Answer. Let X be the number of customers that arrive. X has a Poisson distribution with parameter $\lambda = 9.6$. We want to approximate $P(X \geq 11)$.

$$\begin{aligned} P(X \geq 11) &= P(X \geq 10.5) \approx P\left(Z \geq \frac{10.5 - 9.6}{\sqrt{9.6}}\right) \\ &= P(Z \geq 0.29) = 1 - \Phi(0.29) = 1 - 0.614 = 0.386 \end{aligned}$$

In fact,

$$P(X \geq 11) = \sum_{x=11}^{\infty} \frac{e^{-9.6} 9.6^x}{x!} = 0.367$$