

Chapter 6. Descriptive Statistics



February 16, 2022

- 1 6.1 Numerical Summaries of Data
- 2 6.2 Stem-and-Leaf Diagrams
- 3 6.3 Frequency Distribution and Histogram

6.1 Numerical Summaries of Data

Take a **sample** of n observations (data) x_1, x_2, \dots, x_n from a **population**.

Definition

(1) **Sample mean** $\bar{x} = \frac{x_1 + x_2 + \dots + x_n}{n} = \frac{1}{n} \sum_{i=1}^n x_i$.

(2) **Sample variance**

$$s^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1} = \frac{\sum_{i=1}^n x_i^2 - n\bar{x}^2}{n - 1}.$$

6.1 Numerical Summaries of Data

Take a **sample** of n observations (data) x_1, x_2, \dots, x_n from a **population**.

Definition

(1) **Sample mean** $\bar{x} = \frac{x_1 + x_2 + \dots + x_n}{n} = \frac{1}{n} \sum_{i=1}^n x_i$.

(2) **Sample variance**

$$s^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1} = \frac{\sum_{i=1}^n x_i^2 - n\bar{x}^2}{n - 1}.$$

(3) **Sample standard deviation** $s = \sqrt{s^2}$.

6.1 Numerical Summaries of Data

Take a **sample** of n observations (data) x_1, x_2, \dots, x_n from a **population**.

Definition

(1) **Sample mean** $\bar{x} = \frac{x_1 + x_2 + \dots + x_n}{n} = \frac{1}{n} \sum_{i=1}^n x_i$.

(2) **Sample variance**

$$s^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1} = \frac{\sum_{i=1}^n x_i^2 - n\bar{x}^2}{n - 1}.$$

(3) **Sample standard deviation** $s = \sqrt{s^2}$.

(4) The **sample range** is $r = \max(x_i) - \min(x_i)$.

6.1 Numerical Summaries of Data

Take a **sample** of n observations (data) x_1, x_2, \dots, x_n from a **population**.

Definition

(1) **Sample mean** $\bar{x} = \frac{x_1 + x_2 + \dots + x_n}{n} = \frac{1}{n} \sum_{i=1}^n x_i$.

(2) **Sample variance**

$$s^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1} = \frac{\sum_{i=1}^n x_i^2 - n\bar{x}^2}{n - 1}.$$

(3) **Sample standard deviation** $s = \sqrt{s^2}$.

(4) The **sample range** is $r = \max(x_i) - \min(x_i)$.

These values are good for the estimation of mean, variance (standard deviation) of the investigating population.

6.1 Numerical Summaries of Data

Take a **sample** of n observations (data) x_1, x_2, \dots, x_n from a **population**.

Definition

(1) **Sample mean** $\bar{x} = \frac{x_1 + x_2 + \dots + x_n}{n} = \frac{1}{n} \sum_{i=1}^n x_i$.

(2) **Sample variance**

$$s^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1} = \frac{\sum_{i=1}^n x_i^2 - n\bar{x}^2}{n - 1}.$$

(3) **Sample standard deviation** $s = \sqrt{s^2}$.

(4) The **sample range** is $r = \max(x_i) - \min(x_i)$.

These values are good for the estimation of mean, variance (standard deviation) of the investigating population.

Example

The test scores of 16 students are listed below. Find sample mean and sample standard deviation.

| | | | | | | | |
|----|----|----|----|----|----|----|----|
| 32 | 41 | 47 | 50 | 56 | 58 | 58 | 61 |
| 62 | 68 | 75 | 82 | 85 | 90 | 92 | 98 |

Example

The test scores of 16 students are listed below. Find sample mean and sample standard deviation.

| | | | | | | | |
|----|----|----|----|----|----|----|----|
| 32 | 41 | 47 | 50 | 56 | 58 | 58 | 61 |
| 62 | 68 | 75 | 82 | 85 | 90 | 92 | 98 |

Answer: $\bar{x} = 65.9375, s = \sqrt{376.329} \approx 19.4$.

Example

The test scores of 16 students are listed below. Find sample mean and sample standard deviation.

| | | | | | | | |
|----|----|----|----|----|----|----|----|
| 32 | 41 | 47 | 50 | 56 | 58 | 58 | 61 |
| 62 | 68 | 75 | 82 | 85 | 90 | 92 | 98 |

Answer: $\bar{x} = 65.9375, s = \sqrt{376.329} \approx 19.4$.

$$\bar{x} = \frac{1}{16}(32 + 41 + 47 + 50 + 56 + 58 + 58 + 61 \\ + 62 + 68 + 75 + 82 + 85 + 90 + 92 + 98) = 65.9375$$

$$s^2 = \frac{1}{15} \left(\sum_{i=1}^{16} x_i^2 - 16\bar{x}^2 \right) \\ = \frac{1}{15} (32^2 + 41^2 + 47^2 + 50^2 + 56^2 + 58^2 + 58^2 + 61^2 \\ + 62^2 + 68^2 + 75^2 + 82^2 + 85^2 + 90^2 + 92^2 + 98^2 - 16 * 65.9375^2) \\ = 376.329$$

Example

The test scores of 16 students are listed below. Find sample mean and sample standard deviation.

| | | | | | | | |
|----|----|----|----|----|----|----|----|
| 32 | 41 | 47 | 50 | 56 | 58 | 58 | 61 |
| 62 | 68 | 75 | 82 | 85 | 90 | 92 | 98 |

Answer: $\bar{x} = 65.9375, s = \sqrt{376.329} \approx 19.4$.

$$\bar{x} = \frac{1}{16}(32 + 41 + 47 + 50 + 56 + 58 + 58 + 61 \\ + 62 + 68 + 75 + 82 + 85 + 90 + 92 + 98) = 65.9375$$

$$s^2 = \frac{1}{15} \left(\sum_{i=1}^{16} x_i^2 - 16\bar{x}^2 \right) \\ = \frac{1}{15} (32^2 + 41^2 + 47^2 + 50^2 + 56^2 + 58^2 + 58^2 + 61^2 \\ + 62^2 + 68^2 + 75^2 + 82^2 + 85^2 + 90^2 + 92^2 + 98^2 - 16 * 65.9375^2) \\ = 376.329$$

6.2 Stem-and-Leaf Diagrams

A **stem-and-leaf diagram** is a good way to obtain an informative visual display of a data set where each number x_i consists of at least two digits. To construct a stem-and-leaf diagram, use the following steps:

- (1) Divide each number x_i into two parts: a stem, consisting of one or more of the leading digits, and a leaf, consisting of the remaining digit.

6.2 Stem-and-Leaf Diagrams

A **stem-and-leaf diagram** is a good way to obtain an informative visual display of a data set where each number x_i consists of at least two digits. To construct a stem-and-leaf diagram, use the following steps:

- (1) Divide each number x_i into two parts: a stem, consisting of one or more of the leading digits, and a leaf, consisting of the remaining digit.
- (2) List the stem values in a vertical column.

6.2 Stem-and-Leaf Diagrams

A **stem-and-leaf diagram** is a good way to obtain an informative visual display of a data set where each number x_i consists of at least two digits. To construct a stem-and-leaf diagram, use the following steps:

- (1) Divide each number x_i into two parts: a stem, consisting of one or more of the leading digits, and a leaf, consisting of the remaining digit.
- (2) List the stem values in a vertical column.
- (3) Record the leaf for each observation beside its stem.

6.2 Stem-and-Leaf Diagrams

A **stem-and-leaf diagram** is a good way to obtain an informative visual display of a data set where each number x_i consists of at least two digits. To construct a stem-and-leaf diagram, use the following steps:

- (1) Divide each number x_i into two parts: a stem, consisting of one or more of the leading digits, and a leaf, consisting of the remaining digit.
- (2) List the stem values in a vertical column.
- (3) Record the leaf for each observation beside its stem.
- (4) Write the units for stems and leaves on the display.

6.2 Stem-and-Leaf Diagrams

A **stem-and-leaf diagram** is a good way to obtain an informative visual display of a data set where each number x_i consists of at least two digits. To construct a stem-and-leaf diagram, use the following steps:

- (1) Divide each number x_i into two parts: a stem, consisting of one or more of the leading digits, and a leaf, consisting of the remaining digit.
- (2) List the stem values in a vertical column.
- (3) Record the leaf for each observation beside its stem.
- (4) Write the units for stems and leaves on the display.

6.2 Stem-and-Leaf Diagrams

A **stem-and-leaf diagram** is a good way to obtain an informative visual display of a data set where each number x_i consists of at least two digits. To construct a stem-and-leaf diagram, use the following steps:

- (1) Divide each number x_i into two parts: a stem, consisting of one or more of the leading digits, and a leaf, consisting of the remaining digit.
- (2) List the stem values in a vertical column.
- (3) Record the leaf for each observation beside its stem.
- (4) Write the units for stems and leaves on the display.

Example

The listening scores of 17 students in a TOEIC test are listed below

| | | | | | |
|-----|-----|-----|-----|-----|-----|
| 55 | 115 | 225 | 240 | 330 | 335 |
| 385 | 400 | 405 | 405 | 495 | 495 |

The stem-and-leaf diagram:

| Stem | Leaves |
|------|--------|
| 5 | 5 |
| 11 | 5 |
| 22 | 5 |
| 24 | 0 |
| 33 | 0 5 |
| 38 | 5 |
| 40 | 0 5 5 |
| 49 | 5 5 |

Example

The listening scores of 17 students in a TOEIC test are listed below

55 115 225 240 330 335
385 400 405 405 495 495

The stem-and-leaf diagram:

| Stem | Leaves |
|------|--------|
| 5 | 5 |
| 11 | 5 |
| 22 | 5 |
| 24 | 0 |
| 33 | 0 5 |
| 38 | 5 |
| 40 | 0 5 5 |
| 49 | 5 5 |

Definition

- (1) The **sample median** is a measure of central tendency that divides the data into two equal parts, half below the median and half above. If the number of observations is even, the median is halfway between the two central values.
- (2) The **sample mode** is the most frequently occurring data value.

Definition

- (1) The **sample median** is a measure of central tendency that divides the data into two equal parts, half below the median and half above. If the number of observations is even, the median is halfway between the two central values.
- (2) The **sample mode** is the most frequently occurring data value.
- (3) An ordered set of data is divided into four equal parts, the division points are called **quartiles**:

Definition

- (1) The **sample median** is a measure of central tendency that divides the data into two equal parts, half below the median and half above. If the number of observations is even, the median is halfway between the two central values.
- (2) The **sample mode** is the most frequently occurring data value.
- (3) An ordered set of data is divided into four equal parts, the division points are called **quartiles**:
 - The **first quartile**, q_1 or Q_1 : is a value that has approximately 25% of the observations below.

Definition

- (1) The **sample median** is a measure of central tendency that divides the data into two equal parts, half below the median and half above. If the number of observations is even, the median is halfway between the two central values.
- (2) The **sample mode** is the most frequently occurring data value.
- (3) An ordered set of data is divided into four equal parts, the division points are called **quartiles**:
 - The **first quartile**, q_1 or Q_1 : is a value that has approximately 25% of the observations below.
 - The **sample median** or **second quartile**, q_2 or Q_2 , has approximately 50% of the observations below its value.

Definition

- (1) The **sample median** is a measure of central tendency that divides the data into two equal parts, half below the median and half above. If the number of observations is even, the median is halfway between the two central values.
- (2) The **sample mode** is the most frequently occurring data value.
- (3) An ordered set of data is divided into four equal parts, the division points are called **quartiles**:
 - The **first quartile**, q_1 or Q_1 : is a value that has approximately 25% of the observations below.
 - The **sample median** or **second quartile**, q_2 or Q_2 , has approximately 50% of the observations below its value.
 - The **third quartile**, q_3 or Q_3 , has approximately 75% of the observations below its value.

Definition

- (1) The **sample median** is a measure of central tendency that divides the data into two equal parts, half below the median and half above. If the number of observations is even, the median is halfway between the two central values.
- (2) The **sample mode** is the most frequently occurring data value.
- (3) An ordered set of data is divided into four equal parts, the division points are called **quartiles**:
 - The **first quartile**, q_1 or Q_1 : is a value that has approximately 25% of the observations below.
 - The **sample median** or **second quartile**, q_2 or Q_2 , has approximately 50% of the observations below its value.
 - The **third quartile**, q_3 or Q_3 , has approximately 75% of the observations below its value.
- (4) The **interquartile range**, $IQR = q_3 - q_1$.

Definition

- (1) The **sample median** is a measure of central tendency that divides the data into two equal parts, half below the median and half above. If the number of observations is even, the median is halfway between the two central values.
- (2) The **sample mode** is the most frequently occurring data value.
- (3) An ordered set of data is divided into four equal parts, the division points are called **quartiles**:
 - The **first quartile**, q_1 or Q_1 : is a value that has approximately 25% of the observations below.
 - The **sample median** or **second quartile**, q_2 or Q_2 , has approximately 50% of the observations below its value.
 - The **third quartile**, q_3 or Q_3 , has approximately 75% of the observations below its value.
- (4) The **interquartile range**, $IQR = q_3 - q_1$.

Example

Use the given sample data to find the sample quartiles, the sample mode and the *IQR*.

55, 52, 52, 52, 49, 74, 67, 55.

Example

Use the given sample data to find the sample quartiles, the sample mode and the *IQR*.

55, 52, 52, 52, 49, 74, 67, 55.

Answer: We rearrange the increasing data:

49, 52, 52, 52, 55, 55, 67, 74.

Example

Use the given sample data to find the sample quartiles, the sample mode and the *IQR*.

55, 52, 52, 52, 49, 74, 67, 55.

Answer: We rearrange the increasing data:

49, 52, 52, 52, 55, 55, 67, 74.

We the size of sample $n = 8$.

Example

Use the given sample data to find the sample quartiles, the sample mode and the *IQR*.

55, 52, 52, 52, 49, 74, 67, 55.

Answer: We rearrange the increasing data:

49, 52, 52, 52, 55, 55, 67, 74.

We the size of sample $n = 8$.

(1) $\frac{n+1}{2} = 4.5$, hence the sample median $Q_2 = \frac{x_4 + x_5}{2} = \frac{52 + 55}{2} = 53.5$.

Example

Use the given sample data to find the sample quartiles, the sample mode and the *IQR*.

55, 52, 52, 52, 49, 74, 67, 55.

Answer: We rearrange the increasing data:

49, 52, 52, 52, 55, 55, 67, 74.

We the size of sample $n = 8$.

(1) $\frac{n+1}{2} = 4.5$, hence the sample median $Q_2 = \frac{x_4 + x_5}{2} = \frac{52 + 55}{2} = 53.5$.

(2) $\frac{n+1}{4} = 2.25$, hence the first quartile $Q_1 = \frac{x_2 + x_3}{2} = \frac{52 + 52}{2} = 52$.

Example

Use the given sample data to find the sample quartiles, the sample mode and the *IQR*.

55, 52, 52, 52, 49, 74, 67, 55.

Answer: We rearrange the increasing data:

49, 52, 52, 52, 55, 55, 67, 74.

We the size of sample $n = 8$.

(1) $\frac{n+1}{2} = 4.5$, hence the sample median $Q_2 = \frac{x_4 + x_5}{2} = \frac{52 + 55}{2} = 53.5$.

(2) $\frac{n+1}{4} = 2.25$, hence the first quartile $Q_1 = \frac{x_2 + x_3}{2} = \frac{52 + 52}{2} = 52$.

(3) $\frac{3(n+1)}{4} = 6.75$, hence the third quartile $Q_3 = \frac{x_6 + x_7}{2} = \frac{55 + 67}{2} = 61$.

Example

Use the given sample data to find the sample quartiles, the sample mode and the *IQR*.

55, 52, 52, 52, 49, 74, 67, 55.

Answer: We rearrange the increasing data:

49, 52, 52, 52, 55, 55, 67, 74.

We the size of sample $n = 8$.

(1) $\frac{n+1}{2} = 4.5$, hence the sample median $Q_2 = \frac{x_4 + x_5}{2} = \frac{52 + 55}{2} = 53.5$.

(2) $\frac{n+1}{4} = 2.25$, hence the first quartile $Q_1 = \frac{x_2 + x_3}{2} = \frac{52 + 52}{2} = 52$.

(3) $\frac{3(n+1)}{4} = 6.75$, hence the third quartile $Q_3 = \frac{x_6 + x_7}{2} = \frac{55 + 67}{2} = 61$.

(4) The sample modes: 52.

Example

Use the given sample data to find the sample quartiles, the sample mode and the *IQR*.

55, 52, 52, 52, 49, 74, 67, 55.

Answer: We rearrange the increasing data:

49, 52, 52, 52, 55, 55, 67, 74.

We the size of sample $n = 8$.

(1) $\frac{n+1}{2} = 4.5$, hence the sample median $Q_2 = \frac{x_4 + x_5}{2} = \frac{52 + 55}{2} = 53.5$.

(2) $\frac{n+1}{4} = 2.25$, hence the first quartile $Q_1 = \frac{x_2 + x_3}{2} = \frac{52 + 52}{2} = 52$.

(3) $\frac{3(n+1)}{4} = 6.75$, hence the third quartile $Q_3 = \frac{x_6 + x_7}{2} = \frac{55 + 67}{2} = 61$.

(4) The sample modes: 52.

(5) $IQR = Q_3 - Q_1 = 61 - 52 = 9$.

Example

Use the given sample data to find the sample quartiles, the sample mode and the *IQR*.

55, 52, 52, 52, 49, 74, 67, 55.

Answer: We rearrange the increasing data:

49, 52, 52, 52, 55, 55, 67, 74.

We the size of sample $n = 8$.

(1) $\frac{n+1}{2} = 4.5$, hence the sample median $Q_2 = \frac{x_4 + x_5}{2} = \frac{52 + 55}{2} = 53.5$.

(2) $\frac{n+1}{4} = 2.25$, hence the first quartile $Q_1 = \frac{x_2 + x_3}{2} = \frac{52 + 52}{2} = 52$.

(3) $\frac{3(n+1)}{4} = 6.75$, hence the third quartile $Q_3 = \frac{x_6 + x_7}{2} = \frac{55 + 67}{2} = 61$.

(4) The sample modes: 52.

(5) $IQR = Q_3 - Q_1 = 61 - 52 = 9$.

Example

Use the given sample data to find the sample quartiles, the sample mode and the *IQR*.

55, 52, 52, 52, 49, 74, 67, 55.

Answer: We rearrange the increasing data:

49, 52, 52, 52, 55, 55, 67, 74.

We the size of sample $n = 8$.

(1) $\frac{n+1}{2} = 4.5$, hence the sample median $Q_2 = \frac{x_4 + x_5}{2} = \frac{52 + 55}{2} = 53.5$.

(2) $\frac{n+1}{4} = 2.25$, hence the first quartile $Q_1 = \frac{x_2 + x_3}{2} = \frac{52 + 52}{2} = 52$.

(3) $\frac{3(n+1)}{4} = 6.75$, hence the third quartile $Q_3 = \frac{x_6 + x_7}{2} = \frac{55 + 67}{2} = 61$.

(4) The sample modes: 52.

(5) $IQR = Q_3 - Q_1 = 61 - 52 = 9$.

6.3 Frequency Distribution and Histogram

Construction of **frequency distribution**: divide the range of the data into intervals (called class intervals, cells, or bins). The bins should be of equal width.

Example

We have frequency distribution of MAS291 final scores of the class IS1402.

| scores | frequency | cumulative frequency distribution | relative frequency | cumulative relative frequency |
|--------|-----------|--------------------------------------|-----------------------|----------------------------------|
| 0-2 | 2 | 2 | $2/30$ | $2/30$ |
| 2-4 | 5 | 7 | $5/30$ | $7/30$ |
| 4-6 | 8 | 15 | $8/30$ | $15/30=0.5$ |
| 6-8 | 11 | 26 | $11/30$ | $26/30$ |
| 8-10 | 4 | 30 | $4/30$ | 1 |

6.3 Frequency Distribution and Histogram

Construction of **frequency distribution**: divide the range of the data into intervals (called class intervals, cells, or bins). The bins should be of equal width.

Example

We have frequency distribution of MAS291 final scores of the class IS1402.

| scores | frequency | cumulative frequency distribution | relative frequency | cumulative relative frequency |
|--------|-----------|--------------------------------------|-----------------------|----------------------------------|
| 0-2 | 2 | 2 | $2/30$ | $2/30$ |
| 2-4 | 5 | 7 | $5/30$ | $7/30$ |
| 4-6 | 8 | 15 | $8/30$ | $15/30=0.5$ |
| 6-8 | 11 | 26 | $11/30$ | $26/30$ |
| 8-10 | 4 | 30 | $4/30$ | 1 |

The **histogram** is a visual display of the frequency distribution.

(1) Label the bin (class interval) boundaries on a horizontal scale.

The **histogram** is a visual display of the frequency distribution.

- (1) Label the bin (class interval) boundaries on a horizontal scale.
- (2) Mark and label the vertical scale with the frequencies or the relative frequencies.

The **histogram** is a visual display of the frequency distribution.

- (1) Label the bin (class interval) boundaries on a horizontal scale.
- (2) Mark and label the vertical scale with the frequencies or the relative frequencies.
- (3) Above each bin, draw a rectangle where height is equal to the frequency (or relative frequency) corresponding to that bin.

The **histogram** is a visual display of the frequency distribution.


- (1) Label the bin (class interval) boundaries on a horizontal scale.
- (2) Mark and label the vertical scale with the frequencies or the relative frequencies.
- (3) Above each bin, draw a rectangle where height is equal to the frequency (or relative frequency) corresponding to that bin.

The **histogram** is a visual display of the frequency distribution.

- (1) Label the bin (class interval) boundaries on a horizontal scale.
- (2) Mark and label the vertical scale with the frequencies or the relative frequencies.
- (3) Above each bin, draw a rectangle where height is equal to the frequency (or relative frequency) corresponding to that bin.



Example

| fruits | numbers |
|--------|---------|
| orange | 12 |
| apple | 34 |
| pear | 56 |
| kiwi | 20 |





Example

| fruits | numbers |
|--------|---------|
| orange | 12 |
| apple | 34 |
| pear | 56 |
| kiwi | 20 |




Example

| fruits | numbers |
|--------|---------|
| orange | 12 |
| apple | 34 |
| pear | 56 |
| kiwi | 20 |




| fruits | numbers |
|--------|---------|
| orange | 12 |
| apple | 34 |
| pear | 56 |
| kiwi | 20 |

Piechart



| fruits | numbers |
|--------|---------|
| orange | 12 |
| apple | 34 |
| pear | 56 |
| kiwi | 20 |

Piechart




Example

Given marks of 24 students in a mathematics examination

| | | | | | | | | | | | |
|-----|-----|---|---|-----|---|-----|-----|-----|-----|-----|----|
| 2 | 2.5 | 3 | 3 | 3.5 | 4 | 4 | 4 | 4.5 | 4.5 | 6 | 6 |
| 6.5 | 7 | 7 | 7 | 7.5 | 8 | 8.5 | 8.5 | 9 | 9 | 9.5 | 10 |

Draw a histogram with 5 bins and a histogram with 8 bins.





Example

Given marks of 24 students in a mathematics examination

| | | | | | | | | | | | |
|-----|-----|---|---|-----|---|-----|-----|-----|-----|-----|----|
| 2 | 2.5 | 3 | 3 | 3.5 | 4 | 4 | 4 | 4.5 | 4.5 | 6 | 6 |
| 6.5 | 7 | 7 | 7 | 7.5 | 8 | 8.5 | 8.5 | 9 | 9 | 9.5 | 10 |

Draw a histogram with 5 bins and a histogram with 8 bins.




Example

Given marks of 24 students in a mathematics examination

| | | | | | | | | | | | |
|-----|-----|---|---|-----|---|-----|-----|-----|-----|-----|----|
| 2 | 2.5 | 3 | 3 | 3.5 | 4 | 4 | 4 | 4.5 | 4.5 | 6 | 6 |
| 6.5 | 7 | 7 | 7 | 7.5 | 8 | 8.5 | 8.5 | 9 | 9 | 9.5 | 10 |

Draw histograms of cumulative frequency and cumulative relative frequency with 8 bins.





Example

Given marks of 24 students in a mathematics examination

| | | | | | | | | | | | |
|-----|-----|---|---|-----|---|-----|-----|-----|-----|-----|----|
| 2 | 2.5 | 3 | 3 | 3.5 | 4 | 4 | 4 | 4.5 | 4.5 | 6 | 6 |
| 6.5 | 7 | 7 | 7 | 7.5 | 8 | 8.5 | 8.5 | 9 | 9 | 9.5 | 10 |

Draw histograms of cumulative frequency and cumulative relative frequency with 8 bins.



Example

Given marks of 24 students in a mathematics examination

| | | | | | | | | | | | |
|-----|-----|---|---|-----|---|-----|-----|-----|-----|-----|----|
| 2 | 2.5 | 3 | 3 | 3.5 | 4 | 4 | 4 | 4.5 | 4.5 | 6 | 6 |
| 6.5 | 7 | 7 | 7 | 7.5 | 8 | 8.5 | 8.5 | 9 | 9 | 9.5 | 10 |

Draw histograms of cumulative frequency and cumulative relative frequency with 8 bins.

