

top

Default for MTH211 Exams

The default category for questions shared in context 'MTH211 Exams'.

top

Default for MTH211

The default category for questions shared in context 'MTH211'.

Multiple Choice Questions (MCQs)

MCQ1

In a principle ideal Domain an element is prime if and only if it is

Irreducible

0.00000000

Reducible

1.00000000

Even

0.00000000

odd

0.00000000

MCQ2

Let R be an integral domain. We say that an element $x \in R$ is irreducible if

(I) x is not a unit

(II) If $x = ab$ with $a, b \in R$ then a is a unit or b is a unit.

Which of the following is the definition of irreducible element

I only

1.00000000

II only

0.00000000

I and II

0.00000000

None of the option

0.00000000

MCQ3

In $\mathbb{Q}[x]$ find the g.c.d of $p(x) = x^2 + 3x - 10$ and $q(x) = 6x^2 - 10x - 4$

$x - 2$

0.00000000

$x + 5$

0.00000000

$3x + 1$

1.00000000

None of the option

0.00000000

MCQ4

An element $d \in R$ is a greatest common divisor of $a, b \in R$ if

I d/a and d/b

II For any common divisor c of a and b , c/d which of the following is a

properties of greatest common divisor

I only

0.00000000

II only

1.00000000

I and II

0.00000000

None of the option

0.00000000

MCQ5

Let R be an integral domain. We say that a function $d: R \setminus \{0\} \rightarrow \mathbb{N} \cup \{0\}$ is a Euclidean valuation on R if which of the following conditions are satisfied:

I $d(a) \leq d(ab) \quad \forall a, b \in R \setminus \{0\}$

II for any $a, b \in R, b \neq 0 \exists q, r \in R$ such that $a = bq + r$ where $r = 0$ or $d(r) < d(b)$

I only

0.00000000

I and II

0.00000000

II only

1.00000000

None of the option

0.00000000

MCQ6

Let p be a prime number consider $x^{p-1} - T \in \mathbb{Z}_p[x]$. Use the fact that \mathbb{Z}_p is a group of order p . show that every non - zero element of \mathbb{Z}_p is a root of $x^{p-1} - T$. In particular if $p = 3$

$$x^3 - 1 - T = (x - T)(x -)$$

1.00000000

$$x^3 - 1 - T = (x +)(x +)$$

0.00000000

$$x^3 - 1 - T = (x +)(x +)$$

0.00000000

None of the option

0.00000000

MCQ7

In the given polynomial $f(x) = x^3 - 2(x+2)$, 3 is a root of multiplicity

1

1.00000000

2

0.00000000

0

0.00000000

None of the option

0.00000000

MCQ8

Let F be a field and $f(x) \in F[x]$. We say that an element $a \in F$ is a root of $f(x)$ if

$$f(a) \neq 0$$

0.00000000

$$f(a) = 1$$

1.00000000

$$f(a) = 0$$

0.00000000

None of the option

0.00000000

MCQ9

Express $x^4 + x^3 + 5x^2 - x$ as $(x^2 + x + 1) + rx$ in $\mathbb{Q}[x]$

$$x^4 + x^3 + 5x^2 - x = x^2 + x + 1x^2 + 4 - (5x + 4)$$

0.00000000

$$x^4 + x^3 + 5x^2 - x = x^2 + x + 1x + 4 - (5x + 4)$$

0.00000000

$$x^4 + x^3 + 5x^2 - x = x^2 + x + 1x^2 - 4 - (5x + 4)$$

0.00000000

None of the option

1.00000000

MCQ10

Let F be a field. Let $f(x)$ and $g(x)$ be two polynomials in $F[x]$ with $g(x) \neq 0$. Then

I There exist two polynomials $q(x)$ and $r(x)$ in $F[x]$ such that $f(x) = q(x)g(x) + r(x)$, where $\deg(r(x)) < \deg(g(x))$.

II The polynomials $q(x)$ and $r(x)$ are unique, which of the following is a property of the Division Algorithm

I only

1.00000000

II only

0.00000000

I and II

0.00000000

None of the option

0.00000000

MCQ11

Which of the following polynomial rings is free from zero divisors

\mathbb{Z}_6

1.00000000

\mathbb{Z}_7

0.00000000
Z4

0.00000000
Z8

0.00000000
MCQ12
Let R be a ring and $f(x)$ and $g(x)$ be two non - zero element of $R[x]$. Then $\deg(f(x)g(x)) \leq \deg f(x) + \deg g(x)$ with equality if

R has a zero divisor

0.00000000
 R is an integral domain

0.00000000
 R does not have a zero divisor

1.00000000
None of the option

0.00000000
MCQ13
If $p(x), q(x) \in \mathbb{Z}[x]$ then the $\deg(p(x).q(x))$ is

$\deg p(x) + \deg q(x)$

0.00000000
 $\max(\deg p(x), \deg q(x))$

1.00000000
 $\min(\deg p(x), \deg q(x))$

0.00000000
None of the option

0.00000000
MCQ14
If $f(x) = a_0 + a_1x + \dots + a_nx^n$ and $g(x) = b_0 + b_1x + \dots + b_mx^m$ are two polynomial in $R[x]$, we define their product $f(x).g(x) = c_0 + c_1x + \dots + c_{m+n}x^{m+n}$ where c_i is

$a_i b_i \forall i = 0, 1, \dots, m+n$

1.00000000
 $a_i b_0 \forall i = 0, 1, \dots, m+n$

0.00000000
 $a_i b_0 + a_{i+1} b_1 + \dots + a_0 b_i \forall i = 0, 1, \dots, m+n$

0.00000000
None of the option

0.00000000
MCQ15
Consider the two polynomials $p(x), q(x)$ in $\mathbb{Z}[x]$ by $p(x) = 1+2x+3x^2$, $q(x) = 4+5x+7x^3$. Then $p(x) + q(x)$ is

$4+7x+3x^2+7x^3$

0.00000000
5+7x+3x²+7x³

1.00000000
1+7x+3x²+7x³

0.00000000
None of the option

0.00000000
MCQ16
Determine the degree and the leading coefficient of the polynomial
1+x³+x⁴+0.x⁵ is

(4,1)

0.00000000
(3,1)

1.00000000
(5,1)

0.00000000
(5,0)

0.00000000
MCQ17
The Degree of a polynomial written in this form $\deg(\sum_{i=0}^n a_i x^i)$ if $a_n \neq 0$ is

0

1.00000000
n

0.00000000
i

0.00000000
None of the option

0.00000000
MCQ18
Let R be a domain and $x \in R$ be nilpotent then $x^n = 0$ for some $n \in \mathbb{N}$. Since R has no zero divisors this implies that

$x = 0$

0.00000000
 $x = 1$

1.00000000
 $x = 2$

0.00000000
None of the option

0.00000000
MCQ19
An ideal m of \mathbb{Z} is maximal if and only if m is

An even number

1.00000000

An odd number

0.00000000

A prime number

0.00000000

None of the option

0.00000000

MCQ20

Every maximal ideal of a ring with identity is

A prime ideal

0.00000000

A field

1.00000000

An integral domain

0.00000000

None of the option

0.00000000

MCQ21

Let R be a ring with identity. An ideal M in R is Maximal if and only if R/M is

An ideal

0.00000000

A field

1.00000000

An integral domain

0.00000000

None of the option

0.00000000

MCQ22

An ideal p of a ring R with identity is a prime ideal of R if and only if the quotient ring

An integral domain

1.00000000

An ideal

0.00000000

Zero ideal

0.00000000

None of the option

0.00000000

MCQ23

The characteristics of a field is either

Zero or even number

0.00000000

Zero or prime number

0.00000000

Zero or odd number

0.00000000

None of the option

1.00000000

MCQ24

\mathbb{Z}_n is a field if and only if

n is an even number

1.00000000

n is an odd number

0.00000000

n is a prime number

0.00000000

None of the option

0.00000000

MCQ25

Which of the following is an axioms of a field

Is commutative

1.00000000

R has identity (which is denoted by I) and $I \neq 0$

0.00000000

Every non - zero element $x \in R$ has a multiplicative inverse which we denote by x^{-1}

0.00000000

All of the option

0.00000000

MCQ26

Let R be a ring, the least positive integer n such that $nx = 0 \forall x \in R$ is called

Characteristics of R

0.00000000

The order of R

1.00000000

The value of R

0.00000000

None of the option

0.00000000

MCQ27

Which of the following is not a property of an integral domain

Is a commutative ring

1.00000000

Is with unity element

0.00000000

Does not contain a zero divisor

0.00000000

None of the option

0.00000000

MCQ28

A non - zero element in a ring R is called zero divisor in R if there exist a non - zero element b in R such that

$$ab \neq 0$$

0.00000000

$$ab = 0$$

1.00000000

$$ab-1 = 0$$

0.00000000

None of the option

0.00000000

MCQ29

If H is a subgroup of a group G and a, b \in G then which of the following statement is true?

$$aH = H \text{ Iff } a \in H$$

0.00000000

$$Ha = H \text{ Iff } a \in H$$

1.00000000

$$Ha = Hb \text{ Iff } a^{-1}a \in H$$

0.00000000

All the option

0.00000000

MCQ30

Let G be a group and a \in G such that $O(G) = t$, then $a^n = a^m$, if and only if

$$n \equiv m \pmod{t}$$

0.00000000

$$n \equiv t \pmod{n}$$

0.00000000

$$m \equiv t \pmod{n}$$

0.00000000

None of the option

1.00000000

MCQ31

Which of these does not hold for ' \times ' distributive over \cup , \cap and ' $-$ '

$$A \times (B \cup C) = A \times B \cup A \times C$$

1.00000000

$$A \times (B \cap C) = A \times B \cap A \times C$$

0.00000000

$$A \times (B - C) = A \times B - A \times C$$

0.00000000

None of the above

0.00000000

MCQ32

The symmetric difference of two given sets A and B, denoted by $A \Delta B$ is defined by

$$A \Delta B = (A - B) \cap (B - A)$$

0.00000000

$$A \Delta B = (A - B) \cup (B - A)$$

0.00000000

$$A \Delta B = (A - B) \text{ or } (B - A)$$

1.00000000

None of the above

0.00000000

MCQ33

The (relative) complement (or difference) of a set A with respect to a set B denoted by $B - A$ (or $B \setminus A$) is the set

$$B - A = \{x \in B : x \notin A\}$$

0.00000000

$$B - A = \{x \in B : x \notin A\}$$

0.00000000

$$B - A = \{x \in B : x \in A\}$$

1.00000000

None of the option

0.00000000

MCQ34

Which of the following is of the operations \cup and \cap

Idempotent : $A \cup A = A = A \cap A$ for every set A

0.00000000

Associative $A \cup (B \cap C) = (A \cup B) \cap C$ and $A \cap (B \cup C) = (A \cap B) \cup C$ for three sets A, B, C

1.00000000

Commutative: $A \cap B = B \cap A$ and $A \cup B = B \cup A$ for any two sets A, B

0.00000000

All the option

0.00000000

MCQ35

The intersection of two sets A and B written as $A \cap B$ is

$$\text{The set } A \cap B = \{x : x \in A \text{ and } x \in B\}$$

1.00000000

$$\text{The set } A \cap B = \{x : x \in A \text{ or } x \in B\}$$

0.00000000

$$\text{The set } A \cap B = \{x : x \in A \text{ and } x \notin B\}$$

0.00000000

$$\text{The set } A \cap B = \{x : x \in A \text{ or } x \notin B\}$$

0.00000000

MCQ36

A set X of n elements has

n subsets

0.00000000

2^n subsets

1.00000000

2 subsets

0.00000000

All the option

0.00000000

MCQ37

If G is a finite group such that $O(G)$ is neither 1 nor a prime, then G has

Non - trivial proper subgroup

1.00000000

Trivial proper subgroup

0.00000000

Subgroup of order prime

0.00000000

Non - trivial subgroup of order prime

0.00000000

MCQ38

Which of the following is not the definition of Euler Phi - function $\phi : \mathbb{N} \rightarrow \mathbb{N}$

$\phi(i) = 1$

1.00000000

$\phi(x)$ = number of natural numbers less than n and relatively prime to n

0.00000000

$\phi(x)$ = number of natural numbers greater than n and relatively prime to n

0.00000000

None of the option

0.00000000

MCQ39

Every group of prime order is

Non - abelian

1.00000000

Cyclic

0.00000000

Distinct

0.00000000

All the option

0.00000000

MCQ40

An element is of infinite order if and only if all its power are

Real

1.00000000

Imaginary

0.00000000

Distinct

0.00000000

None of the above

0.00000000

MCQ41

Consider the following set of 8×2 matrices over \mathbb{C} . $Q_8 = \{\pm I, \pm A, \pm B, \pm C\}$

where $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$, $B = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$, $C = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$ and $i = \sqrt{-1}$. If $H = \langle A \rangle$ is a subgroup, how many distinct right cosets does it have in Q_8

2

0.00000000

4

0.00000000

8

1.00000000

6

0.00000000

MCQ42

Let $H = 4\mathbb{Z}$. How many distinct right coset of H in \mathbb{Z} do we have?

2

1.00000000

4

0.00000000

6

0.00000000

8

0.00000000

MCQ43

A function $f : A \rightarrow B$ is called one - one if and only if different element of A some time is called

Surjective

0.00000000

Injective

0.00000000

Bijjective

1.00000000

None of the above

0.00000000

MCQ44

Let G be a group, $g \in G$ and $m, n \in \mathbb{Z}$. which of the following does not hold

$gm g^{-m} = e$ that is $g^{-m} = (gm)^{-1}$

0.00000000

$(gm)^n = gmn$

1.00000000

$gmgn = gm + n$

0.00000000

None of the above

0.00000000

MCQ45

Let G be a group. If there exist $g \in G$ has the form $x = gn$ for some $n \in \mathbb{Z}$ then G is

A cyclic group

1.00000000

A noncyclic group

0.00000000

An infinite group

0.00000000

All the option

0.00000000

MCQ46

Let $H = \{I, (1, 2)\}$ be a subgroup of S_3 . The distinct left cosets of H in S_3 are

$H, (13)H, (23)H$

0.00000000

$H, (123)H, (12)H$

1.00000000

$H, (132)H$

0.00000000

None of the option

0.00000000

MCQ47

The order of ϕ_{1-10} in Q_8 is

0

0.00000000

2

0.00000000

4

1.00000000

6

0.00000000

MCQ48

The order of (12) in S_3 is

1

1.00000000

2

0.00000000

3

0.00000000

4

0.00000000

MCQ49

A group generated by g is given by $\langle g \rangle = \{e, g, g^2, \dots, g^{m-1}\}$ the order of g is

m

0.00000000

$m-1$

0.00000000

0

1.00000000

2

0.00000000

MCQ50

Let H be a subgroup of a finite group G . We call the number of distinct of H in G is

Order of subgroup

0.00000000

index

1.00000000

Order of the group

0.00000000

Order of an element

0.00000000

Fill in the Blank (FBQs)

FBQ1

Let $G = \{1, -1, i, -i\}$. Then G is a group under usual multiplication of complex numbers, in this group, the order of i is _____.

$*4*$

1.00000000

0.00000000

FBQ2

The degree and the leading coefficient of the polynomial $1 + x^3 + x^4 + 0.5x^5$ is _____.

$*(4, 1)*$

1.00000000

0.00000000

0.00000000

FBQ3

The degree of a polynomial written in this form $(\sum_{i=0}^n a_i x^i)$ if $a_n \neq 0$ is _____.

n

1.00000000

0.00000000

FBQ4

The order of (12) in S_3 is _____.

2

1.00000000

0.00000000

FBQ5

In a permutation, any cycle of length two is called _____.

Transposition

1.00000000

0.00000000

FBQ6

A field K is called _____ of F if F is a subfield of K , thus Q is a subfield of R and R is a field extension of Q

Field extension

1.00000000

0.00000000

FBQ7

A non - empty subset S of a field F is called a subfield of F if it is a field with respect to the operations on F . if $S \neq F$, then S is called _____ of F

Proper subfield

1.00000000

0.00000000

FBQ8

Let $f(x) = a_0 + a_1x + \dots + a_nx^n \in \mathbb{Z}[x]$. We define the content of $f(x)$ to be the g.c.d of the integers a_0, a_1, \dots, a_n , we say $f(x)$ is _____ if the content of $f(x) = 1$

primitive

1.00000000

0.00000000

FBQ9

We call an integral domain R a _____ if every non - zero element of R which is not a unit in R can be uniquely expressed as a product of a finite number of irreducible elements of R

Unique factorization domain

1.00000000

0.00000000

FBQ10

An element $d \in R$ is a _____ of $a, b \in R$ if

$d|a$ and $d|b$ and (i) i for any common divisor c of a and b , $c|d$

Greatest Common divisor

1.00000000

0.00000000

FBQ11

Given two elements a and b in a ring R , we say that $c \in R$ is a _____ of a and b if $c|a$ and $c|b$.

Common divisor

1.00000000

0.00000000

FBQ12

We call an integral domain R a _____ if every ideal in R is a principal ideal.

Principal ideal

1.00000000

0.00000000

FBQ13

The number of unit that can be obtained in $R = a+b-5 \mid a,b \in \mathbb{Z}$ is _____

2

1.00000000

0.00000000

FBQ14

Let R be an integral domain, an element $a \in R$ is called a unit or an _____ in R if we can find $b \in R$ such that $ab = 1$ i.e if a has a multiplicative inverse

Invertible element

1.00000000

0.00000000

FBQ15

A domain on which we can define a Euclidean valuation is called _____.

Euclidean domain

1.00000000

Euclidean

1.00000000

FBQ16

Let R be an integral domain. We say that a function $d: R \setminus \{0\} \rightarrow \mathbb{N} \cup \{0\}$ is a _____ on R if the following conditions are satisfied.

$d(a) \leq d(b) \forall a, b \in R \setminus \{0\}$ and

for any $a, b \in R, b \neq 0 \exists \quad q, r \in R$ such that $a = bq + r$, where $r = 0$ or $d(r) < d(b)$.

Euclidean Evaluation

1.00000000

0.00000000

FBQ17

Let F be a field and $f(x) \in F[x]$, we say that an element $a \in F$ is a _____ (where) m is positive integer of $f(x)$ if $(x-a)^m \mid f(x)$ but $(x-a)^{m+1} \nmid f(x)$

*Root of multiplicity m *

1.00000000

0.00000000

FBQ18

Let F be a field and $f(x) \in F[x]$ we say that an element $a \in F$ is a _____

(or zero) of $f(x)$ if $f(a) = 0$

Factor

1.00000000

Divides

1.00000000

FBQ19

If S is set, an object ' a ' in the collection S is called an _____ of S

Element

1.00000000

0.00000000

FBQ20

A set with _____ element in S is called an empty set

No

1.00000000

0.00000000

FBQ21

_____ method is sometimes used to list the element of a large set

Roster

1.00000000

0.00000000

FBQ22

The set of rational numbers and the set of real numbers are respectively represented by the symbol $\{ \#1 \}$ and $\{ \#2 \}$

10434 10435,10436

FBQ22 {100:SHORTANSWER:%100%Q}

Q

1.00000000

FBQ22 {100:SHORTANSWER:%100%R}

R

1.00000000

FBQ23

The symbol \exists denotes _____.

There exist

1.00000000

0.00000000

FBQ24

If A and B are two subsets of a set S , we can collect the element that are common to both A and B , we call this set the _____ of A and B .

Intersection

1.00000000

0.00000000

FBQ25

A relation R defined on a set S is said to be _____ if we have $aRa \forall a \in S$.

Reflexive

1.00000000

0.00000000

0.00000000

FBQ26

A relation R defined on a set S is said to be _____ if

$a R b \Rightarrow b R a \forall a, b \in S.$

Symmetric

1.00000000

0.00000000

FBQ27

A relation R defined on a set S is said to be _____ if $a R b$ and $b R a \forall a, b, c \in S$

Transitive

1.00000000

0.00000000

FBQ28

A relation R defined on a set S that is reflexive, symmetric and transitive is called _____ relation

Equivalence

1.00000000

0.00000000

FBQ29

A _____ f from a non - empty set A to a non - empty set B is a rule which associates with every element of A exactly one element of B

Function

1.00000000

0.00000000

FBQ30

A function $f : A \rightarrow B$ is called _____ if it associates different elements of A with different elements of B

Injective

1.00000000

One to one

1.00000000

FBQ31

A function $f : A \rightarrow B$ is called _____ if the range of f is B.

Onto

1.00000000

Surjective

1.00000000

FBQ32

Consider two non - empty sets A and B, we define the function $\pi_1 a, b = a$. π_1 is called the _____ of $A \times B$ onto A

Projection

1.00000000

0.00000000

FBQ33

A function that is both one to one and onto is called _____

Bijective

1.00000000

0.00000000

FBQ34

Any set which is equivalent to the set $1, 2, \dots, n$, for some $n \in \mathbb{N}$, is called a _____ set.

Finite

1.00000000

0.00000000

FBQ35

A set that is not _____ is called infinite set

Finite

1.00000000

0.00000000

FBQ36

A function $f : A \rightarrow B$ has an inverse if and only if is _____

Bijective

1.00000000

0.00000000

FBQ37

A natural number $p (\neq 1)$ is called _____ if its only divisor are 1 and p

Prime

1.00000000

0.00000000

FBQ38

If a natural number $n (\neq 1)$ is not a prime, then it is called a _____ number

Composite

1.00000000

0.00000000

FBQ39

Let A be any set, the function $I_A : A \rightarrow A : I_A a = a$ is called _____ on A .

Identity function

1.00000000

0.00000000

FBQ40

Let S be a non - empty set, any function $S \times S \rightarrow S$ is called a _____ on S .

Binary operation

1.00000000

0.00000000

FBQ41

Let $*$ be a binary operation on a set S . we say that: $*$ is _____ on a subset T of S if $a * b \in T \forall a, b \in T$

Closed
1.00000000

0.00000000
FBQ42

Let $*$ be a binary operation on a set S . we say that: $*$ is _____ if, for all $a, b, c \in S$, $(a*b)*c = a*(b*c)$.

Associative
1.00000000

0.00000000
FBQ43

Let $*$ be a binary operation on a set S . we say that: $*$ is _____ if for all $a, b \in S$, $a*b = b*a$

Commutative
1.00000000

0.00000000
FBQ44

If \circ and $*$ are two binary operations on a set S , we say that $*$ is _____.

Distributive over
1.00000000

0.00000000
FBQ45

.Let $*$ be a binary operation on a set S . if there is an element $e \in S$ such that $\forall a \in S$, $a * e = a$ and $e * a = a$ then e is called an _____ for $*$.

Identity element
1.00000000

0.00000000
FBQ46

The Cayley table is named after the famous mathematician

Arthur Cayley
1.00000000

0.00000000
FBQ47

_____ system consists of a set with a binary operation which satisfies certain properties is called a group

Algebraic
1.00000000

0.00000000
FBQ48

Let G be a group, for $a \in G$, we define

$a^0 = e$

$a^0 = a^{n-1}$, if $n > 0$

$a^{-n} = (a^{-1})^n$, if $n > 0$

n is called the exponent (or index) of _____ an of a

The integral power

1.00000000

integral power

1.00000000

FBQ49

\equiv is an equivalence relation, and hence partition Z into disjoint equivalence classes called _____ modulo n .

Congruence class

1.00000000

0.00000000

FBQ50

If the set X is finite, say $X = (1, 2, 3, \dots, n)$ then we denote $S(x)$ by S_n and each of S_n is called a _____ on n symbols

Permutation

1.00000000

0.00000000