



NATIONAL OPEN UNIVERSITY OF NIGERIA
14/16 AHMADU BELLO WAY, VICTORIA ISLAND, LAGOS
SCHOOL OF SCIENCE AND TECHNOLOGY
JUNE/JULY EXAMINATION

COURSE CODE: MTH402

COURSE TITLE: GENERAL TOPOLOGY II (3 units)

TIME ALLOWED: 3 HOURS

**INSTRUCTION: COMPLETE ANSWERS TO ANY FIVE (5)
QUESTIONS BEAR FULL MARKS**

- 1(a) Prove that the intersection $\tau = \bigcap_{\alpha} \tau_{\alpha}$ of topologies $\{ \tau_{\alpha} \}_{\alpha \in \Delta}$ on X is itself a topology in X (where Δ is some indexing set)
-7marks
- 1(b) Let X be a set, and let B be a basis for a topology τ on X . Show τ equals the collection of all unions of elements of B .
-7marks
- 2(a) Let B and B' be bases for the topology τ and τ' respectively on X . Show that the following are equivalent.
- i) τ' is finer than τ
 - ii) For each $x \in X$ and each basis element $B \in B$ such that $x \in B$, we know that $B \in \tau$ by definition and that $\tau \subset \tau'$, by condition (i) therefore $B \in \tau'$
- 7marks
- 2(b) Let d be a metric on the set X . show that the collection of all ε - balls $B_d(x, \varepsilon)$ for $x \in X$ and $\varepsilon > 0$ is a basis for a topology on X , called the metric topology induced by d . -7marks
- 3(a) State the properties under which d is a metric on X , given a function $d: X \times X \rightarrow \mathbb{R}$, for all $x, y, z \in X$.
-6marks
- 3(b) Let X and Y be two topological spaces. Let B be the collection of all sets of the form $U \times V$, where U is an open subset of X and V is an open subset of Y i.e. $B = \{ U \times V: U \text{ is open in } X \text{ and } V \text{ is open in } Y \}$. Show that B is the basis for a topology on $X \times Y$. -8marks
- 4(a) Let Y be a subspace of X . If U is open in Y and Y is open in X .

Show that U is open in X .

7marks

- 4(b) Show that the mapping $f: \mathbb{R} \rightarrow \mathbb{R}^+$ defined by $f(x) = e^x$ is a homeomorphism from \mathbb{R} to \mathbb{R}^+ . (Recall that a homeomorphism from one topological space to another is a bijective function f such that f and f^{-1} are both continuous)

-7marks

- 5(a) Give an example to show that every discrete space is Hausdorff.

-2½marks

- 5(b) What does it mean to say that a topological space is Hausdorff?

-2½marks

- 5(c) Let X be the Hausdorff space, then for all $x \in X$, show that the singleton set $\{x\}$ is closed.

-9marks

- 6(a) Let X and Y be topological spaces. When is a function $f: X \rightarrow Y$ said to be

continuous?

7marks

- 6(b) Let X be the subspace of \mathbb{R} given by $X = [0,1] \cup [2,4]$, Define $f: X \rightarrow \mathbb{R}$ by

$$f(x) = \begin{cases} x & \text{if } x \in [0,1] \\ x-1 & \text{if } x \in [2,4] \end{cases}$$

Prove that f is continuous.

-7marks

- 7(a) Let \mathbb{R} be endowed with standard topology. Show that for all $x \in \mathbb{R}$, $W = \{ (x - \epsilon, x + \epsilon) : \epsilon > 0 \}$ is a neighbourhood basis of x .

-7marks

- 7(b) Let X be a topological space and let $x \in X$. Suppose X is first countable, show that there exist a countable basis of x , say $W = \{ W_n : n \geq 1 \}$ such that $W_{n+1} \subset W_n$

-7marks