

Default for MTH251 Exams

The default category for questions shared in context 'MTH251 Exams'.
top

Default for MTH251

The default category for questions shared in context 'MTH251'.

Fill in the Blank (FBQs)

FBQ1

A vector is a quantity specified by ----- and -----

Magnitude and direction

1.00000000

Magnitude direction

1.00000000

Magnitude, direction

1.00000000

Magnitude/direction

1.00000000

FBQ2

The _____ vector coincides with the origin

Zero vector

1.00000000

Zero

1.00000000

FBQ3

Suppose $a=(a_x,a_y,a_z)$ and $b=(b_x,b_y,b_z)$ are two vectors, then the expression
 $a_x+b_x, a_y+b_y, a_z+b_z=$ ____

a+b

1.00000000

a + b

1.00000000

FBQ4

The three unit vectors i, j, k pointing in the directions of the x, y, z axis form
what is known as _____

Orthonormal triad

1.00000000

0.00000000

FBQ5

A vector whose sense is merely conventional, and would be reversed by changing
from a right-hand to a left-hand convention is called _____

Axial vector

1.00000000

0.00000000

FBQ6

The vector product of a vector with itself is the _____ vector

Zero

1.00000000

0

1.00000000

FBQ7

A scalar quantity is specified by _____

Direction

1.00000000

0.00000000

FBQ8

For a vector field A the expression $\nabla \cdot A = i\partial A_x \partial x + j\partial A_y \partial y + k\partial A_z \partial z$ is defined _____.

div A

1.00000000

0.00000000

FBQ9

$\nabla \times A = i\partial A_x \partial y - \partial A_y \partial z + j\partial A_x \partial z - \partial A_z \partial x + k(\partial A_y \partial x - \partial A_x \partial y)$ is the expression for _____

curl A

1.00000000

0.00000000

FBQ10

For two nonparallel vectors a and b drawn from O define a unique axis through O perpendicular to the plane containing a and b , the value $ab \sin \theta$ is the ____

Vector product

1.00000000

0.00000000

FBQ11

If θ is the angle between the vectors a and b , then the ____ of their sum is given by $a^2 + b^2 + 2ab \cos \theta$

Length

1.00000000

0.00000000

FBQ12

Any vector r can be written as a sum of three vectors along the three axes as $R =$ _____

* $x_i + y_j + x_k$ *

1.00000000

0.00000000

FBQ13

The basic equations of electromagnetic theory are _____ equations

Maxwell's

1.00000000

Maxwell

1.00000000

FBQ14

_____ force equation determines the force on a particle of charge q moving with velocity v .

Lorentz

1.00000000

Lorentz's

1.00000000

FBQ15

$A' = A - \nabla \Lambda$ is a ____ equation

Transformation

1.00000000

0.00000000

FBQ16

When there is no electric charge or current density, there are _____dimensional wave equations, which describe a wave propagating with velocity c

Three

1.00000000

3

1.00000000

FBQ17

For a static case, in which all the fields are time-independent; Maxwell's equations separate into a pair of _____equations

Electrostatic

1.00000000

0.00000000

FBQ18

The magnetic dipole field has precisely the same form as the _____dipole field

Electric

1.00000000

0.00000000

FBQ19

The first two members of a family of quantities known as tensors are _____ and _____

Scalars and vectors

1.00000000

Scalars, vectors

1.00000000

FBQ20

Scalars are called tensors of rank _____

Zero

1.00000000

0

1.00000000

FBQ21

The family of tensors of rank 2 are often called _____

Dyadic

1.00000000

0.00000000

FBQ22

Most frequently when one vector b is defined as a linear function of another vector a _____ occurs

Tensor

1.00000000

0.00000000

FBQ23

Tensors are commonly denoted by _____ capitals

Sans-serif capital

1.00000000

0.0000000

FBQ24

A tensor T has _____ components

Nine

1.0000000

9

1.0000000

FBQ25

A vector a is called an/a _____ of T if $Ta = \lambda a$

Eigenvector

1.0000000

0.0000000

FBQ26

For a vector a if $Ta = \lambda a$, λ is called the _____

Eigenvalue

1.0000000

0.0000000

FBQ27

Equivalently, $Ta = \lambda a$ may be written as $T - \lambda I a =$

0

1.0000000

0.0000000

FBQ28

When the density is a constant, the systems is said to be of _____

Uniform

1.0000000

0.0000000

FBQ29

A mass defined per unit volume is called _____ density

Volume

1.0000000

0.0000000

FBQ30

We can define a mass per unit length or linear density when the particles occupy a _____

Line

1.0000000

0.0000000

FBQ31

When _____ system of particles occupies a surface, we can define a surface density or mass per unit area

Continuous

1.0000000

0.0000000

FBQ32

Forces that change the distances between individual particles when applied to systems of particles are called _____ systems

Deformable

1.0000000

0.0000000

FBQ33

A system in which the distance between any two specified particles remains the same regardless of applied forces is called _____

Rigid body

1.0000000

0.0000000

FBQ34

The number of coordinates required to specify the position of a system of one or more particles is called the number of _____ of the system

Degrees of freedom

1.0000000

0.0000000

FBQ35

The number of degrees of freedom for five particles moving freely in a plane is _____

Ten

1.0000000

10

1.0000000

FBQ36

A system consisting of N particles moving freely in space requires 3N_____ to specify its position.

Coordinates

1.0000000

0.0000000

FBQ37

A constraint is _____ if the particle is constrained to move along a surface which is in a plane

Holonomic

1.0000000

0.0000000

FBQ38

The total linear impulse is equal to the change in linear _____

Motion

1.0000000

0.0000000

FBQ39

A rigid body which can move freely in space has _____ degrees of freedom

6

1.0000000

Six

1.0000000

FBQ40

In practice it is fairly simple to go from discrete to continuous systems by merely replacing summations by _____

Integration

1.00000000

0.00000000

FBQ41

If a system of particles is in a uniform gravitational field, the center of mass is sometimes called the _____

Centre of gravity

1.00000000

0.00000000

FBQ42

If $V = dr/dt = r \cdot v$ is the velocity of mv , the $p = \sum u = \sum mv = \sum u = \sum mv = r \cdot v$ defines the ____ of the system

Total momentum

1.00000000

0.00000000

FBQ43

The velocity v - of the ____ is given by $v = dr/dt$

Centre of mass

1.00000000

0.00000000

FBQ44

The total momentum of a system of particles can be found by multiplying the ____ M of the system by the velocity v -

Total mass

1.00000000

0.00000000

FBQ45

If the resultant external force acting on a system of particles is zero, then the total _____ remains constant

Momentum

1.00000000

0.00000000

FBQ46

The quantity $\Omega = \sum v = \sum mv(r \times v)$ is called the total _____ momentum of the system of particles about origin O

angular

1.00000000

0.00000000

FBQ47

The sum $\Lambda = \sum v = \sum Nrv(r \times Fv)$ is called the total external _____ about the origin

Torque

1.00000000

0.00000000

FBQ48

The total external torque on a system of particles is equal to the time rate of change of angular momentum of the system, provided that the internal forces between particles are _____ forces

Central

1.00000000

0.00000000

FBQ49

When all forces, external and internal, are conservative, we can define a total _____ energy V of the system.

potential

1.00000000

0.00000000

FBQ50

If T and V are respectively the total kinetic energy and total potential energy of a system of particles, then $T + V = \text{constant}$ is called the principle of _____ for systems of particles.

Conservation of energy

1.00000000

0.00000000

FBQ17

The first two members of a family of quantities known as tensors are $\{ \#1 \}$ and $\{ \#2 \}$

4689 4690,4691

FBQ17 {100:SHORTANSWER:%100%Scalars}

Scalars

1.00000000

FBQ17 {100:SHORTANSWER:%100%vectors}

vectors

1.00000000

Multiple Choice Questions (MCQs)

MCQ1

Geometrically, a vector is represented by _____

a line

0.00000000

a dot

0.00000000

a curve

0.00000000

an arrow

1.00000000

MCQ2

The scalar product of two vectors a and b is given by _____

$a \cdot b \cos \theta$

1.00000000

$(a+b) \cos \theta$

0.00000000

$ab \cos \theta$

0.00000000

$a \cdot b \sin \theta \cos \theta$

0.00000000

MCQ3

The vector product of a vector with itself is the _____ vector

polar

0.00000000

zero

1.00000000

axial

0.00000000

negative

0.00000000

MCQ4

_____force equation determines the force on a particle of charge q moving with velocity v .

Gauss

0.00000000

Lorenzo

0.00000000

Maxwell's

0.00000000

Lorentz

1.00000000

MCQ5

The transformation $A' = \nabla \Lambda$ is called a _____ transformation.

guage

1.00000000

guag

0.00000000

Gauss

0.00000000

grad

0.00000000

MCQ6

How many coordinates are required to specify the position of a rigid body which moves freely in space?

3

0.00000000

4

0.00000000

6

1.00000000

2

0.00000000

MCQ7

What is the scalar vector of the two vectors $a=(a_x, a_y, a_z)$ and $b=(b_x, b_y, b_z)$?

$a_x b_x + a_y b_y + a_z b_z$

1.00000000
axbz+aybx

0.00000000
axby+aybz

0.00000000
axbx-ayby-azbz

0.00000000
MCQ8

The magnetic dipole field has precisely the same form as the _____dipole field.
electromagnetic

0.00000000
electric

1.00000000
density

0.00000000
electrostatic

0.00000000
MCQ9

The first two members of a family of quantities known as tensors are _____ and _____.

scalars and tensor

0.00000000
vectors and magnitude

0.00000000
scalars and vectors

1.00000000
vectors and tensor

0.00000000
MCQ10

Tensors are commonly denoted by _____ capitals
small

0.00000000
sans

0.00000000
serif

0.00000000
sans-serif

1.00000000
MCQ11

A tensor Thas _____ components.

6

0.00000000
3

0.0000000
9

1.0000000
1

0.0000000
MCQ12

A vector a is called an/a _____ of T if $Ta = \lambda a$

eigenvector

1.0000000
eigenvalue

0.0000000
unit vector

0.0000000
null vector

0.0000000
MCQ13

Given that $Ta = \lambda a$, where T is a tensor, the number λ is called _____

eigenvector

0.0000000
eigenvalue

1.0000000
unit vector

0.0000000
null vector

0.0000000
MCQ14

$Ta = \lambda a$ can also be written as _____

$T - \lambda I = 0$

0.0000000
 $T - \lambda I a = 0$

1.0000000
 $T + \lambda I = 0$

0.0000000
 $T + \lambda a = 0$

0.0000000
MCQ15

The total momentum of a system of particles can be found by multiplying the total mass M of the system by the velocity v of the _____

center of mass

1.0000000
center of gravity

0.0000000
gravitational field

0.0000000
force of attraction

0.0000000
MCQ16
A systems is said to be of uniform density when the density is _____

uniform

0.0000000
dense

0.0000000
constant

1.0000000
normal

0.0000000
MCQ17
Forces applied to systems of particles will change the distances between individual particles. Such systems are often called ____ bodies

deformable

1.0000000
non-deformable

0.0000000
uniform

0.0000000
collapsible

0.0000000
MCQ18
Mass per unit volume is known as _____

continuous system

0.0000000
pressure

0.0000000
surface area

0.0000000
density

1.0000000
MCQ19
Vectors are tensors of rank _____

0

0.0000000
3

0.0000000
4

0.0000000
1

1.00000000

MCQ20

Dyadic are members of the tensor family of rank _____

0

0.00000000

3

0.00000000

2

1.00000000

1

0.00000000

MCQ21

Forces that can change the distances between individual particles of agiven systems are called _____

deformable forces

1.00000000

contact forces

0.00000000

continuous forces

0.00000000

discrete force

0.00000000

MCQ22

A system in which the distance between any two particles remains constant regardless of the applied forces is called _____

rigid body

1.00000000

free body

0.00000000

static body

0.00000000

polar body

0.00000000

MCQ23

The number of degree of freedom of a system is the number of _____ required to specify the position of a system of one or more particles.

forces

0.00000000

system

0.00000000

particles

0.00000000

coordinates

1.00000000

MCQ24

The vector function $\mathbf{r}(t) = x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k}$ from the origin to the particle is called the _____ vector

null

0.00000000
position

1.00000000
commutative

0.00000000
identity

0.00000000

MCQ25

What is the number of degrees of freedom for five particles moving freely in space?

15

1.00000000
10

0.00000000
2

0.00000000
1

0.00000000

MCQ26

A system of N particles moving freely in space requires _____ to specify its position.

2N coordinates

0.00000000
6N coordinates

0.00000000
3Ncoordinates

1.00000000
N coordinates

0.00000000

MCQ27

Surface density is defined as _____

mass per unit area

1.00000000
mass per unit volume

0.00000000
volume per unit area

0.00000000
pressure per unit area

0.00000000

MCQ28

A rigid body which can move freely in space has _____ degrees of freedom.

6

1.00000000

10

0.00000000

2

0.00000000

1

0.00000000

MCQ29

To go from discrete systems to continuous systems, we simply replace summations by _____

multiplications

0.00000000

coordinates

0.00000000

differentiations

0.00000000

integrations

1.00000000

MCQ30

The principle of conservation of momentum states that _____

the resultant of the external forces acting on a system of particles is zero

1.00000000

the sum of some of the external forces acting on a system of particles is zero

0.00000000

system of particles is under the state of motion under the action of an applied force

0.00000000

the resultant of the external forces acting on a system of particles is negligible

0.00000000

MCQ31

The center of mass for a body with uniform gravitational field is called _____

mass center

0.00000000

center of gravity

1.00000000

gravitational attraction

0.00000000

center of attraction

0.00000000

MCQ32

The number of degree of freedom of a particle moving freely in space is _____

5

0.00000000

3

1.00000000

2

0.00000000

1

0.00000000

MCQ33

The velocity v -of the center of mass of an object is given by _____

$v = dr/dt$

0.00000000

$v = ds/dt$

1.00000000

$v = dr/dt$

0.00000000

$v = dr/dv$

0.00000000

MCQ34

Which of the following is not correct?

Total momentum remains constant if the resultant external forces acting on a system of particles is zero

0.00000000

Total momentum is conserved if the resultant external forces acting on a system of particles is zero

0.00000000

If the resultant external forces acting on a system of particles is zero, the center of mass is either at rest or in motion with constant velocity

0.00000000

Total momentum is zero if the resultant external forces acting on a system of particles is conserved

1.00000000

MCQ35

The quantity

$\Omega = \sum v = 1/N m v r v \cdot V v$

Is called _____ of the system of particles about origin O.

the total momentum

0.00000000

the total circular momentum

0.00000000

the total angular momentum

1.00000000

the total angular moment

0.0000000

MCQ36

The total external torque on a system of particles is equal to the time rate of change of angular momentum of the system, provided that the internal forces between particles are _____ forces.

Triangular

0.0000000

circular

0.0000000

angular

0.0000000

central

1.0000000

MCQ37

The total linear impulse of force is equal to the change in linear _____

momentum

1.0000000

torque

0.0000000

energy

0.0000000

force

0.0000000

MCQ38

If T and V are respectively the total kinetic energy and total potential energy of a system of particles, then the formula $T + V = \text{constant}$ is called the _____ for systems of particles.

the principle of conservation of momentum

0.0000000

the principle of conservation of potential energy

0.0000000

the principle of conservation of energy

1.0000000

uniform energy

0.0000000

MCQ39

Assuming that the total mass of a system of particles is located at the center of mass O , then the total kinetic energy equals the kinetic energy of translation plus _____ about the center of mass.

the kinetic energy motion

1.0000000

the kinetic energy force

0.0000000

the kinetic energy momentum

0.00000000
the kinetic energy moment

0.00000000
MCQ40

Which of the following is true for rigid bodies and for motion on curves and surfaces without friction?

the virtual work of the constraint forces is zero

1.00000000
the virtual work of the constraint forces is negligible

0.00000000
the virtual work of the constraint forces is infinity

0.00000000
the virtual work of the constraint nonzero

0.00000000
MCQ41

If F is the total external force acting on a system of particles, then which of the following best describes the total linear impulse of the force?

$\int_{t_1}^{t_2} F dt$

0.00000000
 $\int_{t_1}^{t_2} F dt$

1.00000000
 $\int F dt$

0.00000000
 $\int_{t_1}^{t_2} dF$

0.00000000
MCQ42

Let θ is the angle between two vectors a and b , then the length of their sum $a+b$ is given by

$a^2+b^2+2ab \sin\theta$

0.00000000
 $a^2+b^2+2ab \cos\theta$

1.00000000
 $a^2-b^2+ab \cos\theta$

0.00000000
 $a+b+2ab \cos\theta$

0.00000000
MCQ43

The divergence of the vector $r = 3xyz^2i + 2xy^3j + x^2yzk$ at the point $(1, -1, 1)$ is

4

1.00000000
-4

0.00000000
3

0.0000000
5

0.0000000
MCQ44

What is the speed of a body whose position vector is $\mathbf{r} = \cos t \mathbf{i} + \sin t \mathbf{j} + t \mathbf{k}$?

$\sin t$

0.0000000
 $2\mathbf{j} + 2\mathbf{k}$

0.0000000
2

1.0000000
 $2t \mathbf{i} + 2t \mathbf{j}$

0.0000000
MCQ45

The displacement vector $\Delta \mathbf{r} = \mathbf{r}(t + \Delta t) - \mathbf{r}(t)$ is used to represent

change in speed

0.0000000
change in temperature

0.0000000
change in position

1.0000000
change in speed

0.0000000
MCQ46

A system of particles will be in stable equilibrium if the potential V of the system is _____

a maximum

0.0000000
a minimum.

1.0000000
negative

0.0000000
a torque

0.0000000
MCQ47

The three basic notions for analyzing motion are position, velocity, and _____

acceleration

1.0000000
momentum

0.0000000
viscosity

0.0000000
motion

0.00000000

MCQ48

The rate of change of velocity with respect to time is called the _____

acceleration

1.00000000

momentum

0.00000000

viscosity

0.00000000

motion

0.00000000

MCQ49

The speed v of a particle is defined to be the rate of change of distance along the path with respect to _____.

speed

0.00000000

rate

0.00000000

motion

0.00000000

time

1.00000000

MCQ50

Which is the acceleration of a body whose position vector is $\mathbf{r} = \cos t \mathbf{i} + \sin t \mathbf{j} + t \mathbf{k}$?

$\sin t \mathbf{j}$

0.00000000

$-\cos t \mathbf{i} + \sin t \mathbf{j}$

0.00000000

$-\cos t \mathbf{i} - \sin t \mathbf{j}$

1.00000000

$\cos t \mathbf{i} + \sin t \mathbf{j}$

0.00000000