

COLT— Exercise Sheet 6

Exercise 1. Let \mathcal{C} be a class of hypothesis from X to $\{0, 1\}$. Show that if \mathcal{C} is efficiently agnostic PAC-learnable, then it is efficiently PAC-learnable.

Exercise 2. Hoeffding's inequality states that if X_1, \dots, X_m are independent random variables with $a_i \leq X_i \leq b_i$ almost surely, then for every $\epsilon > 0$, for $S_m = \sum_{i=1}^m X_m$ and $\Delta^2 = \sum_{i=1}^m (b_i - a_i)^2$ we have

$$\begin{aligned}\mathbb{P}(S_m - \mathbb{E}(S_m) \geq \epsilon) &\leq e^{-\frac{2\epsilon^2}{\Delta^2}} \\ \mathbb{P}(S_m - \mathbb{E}(S_m) \leq -\epsilon) &\leq e^{-\frac{2\epsilon^2}{\Delta^2}}\end{aligned}$$

Consider \mathcal{H} being classifiers with the 0-1 loss. Use Hoeffding's inequality to estimate:

- The function $\mathbf{m}^R(\epsilon, \delta) = \min\{m \mid \mathbb{P}_{S \sim D^m}(\text{REP}_D(\mathcal{H}, S) \leq \epsilon) \geq 1 - \delta\}$ for $|\mathcal{H}| = 1$.
- The function $\mathbf{m}^R(\epsilon, \delta)$ for $|\mathcal{H}|$ finite.

Derive the estimate on the function $\epsilon^R(m, \delta) = \min\{\epsilon \mid \mathbf{m}^R(\epsilon, \delta) \leq m\}$.

Exercise 3. Generalize the above assignment to other loss functions. What assumptions do you need for that to work?

Exercise 4. Let D_1, \dots, D_m be a sequence of distributions over an example space X . Let \mathcal{C} be a finite hypothesis class over X . Fix $\mathbf{c} \in \mathcal{C}$. We consider samples S consisting of m examples, where each x_i is drawn with the distribution D_i (the examples are not independent), and then labeled with \mathbf{c} .

Now, for the evaluation we consider the average distribution:

$$D = \frac{D_1 + \dots + D_m}{m}.$$

That is we consider the generalisation error w.r.t. D :

$$\mathcal{R}_D(\mathbf{h}) = \mathbb{P}_{x \sim D}(\mathbf{c}(x) \neq \mathbf{h}(x)).$$

Show that for every $\epsilon > 0$, we have

$$\mathbb{P}_{S \sim (D_1, \dots, D_m)}\left(\exists \mathbf{h} \in \mathcal{C} (\mathcal{R}_D(\mathbf{h}) \geq \epsilon \wedge \widehat{\mathcal{R}}_S(\mathbf{h}) = 0)\right) \leq |\mathcal{C}| \cdot e^{-\epsilon \cdot m}.$$