

# Data-driven subgrid-scale parametrisation for a simple dynamical system

## Honours project update

Thomas Schanzer

School of Physics  
Climate Change Research Centre and ARC Centre of Excellence for Climate Extremes  
University of New South Wales, Sydney, Australia

Thursday 20 July 2023

The complexity of atmospheric models makes parametrisation testing difficult.

I aim to:

1. Use a simpler dynamical system as a test-bed
2. Build data-driven parametrisation schemes from scratch, with/without
  - i Stochasticity
  - ii Memory
3. Evaluate the schemes' performance

# Rayleigh-Bénard convection in 2D

$$\begin{aligned}\frac{\partial \mathbf{u}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{u} &= -\nabla \pi + \left(\frac{\text{Pr}}{\text{Ra}}\right)^{1/2} (1 + \nu^* |\nabla^2 \mathbf{u}|) \nabla^2 \mathbf{u} + \theta \hat{\mathbf{z}} \\ \frac{\partial \theta}{\partial t} + \mathbf{u} \cdot \nabla \theta &= (\text{Ra Pr})^{-1/2} (1 + \nu^* |\nabla^2 \theta|) \nabla^2 \theta \\ \nabla \cdot \mathbf{u} &= 0\end{aligned}$$

- Rayleigh number  $\text{Ra} = 10^9 \Rightarrow$  turbulent
- Prandtl number  $\text{Pr} = 1$
- **Nonlinear hyperdiffusion** for model stability at low resolution ( $\nu^* = 10^{-3}$ )
- No-slip, isothermal top/bottom boundaries; periodic lateral boundaries

# Numerical methods

- Equations solved using `dedalus v3`: spectral PDE solver for Python
  - Highly flexible
  - Designed to run in parallel
- Horizontal Fourier series expansion
- Vertical Chebyshev polynomial expansion

# Animation

Integration for 100 time units with 1024 horizontal and 128 vertical modes:

<https://youtu.be/SU8Co1UYY1A>

Key properties:

- Turbulent flow
- Large and small scale features
- Only two prognostic variables
- Comparatively inexpensive (13 cpu-hr for the above run)

⇒ A simple dynamical system analogue of the atmosphere

I need:

1. A reasonably accurate high-resolution model to treat as “truth”
2. A coarse-resolution model to correct using parametrisation
3. Evidence that under-resolution introduces biases that are:
  - i *Systematic*: consistent over- or under-estimation, not mere numerical error
  - ii *Statistically significant*

## Quantities of interest

Nusselt number (dimensionless rate of vertical heat transport):

$$\text{Nu}(z, t) \equiv (\text{Ra Pr})^{1/2} \langle w\theta \rangle_x - \frac{\partial \langle \theta \rangle_x}{\partial z}$$


Root-mean-square velocity:

$$u_{\text{rms}}(t) = \sqrt{\langle \mathbf{u} \cdot \mathbf{u} \rangle_{x,z}}$$

Mean values are obtained via:


1. 1000 time unit spin-up at coarse resolution
2. 300-time-unit integration at new resolution
3. Averaging over final 275 time units

# Mean Nusselt number





# Mean RMS velocity



## Next steps: building the parametrisation

1. Given the state  $S_t$  of the fine model at time  $t$ ,
  - i Advance it by  $\Delta t$  using the fine model  $\mathcal{F}_{\Delta t}$ , giving  $\mathcal{F}_{\Delta t}S_t$
  - ii Coarse-grain it and advance it by  $\Delta t$  using the coarse model  $\mathcal{C}_{\Delta t}$ , giving  $\mathcal{C}_{\Delta t}\langle S_t \rangle$
2. Determine the *unresolved tendency*

$$T = \frac{\langle \mathcal{F}_{\Delta t} S_t \rangle - \mathcal{C}_{\Delta t} \langle S_t \rangle}{\Delta t}$$

3. Fit a model to the dataset of  $(\langle S_t \rangle, T)$
4. Plug this into the coarse model as an external forcing and assess the results