

Table 2. Recommendation adherence

Recommendation adherence by recommendation window and day type						
	Recommendation	Observed minute-level adherence	Site-average recommendation-window model			
		Minutes meeting recommendation, n/N (%)	Work day, % (95% CI)	Free day, % (95% CI)	Free minus Work, percentage points (95% CI)	FDR-adjusted p
Daytime	At least 250 lx melanopic EDI during wake, excluding the three hours before sleep	137,792/573,712 (24.0%)	26.5 (24.4 to 28.6)	21.6 (19.3 to 23.9)	-4.9 (-7.6 to -2.2)	<0.001
Pre-sleep	No more than 10 lx melanopic EDI during the three hours before reported sleep	81,894/129,390 (63.3%)	61.9 (58.5 to 65.4)	69.9 (66.3 to 73.5)	+8.0 (+3.3 to +12.7)	<0.001
Sleep	No more than 1 lx melanopic EDI during reported sleep; bedside sleep environment	336,052/383,366 (87.7%)	90.3 (88.7 to 91.8)	84.1 (81.6 to 86.7)	-6.1 (-8.8 to -3.5)	<0.001

Observed pooled-minute fractions and site-average model estimates are distinct estimands. Each of the nine sites contributes equally to the model estimates.

Table 2: Recommendation adherence by Brown et al. recommendation window and day type. Daytime, Pre-sleep and Sleep identify the recommendation windows used for comparison. Daytime comprises diary-defined wake excluding the three hours before Sleep, Pre-sleep comprises those three hours, and Sleep uses the reported sleep interval and therefore describes the bedside sleep environment. Pooled minutes meeting recommendation report the number and percentage of valid measured minutes whose exposure lay within the applicable range. Work-day and free-day percentages, differences, 95% confidence intervals and false-discovery-rate-adjusted p-values are site-average estimates from the main recommendation-window model, with each of the nine sites weighted equally. The pooled-minute fractions and modelled recommendation-window estimates are distinct estimands. The main model included 2,216 recommendation-window periods from 140 participants and 782 behavioural cycles, representing 1,028,958 valid minutes. Adherence is defined at the measured-minute or recommendation-window-period level and does not classify participants.