

Table S11b. Light-exposure behaviour and awareness, part 2

	F2	F3	F4	F5
Midpoint of the brightest 10 hours	+0.049 h (-0.132–+0.229 h) FDR-adjusted p = 0.747	+0.133 h (-0.042–+0.309 h) FDR-adjusted p = 0.375	+0.030 h (-0.153–+0.213 h) FDR-adjusted p = 0.868	-0.083 h (-0.270–+0.105 h) FDR-adjusted p = 0.601
Midpoint of the darkest 10 hours	-0.136 h (-0.317–+0.044 h) FDR-adjusted p = 0.375	+0.080 h (-0.097–+0.258 h) FDR-adjusted p = 0.594	-0.020 h (-0.204–+0.164 h) FDR-adjusted p = 0.935	+0.132 h (-0.056–+0.320 h) FDR-adjusted p = 0.379
Mean timing of exposure above 250 lx melEDI	+0.020 h (-0.152–+0.192 h) FDR-adjusted p = 0.935	+0.086 h (-0.081–+0.253 h) FDR-adjusted p = 0.558	-0.097 h (-0.267–+0.074 h) FDR-adjusted p = 0.510	-0.166 h (-0.339–+0.008 h) FDR-adjusted p = 0.301
First light timing above 250 lx melEDI	-0.219 h (-0.489–+0.052 h) FDR-adjusted p = 0.362	+0.324 h (+0.062–+0.585 h) FDR-adjusted p = 0.179	+0.019 h (-0.252–+0.291 h) FDR-adjusted p = 0.958	-0.097 h (-0.380–+0.185 h) FDR-adjusted p = 0.679
Last light timing above 250 lx melEDI	+0.308 h (+0.043–+0.573 h) FDR-adjusted p = 0.202	-0.066 h (-0.328–+0.196 h) FDR-adjusted p = 0.760	-0.011 h (-0.280–+0.259 h) FDR-adjusted p = 0.958	-0.081 h (-0.361–+0.200 h) FDR-adjusted p = 0.743
melEDI dose	×1.278 (1.079–1.514) FDR-adjusted p = 0.124	×0.885 (0.748–1.048) FDR-adjusted p = 0.379	×0.872 (0.732–1.038) FDR-adjusted p = 0.375	×0.881 (0.736–1.054) FDR-adjusted p = 0.379
Melanopic daylight efficacy ratio	+0.004 (-0.011–+0.019) FDR-adjusted p = 0.747	-0.004 (-0.019–+0.011) FDR-adjusted p = 0.747	-0.010 (-0.026–+0.005) FDR-adjusted p = 0.380	-0.020 (-0.035–+0.004) FDR-adjusted p = 0.169

Associations are reported per one participant-level SD of the LEBA score with 95% CIs. FDR-adjusted p is bold only at alpha = 0.050 across all 68 tests. Sleep-environment cells are unfit for inference and suppressed.