

# A Simpler Prediction Rule for Rebound Hyperbilirubinemia

Pearl W. Chang, MD,<sup>a</sup> Thomas B. Newman, MD, MPH<sup>b,c,d</sup>

abstract

**OBJECTIVES:** We previously reported a clinical prediction rule to estimate the probability of rebound hyperbilirubinemia using gestational age (GA), age at phototherapy initiation, and total serum bilirubin (TSB) relative to the treatment threshold at phototherapy termination. We investigated (1) how a simpler 2-variable model would perform and (2) the absolute rebound risk if phototherapy were stopped at 2 mg/dL below the threshold for treatment initiation.

**METHODS:** Subjects for this retrospective cohort study were infants born 2012–2014 at  $\geq 35$  weeks' gestation at 1 of 17 Kaiser Permanente hospitals who underwent inpatient phototherapy before age 14 days. TSB reaching the phototherapy threshold within 72 hours of phototherapy termination was considered rebound. We simplified by using the difference between the TSB level at the time of phototherapy termination and the treatment threshold at the time of phototherapy initiation as 1 predictor, and kept GA as the other predictor.

**RESULTS:** Of the 7048 infants treated with phototherapy, 4.6% had rebound hyperbilirubinemia. The area under the receiver operating characteristic curve was 0.876 (95% confidence interval, 0.854 to 0.899) for the 2-variable model versus 0.881 (95% confidence interval, 0.859 to 0.903) for the 3-variable model. The rebound probability after stopping phototherapy at 2 mg/dL below the starting threshold was 2.5% for infants  $\geq 38$  weeks' GA and 10.2% for infants  $< 38$  weeks' GA.

**CONCLUSIONS:** Rebound hyperbilirubinemia can be predicted by a simpler 2-variable model consisting of GA and the starting threshold–ending TSB difference. Infants  $< 38$  weeks' gestation may need longer phototherapy because of their higher rebound risk.



<sup>a</sup>Department of Pediatrics, Seattle Children's Hospital, Seattle, Washington; Departments of <sup>b</sup>Epidemiology and Biostatistics and <sup>c</sup>Pediatrics, University of California, San Francisco, San Francisco, California; and <sup>d</sup>Division of Research, Kaiser Permanente Northern California, Oakland, California

Dr Chang conducted the statistical analysis and interpretation of data and drafted the initial manuscript; Dr Newman conceptualized and designed the study and guided the statistical analysis and interpretation of data; and both authors revised and reviewed the manuscript and approved the final manuscript as submitted.

**DOI:** <https://doi.org/10.1542/peds.2018-3712>

Accepted for publication Apr 16, 2019

Address correspondence to Pearl W. Chang, MD, Department of Pediatrics, Seattle Children's Hospital, Mailstop FA.2.115, PO Box 5371, Seattle, WA 98145-5005. E-mail: [pearlchangmd@gmail.com](mailto:pearlchangmd@gmail.com)

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2019 by the American Academy of Pediatrics

**FINANCIAL DISCLOSURE:** The authors have indicated they have no financial relationships relevant to this article to disclose.

**FUNDING:** No external funding.

**WHAT'S KNOWN ON THIS SUBJECT:** We previously reported a 3-variable clinical prediction rule that quantifies the risk of the bilirubin level rebounding to phototherapy levels after inpatient phototherapy.

**WHAT THIS STUDY ADDS:** We simplified the rule while maintaining excellent discrimination using as predictors only gestational age and the difference between the current bilirubin level and the treatment threshold at the time phototherapy was started.

**To cite:** Chang PW and Newman TB. A Simpler Prediction Rule for Rebound Hyperbilirubinemia. *Pediatrics*. 2019; 144(1):e20183712

Predicting rebound hyperbilirubinemia can help clinicians decide when to discontinue phototherapy for infants undergoing treatment of neonatal jaundice. We recently devised a 3-variable clinical prediction rule that quantifies the rebound risk after the first inpatient phototherapy according to an infant's gestational age (GA), age at phototherapy initiation, and total serum bilirubin (TSB) level relative to the American Academy of Pediatrics (AAP) phototherapy threshold at treatment termination.<sup>1</sup> An alternative approach is to discontinue phototherapy when the TSB is at least 2 mg/dL below the treatment threshold at the age of phototherapy initiation (M.J. Maisels, MB, BCh, DSc, personal communication, 2018). We sought to determine (1) if a 2-variable model, with just GA and the TSB relative to the threshold at phototherapy initiation, would perform as well as our previous 3-variable model and (2) the absolute risk of rebound if phototherapy were stopped as suggested.

## METHODS

### Study Cohort

Our retrospective cohort consisted of 7048 newborns born in Kaiser Permanente Northern California hospitals between 2012 and 2014 at  $\geq 35$  weeks' GA who underwent their first inpatient phototherapy before age 14 days. We excluded infants who did not have at least 2 TSB levels before phototherapy termination and infants with a conjugated bilirubin level of  $\geq 2$  mg/dL before or during their first phototherapy admission. Details of the cohort have been previously described.<sup>1</sup> In brief, 39.6% of the infants were  $< 38$  weeks' GA, and 14.5% had positive direct antiglobulin test (DAT) results. Phototherapy was initiated at a mean age of 2.3 (SD 1.3) days and terminated at a mean age of 3.6 (SD 1.3) days. The mean TSB level at

phototherapy termination was 9.8 (SD 2.7) mg/dL. Approximately 90% of infants had a TSB measurement after inpatient phototherapy termination. There was an order for home phototherapy equipment during the hospitalization for 4.4% of infants, who were thus assumed to have continued on home phototherapy after inpatient treatment. An additional 6.6% of infants had an order for home equipment after hospitalization and were thus assumed to have restarted on home phototherapy after discharge.

### Predictor and Outcome Variables

All of the variables were derived from electronic data sources.<sup>1</sup> We used the time of the first order for phototherapy as the time of phototherapy initiation. We estimated the time of phototherapy termination by using (1) nursing flowsheet documentation of discontinuation of phototherapy (25% of cohort); (2) if no such documentation was available, we used the time stamp of the discontinuation order (41%); and (3) if there was neither a nursing flowsheet nor order, we used the 1 hour before discharge time (34%). To estimate the TSB level at the time of phototherapy termination, we used the TSB level closest to the time of phototherapy discontinuation if 1 was measured between 3 hours before and 1 hour after treatment termination (24.6% of the cohort). If no TSB level was measured in this time window, we estimated a TSB at 3 hours before phototherapy termination by linear extrapolation using the last 2 TSB levels before termination. As previously reported, the average difference between an infant's extrapolated TSB value and his or her last measured TSB was 0.4 (SD 1.1) mg/dL. For this analysis, we subtracted the estimated TSB at phototherapy termination (ending TSB) from the AAP treatment threshold at phototherapy initiation (starting threshold) and used this

difference and the GA (dichotomized at  $< 38$  weeks) to form a 2-variable model. Rebound hyperbilirubinemia was defined as the TSB reaching or exceeding the hour-specific AAP phototherapy threshold within 72 hours of discontinuing phototherapy, as in our previous report. In our study cohort, 4.6% of infants had rebound hyperbilirubinemia.

### Statistical Analyses


To derive and validate this 2-variable prediction rule, we used the same random-split samples previously used to derive the 3-variable prediction rule. As before,<sup>1</sup> we summed the variables multiplied by 10 times their logistic regression coefficients to formulate a prediction score. Because the logistic coefficients are equal to the logarithm of the odds ratios, summing them is equivalent to multiplying their odds ratios. We assessed model fit in the validation data set with a calibration plot and the Hosmer-Lemeshow test (10 groups), and we assessed discrimination with the area under the receiver operating characteristic (AUROC) curve. We performed analyses using Stata 14.2 (Stata Corp, College Station, TX).

## RESULTS

Of the 7048 infants, the mean difference between the starting threshold and ending TSB was 4.4 (SD 3.5) mg/dL. The logistic coefficients from the derivation data set were 1.55 (95% confidence interval [CI], 1.02 to 2.08) for GA  $< 38$  weeks and  $-0.43$  per mg/dL (95% CI,  $-0.52$  to  $-0.34$ ) for the starting threshold–ending TSB difference. The equation for the score is thus:

$$\text{Score} = 15.5 \text{ (if GA } < 38 \text{ weeks)} \\ - 4.3 \times (\text{starting threshold} - \text{ending TSB})$$

The Hosmer-Lemeshow  $\chi^2$  (8 degrees of freedom) was 9.21



**FIGURE 1**

Calibration curve of the 2-variable prediction rule. Each point represents 1 decile of predicted probability of rebound.

( $P = .33$ ) in the validation data set ( $n = 3530$ ). Figure 1 shows the calibration curve on the validation data set. The discrimination of this 2-variable model was similar to that of the 3-variable model. In the derivation data set ( $n = 3518$ ), the AUROC of the 2-variable model was 0.877 (95% CI, 0.856 to 0.899) compared with 0.887 (95% CI, 0.864 to 0.910) in the 3-variable model. In the validation data set, the AUROC was 0.876 (95% CI, 0.854 to 0.899) compared with 0.881 (95% CI, 0.859 to 0.903). In the subset of infants with a measured TSB at phototherapy termination ( $n = 1737$ ), the AUROC was slightly higher at 0.901 (95% CI, 0.874 to 0.927).

The score can be translated into a predicted probability of rebound hyperbilirubinemia by consulting Fig 2. Alternatively, the following

equation can be programmed into a spreadsheet, mobile application, or Web site.

$$\begin{aligned} \text{Predicted probability of rebound} \\ = 1 / (1 \\ + \exp(-(0.097 * \text{score} - 2.84))) \end{aligned}$$


The mean difference of an infant's predicted probability of rebound hyperbilirubinemia using the 2 different models was 0% (SD 4.6%). Stopping phototherapy at 2 mg/dL below the starting threshold gave a rebound probability of 2.5% for infants  $\geq 38$  weeks' gestation and 10.2% for infants  $< 38$  weeks' gestation. For infants  $< 38$  weeks' gestation, phototherapy would need to be stopped at 5.5 mg/dL below the starting threshold to have a rebound probability (2.6%) similar to that of infants  $\geq 38$  weeks' GA who stopped phototherapy at 2 mg/dL below the starting threshold.

## DISCUSSION

We originally evaluated 10 variables (including race and ethnicity, DAT status, and formula and home phototherapy use) and by backward logistic regression devised a 3-variable clinical prediction rule that estimates the probability of rebound hyperbilirubinemia for an individual infant. That rule was based on (1) GA, (2) age at phototherapy initiation, and (3) TSB relative to the treatment threshold at phototherapy termination.<sup>1</sup>

Here, using the same large infant cohort, we turned the previous 3-variable model into a 2-variable model by using the difference between the starting threshold and ending TSB as 1 predictor, effectively combining the age at initiation with the relative TSB at termination. We kept GA as the other predictor. The AUROC of this simpler model was excellent and only minimally lower than that of the 3-variable model. Similarly, the calibration is excellent. The worst point on the calibration plot is when the predicted risk of rebound is  $\sim 5\%$  and the observed risk was 7%, a small absolute difference (Fig 1). However, calibration should be assessed in additional populations.

The rebound probability predicted by the 2-variable model may differ from that of the 3-variable model by up to  $\sim 5\%$ , more noticeably for infants  $< 38$  weeks' GA. Possible scenarios include the 2-variable model's estimates being lower for those who start phototherapy younger and have relatively short treatment durations and higher for those who stay on phototherapy longer. For example, if a DAT-negative infant of 37 weeks' gestation starts phototherapy at 24 hours of age and stops at 48 hours of age at an ending TSB of 7.9 (2 mg/dL below the starting threshold), then the predicted



**FIGURE 2**

Probability of rebound hyperbilirubinemia by score. Score =  $15.5 \text{ (if GA } < 38 \text{ weeks)} - 4.3 \times (\text{AAP phototherapy threshold at initiation} - \text{TSB at phototherapy termination})$ .

rebound probability would be ~10.2% on the basis of the new model compared with 15% on the basis of the previous model.

If phototherapy were discontinued at 2 mg/dL below the threshold for treatment initiation, the predicted probability of rebound is 2.5% for infants  $\geq 38$  weeks' gestation but 10.2% for infants  $< 38$  weeks' gestation. Lower GA has been well established as a risk factor for both hyperbilirubinemia and rebound hyperbilirubinemia.<sup>2-4</sup> In most previous studies of rebound hyperbilirubinemia, phototherapy was stopped when the TSB decreased below a fixed threshold or at the physicians' discretion.<sup>2,3,5,6</sup>

A randomized controlled trial from Israel compared discontinuing phototherapy at 3 vs 1 mg/dL below the threshold for phototherapy initiation. They reported no significant difference in the occurrence of rebound hyperbilirubinemia within ~24 hours.<sup>7</sup> However, 19% of infants in that study had rebound, which is higher than estimated by our prediction rule. The trial, however, had a small sample size ( $n = 52$ ), and 15% of the infants had glucose-6-phosphate dehydrogenase (G6PD) deficiency.<sup>7</sup> The prevalence of G6PD deficiency in our population is unknown but likely lower.<sup>8,9</sup> G6PD deficiency is

a known cause of hemolysis and severe hyperbilirubinemia<sup>10,11</sup> but has not been well characterized in rebound hyperbilirubinemia.

As before, this 2-variable prediction rule has several limitations.<sup>1,12</sup> Notably, it does not capture the rebound risk after the second episode of inpatient phototherapy, and because almost 97% of the cohort started treatment before age 5 days, it does not capture the rebound probability of older infants. The prediction rule also has not yet been externally validated.

## CONCLUSIONS

Rebound hyperbilirubinemia after inpatient phototherapy can be predicted by a simple 2-variable model consisting of GA and the difference between the starting treatment threshold and the ending TSB level. For infants  $< 38$  weeks' GA, it may be prudent to continue phototherapy longer because of their higher risk of rebound hyperbilirubinemia.

## ABBREVIATIONS

AAP: American Academy of Pediatrics  
AUROC: area under the receiver operating characteristic  
CI: confidence interval  
DAT: direct antiglobulin test  
GA: gestational age  
G6PD: glucose-6-phosphate dehydrogenase  
TSB: total serum bilirubin

**POTENTIAL CONFLICT OF INTEREST:** The authors have indicated they have no potential conflicts of interest to disclose.

## REFERENCES

- Chang PW, Kuzniewicz MW, McCulloch CE, Newman TB. A clinical prediction rule for rebound hyperbilirubinemia following inpatient phototherapy. *Pediatrics*. 2017;139(3):e20162896
- Kaplan M, Kaplan E, Hammerman C, et al. Post-phototherapy neonatal bilirubin rebound: a potential cause of

- significant hyperbilirubinaemia. *Arch Dis Child*. 2006;91(1):31–34
3. Bansal A, Jain S, Parmar VR, Chawla D. Bilirubin rebound after intensive phototherapy for neonatal jaundice. *Indian Pediatr*. 2010;47(7):607–609
  4. Stevenson DK, Maisels MJ, Watchko JF. *Care of the Jaundiced Neonate*. New York, NY: McGraw-Hill; 2012
  5. Maisels MJ, Kring E. Rebound in serum bilirubin level following intensive phototherapy. *Arch Pediatr Adolesc Med*. 2002;156(7):669–672
  6. Lazar L, Litwin A, Merlob P. Phototherapy for neonatal nonhemolytic hyperbilirubinemia. Analysis of rebound and indications for discontinuing therapy. *Clin Pediatr (Phila)*. 1993;32(5):264–267
  7. Barak M, Berger I, Dollberg S, Mimouni FB, Mandel D. When should phototherapy be stopped? A pilot study comparing two targets of serum bilirubin concentration. *Acta Paediatr*. 2009;98(2):277–281
  8. Wickremasinghe AC, Risley RJ, Kuzniewicz MW, et al. Risk of sensorineural hearing loss and bilirubin exchange transfusion thresholds. *Pediatrics*. 2015;136(3):505–512
  9. Nkhoma ET, Poole C, Vannappagari V, Hall SA, Beutler E. The global prevalence of glucose-6-phosphate dehydrogenase deficiency: a systematic review and meta-analysis. *Blood Cells Mol Dis*. 2009;42(3):267–278
  10. Kuzniewicz MW, Wickremasinghe AC, Wu YW, et al. Incidence, etiology, and outcomes of hazardous hyperbilirubinemia in newborns. *Pediatrics*. 2014;134(3):504–509
  11. Manning D, Todd P, Maxwell M, Jane Platt M. Prospective surveillance study of severe hyperbilirubinaemia in the newborn in the UK and Ireland. *Arch Dis Child Fetal Neonatal Ed*. 2007;92(5):F342–F346
  12. Chang PW. Authors' response. *Pediatrics*. 2017;140(2):e20171694B

## A Simpler Prediction Rule for Rebound Hyperbilirubinemia

Pearl W. Chang and Thomas B. Newman

*Pediatrics* 2019;144;

DOI: 10.1542/peds.2018-3712 originally published online June 13, 2019;

### Updated Information & Services

including high resolution figures, can be found at:  
<http://pediatrics.aappublications.org/content/144/1/e20183712>

### References

This article cites 11 articles, 5 of which you can access for free at:  
<http://pediatrics.aappublications.org/content/144/1/e20183712#BIBL>

### Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):  
**Fetus/Newborn Infant**  
[http://www.aappublications.org/cgi/collection/fetus:newborn\\_infant\\_sub](http://www.aappublications.org/cgi/collection/fetus:newborn_infant_sub)  
**Hyperbilirubinemia**  
[http://www.aappublications.org/cgi/collection/hyperbilirubinemia\\_sub](http://www.aappublications.org/cgi/collection/hyperbilirubinemia_sub)

### Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:  
<http://www.aappublications.org/site/misc/Permissions.xhtml>

### Reprints

Information about ordering reprints can be found online:  
<http://www.aappublications.org/site/misc/reprints.xhtml>

# American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **A Simpler Prediction Rule for Rebound Hyperbilirubinemia**

Pearl W. Chang and Thomas B. Newman

*Pediatrics* 2019;144;

DOI: 10.1542/peds.2018-3712 originally published online June 13, 2019;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/144/1/e20183712>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2019 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

## American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®

