

Today

Probability:

- Keep building it formally..

- And our intuition.

Poll: blows my mind.

Flip 300 million coins.

Poll: blows my mind.

Flip 300 million coins.

Which is more likely?

(A) 300 million heads.

(B) 300 million tails.

(C) Alternating heads and tails.

(D) A tail every third spot.

Poll: blows my mind.

Flip 300 million coins.

Which is more likely?

(A) 300 million heads.

(B) 300 million tails.

(C) Alternating heads and tails.

(D) A tail every third spot.

Given the history of the universe up to right now.

Poll: blows my mind.

Flip 300 million coins.

Which is more likely?

- (A) 300 million heads.
- (B) 300 million tails.
- (C) Alternating heads and tails.
- (D) A tail every third spot.

Given the history of the universe up to right now.

What is the likelihood of our universe?

- (A) The likelihood is 1. Cuz here it is.
- (B) As likely as any other. Cuz of probability.
- (C) Well. Quantum. IDK- TBH.

Poll: blows my mind.

Flip 300 million coins.

Which is more likely?

(A) 300 million heads.

(B) 300 million tails.

(C) Alternating heads and tails.

(D) A tail every third spot.

Given the history of the universe up to right now.

What is the likelihood of our universe?

(A) The likelihood is 1. Cuz here it is.

(B) As likely as any other. Cuz of probability.

(C) Well. Quantum. IDK- TBH.

Perhaps a philosophical (“wastebasket”) question.

Poll: blows my mind.

Flip 300 million coins.

Which is more likely?

- (A) 300 million heads.
- (B) 300 million tails.
- (C) Alternating heads and tails.
- (D) A tail every third spot.

Given the history of the universe up to right now.

What is the likelihood of our universe?

- (A) The likelihood is 1. Cuz here it is.
- (B) As likely as any other. Cuz of probability.
- (C) Well. Quantum. IDK- TBH.

Perhaps a philosophical (“wastebasket”) question.

Also, “cuz” == “because”

Probability Basics.

Probability Space.

Probability Basics.

Probability Space.

1. **Sample Space:** Set of outcomes, Ω .

Probability Basics.

Probability Space.

1. **Sample Space:** Set of outcomes, Ω .
2. **Probability:** $Pr[\omega]$ for all $\omega \in \Omega$.

Probability Basics.

Probability Space.

1. **Sample Space:** Set of outcomes, Ω .
2. **Probability:** $Pr[\omega]$ for all $\omega \in \Omega$.
 - 2.1 $0 \leq Pr[\omega] \leq 1$.

Probability Basics.

Probability Space.

1. **Sample Space:** Set of outcomes, Ω .

2. **Probability:** $Pr[\omega]$ for all $\omega \in \Omega$.

2.1 $0 \leq Pr[\omega] \leq 1$.

2.2 $\sum_{\omega \in \Omega} Pr[\omega] = 1$.

Probability Basics.

Probability Space.

1. **Sample Space:** Set of outcomes, Ω .

2. **Probability:** $Pr[\omega]$ for all $\omega \in \Omega$.

2.1 $0 \leq Pr[\omega] \leq 1$.

2.2 $\sum_{\omega \in \Omega} Pr[\omega] = 1$.

Example: Two coins.

1. $\Omega = \{HH, HT, TH, TT\}$

Probability Basics.

Probability Space.

1. **Sample Space:** Set of outcomes, Ω .

2. **Probability:** $Pr[\omega]$ for all $\omega \in \Omega$.

2.1 $0 \leq Pr[\omega] \leq 1$.

2.2 $\sum_{\omega \in \Omega} Pr[\omega] = 1$.

Example: Two coins.

1. $\Omega = \{HH, HT, TH, TT\}$

(Note: **Not** $\Omega = \{H, T\}$ with two picks!)

Probability Basics.

Probability Space.

1. **Sample Space:** Set of outcomes, Ω .
2. **Probability:** $Pr[\omega]$ for all $\omega \in \Omega$.
 - 2.1 $0 \leq Pr[\omega] \leq 1$.
 - 2.2 $\sum_{\omega \in \Omega} Pr[\omega] = 1$.

Example: Two coins.

1. $\Omega = \{HH, HT, TH, TT\}$
(Note: **Not** $\Omega = \{H, T\}$ with two picks!)
2. $Pr[HH] = \dots = Pr[TT] = 1/4$

Probability Basics.

Probability Space.

1. **Sample Space:** Set of outcomes, Ω .
2. **Probability:** $Pr[\omega]$ for all $\omega \in \Omega$.
 - 2.1 $0 \leq Pr[\omega] \leq 1$.
 - 2.2 $\sum_{\omega \in \Omega} Pr[\omega] = 1$.

Example: Two coins.

1. $\Omega = \{HH, HT, TH, TT\}$
(Note: **Not** $\Omega = \{H, T\}$ with two picks!)
2. $Pr[HH] = \dots = Pr[TT] = 1/4$

Consequences of Additivity

Theorem

Consequences of Additivity

Theorem

(a) Inclusion/Exclusion: $Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B]$;

Consequences of Additivity

Theorem

(a) **Inclusion/Exclusion:** $Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B]$;

(b) **Union Bound:** $Pr[A_1 \cup \dots \cup A_n] \leq Pr[A_1] + \dots + Pr[A_n]$;

Consequences of Additivity

Theorem

(a) Inclusion/Exclusion: $Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B]$;

(b) Union Bound: $Pr[A_1 \cup \dots \cup A_n] \leq Pr[A_1] + \dots + Pr[A_n]$;

(c) Law of Total Probability:

If A_1, \dots, A_N are a partition of Ω ,

Consequences of Additivity

Theorem

(a) Inclusion/Exclusion: $Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B]$;

(b) Union Bound: $Pr[A_1 \cup \dots \cup A_n] \leq Pr[A_1] + \dots + Pr[A_n]$;

(c) Law of Total Probability:

If A_1, \dots, A_N are a partition of Ω , i.e.,

pairwise disjoint and $\cup_{m=1}^N A_m = \Omega$,

Consequences of Additivity

Theorem

- (a) Inclusion/Exclusion: $Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B]$;
- (b) Union Bound: $Pr[A_1 \cup \dots \cup A_n] \leq Pr[A_1] + \dots + Pr[A_n]$;
- (c) Law of Total Probability:

If A_1, \dots, A_N are a partition of Ω , i.e.,
pairwise disjoint and $\cup_{m=1}^N A_m = \Omega$, then

$$Pr[B] = Pr[B \cap A_1] + \dots + Pr[B \cap A_N].$$

Consequences of Additivity

Theorem

(a) **Inclusion/Exclusion:** $Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B]$;

(b) **Union Bound:** $Pr[A_1 \cup \dots \cup A_n] \leq Pr[A_1] + \dots + Pr[A_n]$;

(c) **Law of Total Probability:**

If A_1, \dots, A_N are a **partition** of Ω , i.e.,

pairwise disjoint and $\cup_{m=1}^N A_m = \Omega$, then

$$Pr[B] = Pr[B \cap A_1] + \dots + Pr[B \cap A_N].$$

Proof Idea: Total probability.

Consequences of Additivity


Theorem

- (a) **Inclusion/Exclusion:** $Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B]$;
- (b) **Union Bound:** $Pr[A_1 \cup \dots \cup A_n] \leq Pr[A_1] + \dots + Pr[A_n]$;
- (c) **Law of Total Probability:**

If A_1, \dots, A_N are a **partition** of Ω , i.e.,
pairwise disjoint and $\cup_{m=1}^N A_m = \Omega$, then

$$Pr[B] = Pr[B \cap A_1] + \dots + Pr[B \cap A_N].$$

Proof Idea: Total probability.



Add it up!

Add it up. Poll.

What does Rao mean by “Add it up.”

Add it up. Poll.

What does Rao mean by “Add it up.”

- (A) Organize intuitions/proofs around $Pr[\omega]$.
- (B) Organize intuition/proofs around $Pr[A]$.
- (C) Some weird song whose refrain he heard in his youth.

Add it up. Poll.

What does Rao mean by “Add it up.”

(A) Organize intuitions/proofs around $Pr[\omega]$.

(B) Organize intuition/proofs around $Pr[A]$.


(C) Some weird song whose refrain he heard in his youth.

(A), (B), and (C)

Conditional Probability.

Definition: The **conditional probability** of B given A is

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}$$




Note also:

$$Pr[A \cap B] = Pr[B|A]Pr[A]$$

Conditional Probability.

Definition: The **conditional probability** of B given A is

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}$$



In A !


Note also:

$$Pr[A \cap B] = Pr[B|A]Pr[A]$$

Conditional Probability.

Definition: The **conditional probability** of B given A is

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}$$



In A !

In B ?


Note also:

$$Pr[A \cap B] = Pr[B|A]Pr[A]$$

Conditional Probability.

Definition: The **conditional probability** of B given A is

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}$$



In A !
In B ?
Must be in $A \cap B$.


Note also:

$$Pr[A \cap B] = Pr[B|A]Pr[A]$$

Conditional Probability.

Definition: The **conditional probability** of B given A is

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}$$




In A !
In B ?
Must be in $A \cap B$.

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}.$$

Conditional Probability.

Definition: The **conditional probability** of B given A is

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}$$



In A !

In B ?

Must be in $A \cap B$.


$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}.$$

Note also:

Conditional Probability.

Definition: The **conditional probability** of B given A is

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}$$



In A !
In B ?
Must be in $A \cap B$.

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}.$$

Note also:

$$Pr[A \cap B] = Pr[B|A]Pr[A]$$

Product Rule

Def: $Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}.$

Product Rule

Def: $Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}$.

Also: $Pr[A \cap B] = Pr[B|A]Pr[A]$

Product Rule

Def: $Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}$.

Also: $Pr[A \cap B] = Pr[B|A]Pr[A]$

Theorem Product Rule

Let A_1, A_2, \dots, A_n be events. Then

Product Rule

Def: $Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}$.

Also: $Pr[A \cap B] = Pr[B|A]Pr[A]$

Theorem Product Rule

Let A_1, A_2, \dots, A_n be events. Then

$$Pr[A_1 \cap \dots \cap A_n] = Pr[A_1]Pr[A_2|A_1] \dots Pr[A_n|A_1 \cap \dots \cap A_{n-1}].$$

Simple Bayes Rule.

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}, \quad Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}.$$

Simple Bayes Rule.

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}, \quad Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}.$$

$$Pr[A \cap B] = Pr[A|B]Pr[B] = Pr[B|A]Pr[A].$$

Simple Bayes Rule.

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}, \quad Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}.$$

$$Pr[A \cap B] = Pr[A|B]Pr[B] = Pr[B|A]Pr[A].$$

$$\text{Bayes Rule: } Pr[A|B] = \frac{Pr[B|A]Pr[A]}{Pr[B]}.$$

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair',

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

We know $P[B|A] =$

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

We know $P[B|A] = 1/2$, $P[B|\bar{A}] =$

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

We know $P[B|A] = 1/2$, $P[B|\bar{A}] = 0.6$,

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

We know $P[B|A] = 1/2$, $P[B|\bar{A}] = 0.6$, $Pr[A] =$

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

We know $P[B|A] = 1/2$, $P[B|\bar{A}] = 0.6$, $Pr[A] = 1/2$

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

We know $P[B|A] = 1/2$, $P[B|\bar{A}] = 0.6$, $Pr[A] = 1/2 = Pr[\bar{A}]$

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

We know $P[B|A] = 1/2$, $P[B|\bar{A}] = 0.6$, $Pr[A] = 1/2 = Pr[\bar{A}]$

Now,

$$Pr[B] = Pr[A \cap B] + Pr[\bar{A} \cap B] =$$

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

We know $P[B|A] = 1/2$, $P[B|\bar{A}] = 0.6$, $Pr[A] = 1/2 = Pr[\bar{A}]$

Now,

$$Pr[B] = Pr[A \cap B] + Pr[\bar{A} \cap B] = Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}]$$

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

We know $P[B|A] = 1/2$, $P[B|\bar{A}] = 0.6$, $Pr[A] = 1/2 = Pr[\bar{A}]$

Now,

$$\begin{aligned} Pr[B] &= Pr[A \cap B] + Pr[\bar{A} \cap B] = Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}] \\ &= (1/2)(1/2) + (1/2)0.6 = 0.55. \end{aligned}$$

Is your coin loaded?

Your coin is fair w.p. $1/2$ or such that $Pr[H] = 0.6$, otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

Analysis:

$A =$ 'coin is fair', $B =$ 'outcome is heads'

We want to calculate $P[A|B]$.

We know $P[B|A] = 1/2$, $P[B|\bar{A}] = 0.6$, $Pr[A] = 1/2 = Pr[\bar{A}]$

Now,

$$\begin{aligned} Pr[B] &= Pr[A \cap B] + Pr[\bar{A} \cap B] = Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}] \\ &= (1/2)(1/2) + (1/2)0.6 = 0.55. \end{aligned}$$

Thus,

$$Pr[A|B] = \frac{Pr[A]Pr[B|A]}{Pr[B]} = \frac{(1/2)(1/2)}{(1/2)(1/2) + (1/2)0.6} \approx 0.45.$$

Independence

Definition: Two events A and B are **independent** if

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, A = sum is 7 and B = red die is 1 are

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, $A = \text{sum is 7}$ and $B = \text{red die is 1}$ are independent;

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, $A = \text{sum is 7}$ and $B = \text{red die is 1}$ are independent; $Pr[A \cap B] = \frac{1}{36}$, $Pr[A]Pr[B] = \left(\frac{1}{6}\right)\left(\frac{1}{6}\right)$.

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, $A = \text{sum is 7}$ and $B = \text{red die is 1}$ are independent; $Pr[A \cap B] = \frac{1}{36}$, $Pr[A]Pr[B] = \left(\frac{1}{6}\right)\left(\frac{1}{6}\right)$.
- ▶ When rolling two dice, $A = \text{sum is 3}$ and $B = \text{red die is 1}$ are

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, $A = \text{sum is 7}$ and $B = \text{red die is 1}$ are independent; $Pr[A \cap B] = \frac{1}{36}$, $Pr[A]Pr[B] = \left(\frac{1}{6}\right)\left(\frac{1}{6}\right)$.
- ▶ When rolling two dice, $A = \text{sum is 3}$ and $B = \text{red die is 1}$ are **not** independent;

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, $A = \text{sum is 7}$ and $B = \text{red die is 1}$ are independent; $Pr[A \cap B] = \frac{1}{36}$, $Pr[A]Pr[B] = (\frac{1}{6})(\frac{1}{6})$.
- ▶ When rolling two dice, $A = \text{sum is 3}$ and $B = \text{red die is 1}$ are **not** independent; $Pr[A \cap B] = \frac{1}{36}$, $Pr[A]Pr[B] = (\frac{2}{36})(\frac{1}{6})$.

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, A = sum is 7 and B = red die is 1 are independent; $Pr[A \cap B] = \frac{1}{36}$, $Pr[A]Pr[B] = (\frac{1}{6})(\frac{1}{6})$.
- ▶ When rolling two dice, A = sum is 3 and B = red die is 1 are **not** independent; $Pr[A \cap B] = \frac{1}{36}$, $Pr[A]Pr[B] = (\frac{2}{36})(\frac{1}{6})$.
- ▶ When flipping coins, A = coin 1 yields heads and B = coin 2 yields tails are

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, A = sum is 7 and B = red die is 1 are independent; $Pr[A \cap B] = \frac{1}{36}$, $Pr[A]Pr[B] = (\frac{1}{6})(\frac{1}{6})$.
- ▶ When rolling two dice, A = sum is 3 and B = red die is 1 are **not** independent; $Pr[A \cap B] = \frac{1}{36}$, $Pr[A]Pr[B] = (\frac{2}{36})(\frac{1}{6})$.
- ▶ When flipping coins, A = coin 1 yields heads and B = coin 2 yields tails are independent;

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, A = sum is 7 and B = red die is 1 are independent; $Pr[A \cap B] = \frac{1}{36}$, $Pr[A]Pr[B] = (\frac{1}{6})(\frac{1}{6})$.
- ▶ When rolling two dice, A = sum is 3 and B = red die is 1 are **not** independent; $Pr[A \cap B] = \frac{1}{36}$, $Pr[A]Pr[B] = (\frac{2}{36})(\frac{1}{6})$.
- ▶ When flipping coins, A = coin 1 yields heads and B = coin 2 yields tails are independent; $Pr[A \cap B] = \frac{1}{4}$, $Pr[A]Pr[B] = (\frac{1}{2})(\frac{1}{2})$.

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, A = sum is 7 and B = red die is 1 are independent; $Pr[A \cap B] = \frac{1}{36}$, $Pr[A]Pr[B] = (\frac{1}{6})(\frac{1}{6})$.
- ▶ When rolling two dice, A = sum is 3 and B = red die is 1 are **not** independent; $Pr[A \cap B] = \frac{1}{36}$, $Pr[A]Pr[B] = (\frac{2}{36})(\frac{1}{6})$.
- ▶ When flipping coins, A = coin 1 yields heads and B = coin 2 yields tails are independent; $Pr[A \cap B] = \frac{1}{4}$, $Pr[A]Pr[B] = (\frac{1}{2})(\frac{1}{2})$.
- ▶ When throwing 3 balls into 3 bins, A = bin 1 is empty and B = bin 2 is empty are

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, A = sum is 7 and B = red die is 1 are independent; $Pr[A \cap B] = \frac{1}{36}$, $Pr[A]Pr[B] = (\frac{1}{6})(\frac{1}{6})$.
- ▶ When rolling two dice, A = sum is 3 and B = red die is 1 are **not** independent; $Pr[A \cap B] = \frac{1}{36}$, $Pr[A]Pr[B] = (\frac{2}{36})(\frac{1}{6})$.
- ▶ When flipping coins, A = coin 1 yields heads and B = coin 2 yields tails are independent; $Pr[A \cap B] = \frac{1}{4}$, $Pr[A]Pr[B] = (\frac{1}{2})(\frac{1}{2})$.
- ▶ When throwing 3 balls into 3 bins, A = bin 1 is empty and B = bin 2 is empty are **not** independent; $Pr[A \cap B] = \frac{1}{27}$, $Pr[A]Pr[B] = (\frac{8}{27})(\frac{8}{27})$.

Independence

Definition: Two events A and B are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice, A = sum is 7 and B = red die is 1 are independent; $Pr[A \cap B] = \frac{1}{36}$, $Pr[A]Pr[B] = (\frac{1}{6})(\frac{1}{6})$.
- ▶ When rolling two dice, A = sum is 3 and B = red die is 1 are **not** independent; $Pr[A \cap B] = \frac{1}{36}$, $Pr[A]Pr[B] = (\frac{2}{36})(\frac{1}{6})$.
- ▶ When flipping coins, A = coin 1 yields heads and B = coin 2 yields tails are independent; $Pr[A \cap B] = \frac{1}{4}$, $Pr[A]Pr[B] = (\frac{1}{2})(\frac{1}{2})$.
- ▶ When throwing 3 balls into 3 bins, A = bin 1 is empty and B = bin 2 is empty are **not** independent; $Pr[A \cap B] = \frac{1}{27}$, $Pr[A]Pr[B] = (\frac{8}{27})(\frac{8}{27})$.

Independence and conditional probability

Fact: Two events A and B are **independent** if and only if

Independence and conditional probability

Fact: Two events A and B are **independent** if and only if

$$Pr[A|B] = Pr[A].$$

Independence and conditional probability

Fact: Two events A and B are **independent** if and only if

$$Pr[A|B] = Pr[A].$$

Indeed:

Independence and conditional probability

Fact: Two events A and B are **independent** if and only if

$$Pr[A|B] = Pr[A].$$

Indeed: $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$, so that

Independence and conditional probability

Fact: Two events A and B are **independent** if and only if

$$Pr[A|B] = Pr[A].$$

Indeed: $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$, so that

$$Pr[A|B] = Pr[A] \Leftrightarrow \frac{Pr[A \cap B]}{Pr[B]} = Pr[A]$$

Independence and conditional probability

Fact: Two events A and B are **independent** if and only if

$$Pr[A|B] = Pr[A].$$

Indeed: $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$, so that

$$Pr[A|B] = Pr[A] \Leftrightarrow \frac{Pr[A \cap B]}{Pr[B]} = Pr[A] \Leftrightarrow Pr[A \cap B] = Pr[A]Pr[B].$$

Conditional Probability: Review

Conditional Probability: Review

Recall:

$$\blacktriangleright \Pr[A|B] = \frac{\Pr[A \cap B]}{\Pr[B]}.$$

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A],$

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,
i.e., if $Pr[A \cap B] < Pr[A]Pr[B].$

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,
i.e., if $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶ A and B are *independent* if $Pr[A|B] = Pr[A],$

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,
i.e., if $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶ A and B are *independent* if $Pr[A|B] = Pr[A]$,
i.e., if $Pr[A \cap B] = Pr[A]Pr[B].$

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,
i.e., if $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶ A and B are *independent* if $Pr[A|B] = Pr[A]$,
i.e., if $Pr[A \cap B] = Pr[A]Pr[B].$
- ▶ Note: $B \subset A \Rightarrow A$ and B are

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,
i.e., if $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶ A and B are *independent* if $Pr[A|B] = Pr[A]$,
i.e., if $Pr[A \cap B] = Pr[A]Pr[B].$
- ▶ Note: $B \subset A \Rightarrow A$ and B are positively correlated.

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,
i.e., if $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶ A and B are *independent* if $Pr[A|B] = Pr[A]$,
i.e., if $Pr[A \cap B] = Pr[A]Pr[B].$
- ▶ Note: $B \subset A \Rightarrow A$ and B are positively correlated.
($Pr[A|B] = 1 > Pr[A]$)

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,
i.e., if $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶ A and B are *independent* if $Pr[A|B] = Pr[A]$,
i.e., if $Pr[A \cap B] = Pr[A]Pr[B].$
- ▶ Note: $B \subset A \Rightarrow A$ and B are positively correlated.
($Pr[A|B] = 1 > Pr[A]$)
- ▶ Note: $A \cap B = \emptyset \Rightarrow A$ and B are

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,
i.e., if $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶ A and B are *independent* if $Pr[A|B] = Pr[A]$,
i.e., if $Pr[A \cap B] = Pr[A]Pr[B].$
- ▶ Note: $B \subset A \Rightarrow A$ and B are positively correlated.
($Pr[A|B] = 1 > Pr[A]$)
- ▶ Note: $A \cap B = \emptyset \Rightarrow A$ and B are negatively correlated.

Conditional Probability: Review




Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,
i.e., if $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶ A and B are *independent* if $Pr[A|B] = Pr[A]$,
i.e., if $Pr[A \cap B] = Pr[A]Pr[B].$
- ▶ Note: $B \subset A \Rightarrow A$ and B are positively correlated.
($Pr[A|B] = 1 > Pr[A]$)
- ▶ Note: $A \cap B = \emptyset \Rightarrow A$ and B are negatively correlated.
($Pr[A|B] = 0 < Pr[A]$)

Conditional Probability: Pictures/Poll.


Conditional Probability: Pictures/Poll.

Illustrations: Pick a point uniformly in the unit square



Conditional Probability: Pictures/Poll.


Illustrations: Pick a point uniformly in the unit square



Which A and B are independent?

Conditional Probability: Pictures/Poll.

Illustrations: Pick a point uniformly in the unit square




Which A and B are independent?

- (A) Left.
- (B) Middle.
- (B) Right.

Conditional Probability: Pictures/Poll.

Illustrations: Pick a point uniformly in the unit square



Which A and B are independent?




- (A) Left.
- (B) Middle.
- (B) Right.

See next slide.

Conditional Probability: Pictures




Conditional Probability: Pictures

Illustrations: Pick a point uniformly in the unit square



Conditional Probability: Pictures




Illustrations: Pick a point uniformly in the unit square



► Left: A and B are

Conditional Probability: Pictures




Illustrations: Pick a point uniformly in the unit square



- Left: A and B are independent.

Conditional Probability: Pictures




Illustrations: Pick a point uniformly in the unit square



- Left: A and B are independent. $Pr[B] =$

Conditional Probability: Pictures




Illustrations: Pick a point uniformly in the unit square



- Left: A and B are independent. $Pr[B] = b$;

Conditional Probability: Pictures




Illustrations: Pick a point uniformly in the unit square



- Left: A and B are independent. $Pr[B] = b$; $Pr[B|A] =$

Conditional Probability: Pictures




Illustrations: Pick a point uniformly in the unit square



- Left: A and B are independent. $Pr[B] = b$; $Pr[B|A] = b$.

Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



- ▶ Left: A and B are independent. $Pr[B] = b$; $Pr[B|A] = b$.
- ▶ Middle: A and B are

Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



- ▶ Left: A and B are independent. $Pr[B] = b$; $Pr[B|A] = b$.
- ▶ Middle: A and B are positively correlated.

Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



- ▶ Left: A and B are independent. $Pr[B] = b$; $Pr[B|A] = b$.
- ▶ Middle: A and B are positively correlated.
 $Pr[B|A] =$

Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



- ▶ Left: A and B are independent. $Pr[B] = b$; $Pr[B|A] = b$.
- ▶ Middle: A and B are positively correlated.
 $Pr[B|A] = b_1 > Pr[B|\bar{A}] =$

Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



- ▶ Left: A and B are independent. $Pr[B] = b$; $Pr[B|A] = b$.
- ▶ Middle: A and B are positively correlated.
 $Pr[B|A] = b_1 > Pr[B|\bar{A}] = b_2$.

Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



- ▶ Left: A and B are independent. $Pr[B] = b$; $Pr[B|A] = b$.
- ▶ Middle: A and B are positively correlated.
 $Pr[B|A] = b_1 > Pr[B|\bar{A}] = b_2$. Note: $Pr[B] \in (b_2, b_1)$.

Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



- ▶ Left: A and B are independent. $Pr[B] = b$; $Pr[B|A] = b$.
- ▶ Middle: A and B are positively correlated.
 $Pr[B|A] = b_1 > Pr[B|\bar{A}] = b_2$. Note: $Pr[B] \in (b_2, b_1)$.
- ▶ Right: A and B are

Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



- ▶ Left: A and B are independent. $Pr[B] = b$; $Pr[B|A] = b$.
- ▶ Middle: A and B are positively correlated.
 $Pr[B|A] = b_1 > Pr[B|\bar{A}] = b_2$. Note: $Pr[B] \in (b_2, b_1)$.
- ▶ Right: A and B are negatively correlated.

Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



- ▶ Left: A and B are independent. $Pr[B] = b$; $Pr[B|A] = b$.
- ▶ Middle: A and B are positively correlated.
 $Pr[B|A] = b_1 > Pr[B|\bar{A}] = b_2$. Note: $Pr[B] \in (b_2, b_1)$.
- ▶ Right: A and B are negatively correlated.
 $Pr[B|A] = b_1 < Pr[B|\bar{A}] = b_2$.

Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square




- ▶ Left: A and B are independent. $Pr[B] = b$; $Pr[B|A] = b$.
- ▶ Middle: A and B are positively correlated.
 $Pr[B|A] = b_1 > Pr[B|\bar{A}] = b_2$. Note: $Pr[B] \in (b_2, b_1)$.
- ▶ Right: A and B are negatively correlated.
 $Pr[B|A] = b_1 < Pr[B|\bar{A}] = b_2$. Note: $Pr[B] \in (b_1, b_2)$.

Bayes and Biased Coin

Bayes and Biased Coin




Bayes and Biased Coin



Pick a point uniformly at random in the unit square. Then


Bayes and Biased Coin



Pick a point uniformly at random in the unit square. Then

$$Pr[A] =$$


Bayes and Biased Coin



Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5;$$


Bayes and Biased Coin



Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] =$$


Bayes and Biased Coin



Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

Bayes and Biased Coin




Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] =$$

Bayes and Biased Coin




Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5;$$

Bayes and Biased Coin




Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] =$$

Bayes and Biased Coin




Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6;$$

Bayes and Biased Coin




Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6; Pr[A \cap B] =$$

Bayes and Biased Coin




Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6; Pr[A \cap B] = 0.5 \times 0.5$$

Bayes and Biased Coin




Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6; Pr[A \cap B] = 0.5 \times 0.5$$

$$Pr[B] =$$

Bayes and Biased Coin




Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6; Pr[A \cap B] = 0.5 \times 0.5$$

$$Pr[B] = 0.5 \times 0.5 + 0.5 \times 0.6$$

Bayes and Biased Coin




Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6; Pr[A \cap B] = 0.5 \times 0.5$$

$$Pr[B] = 0.5 \times 0.5 + 0.5 \times 0.6 = Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}]$$

Bayes and Biased Coin



Pick a point uniformly at random in the unit square. Then


$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6; Pr[A \cap B] = 0.5 \times 0.5$$

$$Pr[B] = 0.5 \times 0.5 + 0.5 \times 0.6 = Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}]$$

$$Pr[A|B] = \frac{0.5 \times 0.5}{0.5 \times 0.5 + 0.5 \times 0.6}$$

Bayes and Biased Coin



Pick a point uniformly at random in the unit square. Then


$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6; Pr[A \cap B] = 0.5 \times 0.5$$

$$Pr[B] = 0.5 \times 0.5 + 0.5 \times 0.6 = Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}]$$

$$Pr[A|B] = \frac{0.5 \times 0.5}{0.5 \times 0.5 + 0.5 \times 0.6} = \frac{Pr[A]Pr[B|A]}{Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}]}$$

Bayes and Biased Coin



Pick a point uniformly at random in the unit square. Then


$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6; Pr[A \cap B] = 0.5 \times 0.5$$

$$Pr[B] = 0.5 \times 0.5 + 0.5 \times 0.6 = Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}]$$

$$Pr[A|B] = \frac{0.5 \times 0.5}{0.5 \times 0.5 + 0.5 \times 0.6} = \frac{Pr[A]Pr[B|A]}{Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}]}$$
$$\approx 0.46$$

Bayes and Biased Coin



Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6; Pr[A \cap B] = 0.5 \times 0.5$$


$$Pr[B] = 0.5 \times 0.5 + 0.5 \times 0.6 = Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}]$$

$$Pr[A|B] = \frac{0.5 \times 0.5}{0.5 \times 0.5 + 0.5 \times 0.6} = \frac{Pr[A]Pr[B|A]}{Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}]}$$


≈ 0.46 = fraction of B that is inside A

Bayes: General Case

Bayes: General Case



Event B





$$p_n = Pr[A_n]$$

$$q_n = Pr[B|A_n]$$


A_1, \dots, A_N disjoint

$$A_1 \cup \dots \cup A_N = \Omega$$


Bayes: General Case



Bayes: General Case




Event B




Pick a point uniformly at random in the unit square. Then

Bayes: General Case




Event B




Pick a point uniformly at random in the unit square. Then

$$Pr[A_n] = p_n, n = 1, \dots, N$$

Bayes: General Case



Event B




Pick a point uniformly at random in the unit square. Then


$$Pr[A_n] = p_n, n = 1, \dots, N$$

$$Pr[B|A_n] = q_n, n = 1, \dots, N;$$

Bayes: General Case



Event B




Pick a point uniformly at random in the unit square. Then


$$Pr[A_n] = p_n, n = 1, \dots, N$$

$$Pr[B|A_n] = q_n, n = 1, \dots, N; Pr[A_n \cap B] =$$

Bayes: General Case



Event B




Pick a point uniformly at random in the unit square. Then

$$Pr[A_n] = p_n, n = 1, \dots, N$$

$$Pr[B|A_n] = q_n, n = 1, \dots, N; Pr[A_n \cap B] = p_n q_n$$

Bayes: General Case




Pick a point uniformly at random in the unit square. Then

$$Pr[A_n] = p_n, n = 1, \dots, N$$

$$Pr[B|A_n] = q_n, n = 1, \dots, N; Pr[A_n \cap B] = p_n q_n$$

$$Pr[B] = p_1 q_1 + \dots + p_N q_N$$

Bayes: General Case



Pick a point uniformly at random in the unit square. Then


$$\Pr[A_n] = p_n, n = 1, \dots, N$$

$$\Pr[B|A_n] = q_n, n = 1, \dots, N; \Pr[A_n \cap B] = p_n q_n$$

$$\Pr[B] = p_1 q_1 + \dots + p_N q_N$$

$$\Pr[A_n|B] = \frac{p_n q_n}{p_1 q_1 + \dots + p_N q_N}$$

Bayes: General Case



Pick a point uniformly at random in the unit square. Then

$$\Pr[A_n] = p_n, n = 1, \dots, N$$

$$\Pr[B|A_n] = q_n, n = 1, \dots, N; \Pr[A_n \cap B] = p_n q_n$$

$$\Pr[B] = p_1 q_1 + \dots + p_N q_N$$


$$\Pr[A_n|B] = \frac{p_n q_n}{p_1 q_1 + \dots + p_N q_N} = \text{fraction of } B \text{ inside } A_n.$$

Bayes Rule

A general picture: We imagine that there are N possible causes A_1, \dots, A_N .


Bayes Rule

A general picture: We imagine that there are N possible causes A_1, \dots, A_N .



Bayes Rule


A general picture: We imagine that there are N possible causes A_1, \dots, A_N .



100 situations: $100p_nq_n$ where A_n and B occur, for $n = 1, \dots, N$.

Bayes Rule


A general picture: We imagine that there are N possible causes A_1, \dots, A_N .



100 situations: $100p_nq_n$ where A_n and B occur, for $n = 1, \dots, N$.
In $100 \sum_m p_m q_m$ occurrences of B , $100p_nq_n$ occurrences of A_n .

Bayes Rule

A general picture: We imagine that there are N possible causes A_1, \dots, A_N .




100 situations: $100p_nq_n$ where A_n and B occur, for $n = 1, \dots, N$.
In $100\sum_m p_m q_m$ occurrences of B , $100p_nq_n$ occurrences of A_n .

Hence,

Bayes Rule

A general picture: We imagine that there are N possible causes A_1, \dots, A_N .




100 situations: $100p_nq_n$ where A_n and B occur, for $n = 1, \dots, N$.
In $100\sum_m p_mq_m$ occurrences of B , $100p_nq_n$ occurrences of A_n .

Hence,

$$\Pr[A_n|B] = \frac{p_nq_n}{\sum_m p_mq_m}.$$

Bayes Rule

A general picture: We imagine that there are N possible causes A_1, \dots, A_N .



100 situations: $100p_nq_n$ where A_n and B occur, for $n = 1, \dots, N$.
In $100\sum_m p_m q_m$ occurrences of B , $100p_nq_n$ occurrences of A_n .


Hence,

$$\Pr[A_n|B] = \frac{p_n q_n}{\sum_m p_m q_m}.$$

But, $p_n = \Pr[A_n]$, $q_n = \Pr[B|A_n]$, $\sum_m p_m q_m = \Pr[B]$, hence,

Bayes Rule

A general picture: We imagine that there are N possible causes A_1, \dots, A_N .



100 situations: $100p_nq_n$ where A_n and B occur, for $n = 1, \dots, N$.
 In $100\sum_m p_m q_m$ occurrences of B , $100p_nq_n$ occurrences of A_n .


Hence,

$$\Pr[A_n|B] = \frac{p_n q_n}{\sum_m p_m q_m}.$$


But, $p_n = \Pr[A_n]$, $q_n = \Pr[B|A_n]$, $\sum_m p_m q_m = \Pr[B]$, hence,

$$\Pr[A_n|B] = \frac{\Pr[B|A_n]\Pr[A_n]}{\Pr[B]}.$$


Why do you have a fever?



Why do you have a fever?




Why do you have a fever?



Using Bayes' rule, we find

$$Pr[\text{Flu}|\text{High Fever}] = \frac{0.15 \times 0.80}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.58$$

Why do you have a fever?




Using Bayes' rule, we find

$$Pr[\text{Flu}|\text{High Fever}] = \frac{0.15 \times 0.80}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.58$$

$$Pr[\text{Ebola}|\text{High Fever}] = \frac{10^{-8} \times 1}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 5 \times 10^{-8}$$

Why do you have a fever?




Using Bayes' rule, we find

$$Pr[\text{Flu}|\text{High Fever}] = \frac{0.15 \times 0.80}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.58$$

$$Pr[\text{Ebola}|\text{High Fever}] = \frac{10^{-8} \times 1}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 5 \times 10^{-8}$$

$$Pr[\text{Other}|\text{High Fever}] = \frac{0.85 \times 0.1}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.42$$

Why do you have a fever?



Using Bayes' rule, we find

$$Pr[\text{Flu}|\text{High Fever}] = \frac{0.15 \times 0.80}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.58$$

$$Pr[\text{Ebola}|\text{High Fever}] = \frac{10^{-8} \times 1}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 5 \times 10^{-8}$$

$$Pr[\text{Other}|\text{High Fever}] = \frac{0.85 \times 0.1}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.42$$

The values 0.58, 5×10^{-8} , 0.42 are the **posterior probabilities**.



Why do you have a fever?

Why do you have a fever?

Our “Bayes’ Square” picture:

Why do you have a fever?


Our “Bayes’ Square” picture:



58% of Fever = Flu
 $\approx 0\%$ of Fever = Ebola
42% of Fever = Other

Why do you have a fever?


Our “Bayes’ Square” picture:



Note that even though $Pr[\text{Fever}|\text{Ebola}] = 1$,

Why do you have a fever?

Our “Bayes’ Square” picture:




Note that even though $Pr[\text{Fever}|\text{Ebola}] = 1$, one has

$$Pr[\text{Ebola}|\text{Fever}] \approx 0.$$

Why do you have a fever?

Our “Bayes’ Square” picture:



Note that even though $Pr[\text{Fever}|\text{Ebola}] = 1$, one has

$$Pr[\text{Ebola}|\text{Fever}] \approx 0.$$

This example shows the importance of the prior probabilities.

Why do you have a fever?

We found

Why do you have a fever?

We found

$$Pr[\text{Flu}|\text{High Fever}] \approx 0.58,$$

$$Pr[\text{Ebola}|\text{High Fever}] \approx 5 \times 10^{-8},$$

$$Pr[\text{Other}|\text{High Fever}] \approx 0.42$$

Why do you have a fever?

We found

$$Pr[\text{Flu}|\text{High Fever}] \approx 0.58,$$

$$Pr[\text{Ebola}|\text{High Fever}] \approx 5 \times 10^{-8},$$

$$Pr[\text{Other}|\text{High Fever}] \approx 0.42$$

'Flu' is **Most Likely a Posteriori** (MAP) cause of high fever.

Why do you have a fever?

We found

$$Pr[\text{Flu}|\text{High Fever}] \approx 0.58,$$

$$Pr[\text{Ebola}|\text{High Fever}] \approx 5 \times 10^{-8},$$

$$Pr[\text{Other}|\text{High Fever}] \approx 0.42$$

'Flu' is **Most Likely a Posteriori** (MAP) cause of high fever.

'Ebola' is **Maximum Likelihood Estimate** (MLE) of cause:
causes fever with largest probability.

Why do you have a fever?

We found

$$Pr[\text{Flu}|\text{High Fever}] \approx 0.58,$$

$$Pr[\text{Ebola}|\text{High Fever}] \approx 5 \times 10^{-8},$$

$$Pr[\text{Other}|\text{High Fever}] \approx 0.42$$

'Flu' is **Most Likely a Posteriori** (MAP) cause of high fever.

'Ebola' is **Maximum Likelihood Estimate** (MLE) of cause:
causes fever with largest probability.

Recall that

$$p_m = Pr[A_m], q_m = Pr[B|A_m], Pr[A_m|B] = \frac{p_m q_m}{p_1 q_1 + \cdots + p_M q_M}.$$

Why do you have a fever?

We found

$$Pr[\text{Flu}|\text{High Fever}] \approx 0.58,$$

$$Pr[\text{Ebola}|\text{High Fever}] \approx 5 \times 10^{-8},$$

$$Pr[\text{Other}|\text{High Fever}] \approx 0.42$$

'Flu' is **Most Likely a Posteriori** (MAP) cause of high fever.

'Ebola' is **Maximum Likelihood Estimate** (MLE) of cause:
causes fever with largest probability.

Recall that

$$p_m = Pr[A_m], q_m = Pr[B|A_m], Pr[A_m|B] = \frac{p_m q_m}{p_1 q_1 + \dots + p_M q_M}.$$

Thus,

- ▶ MAP = value of m that maximizes $p_m q_m$.

Why do you have a fever?

We found

$$Pr[\text{Flu}|\text{High Fever}] \approx 0.58,$$

$$Pr[\text{Ebola}|\text{High Fever}] \approx 5 \times 10^{-8},$$

$$Pr[\text{Other}|\text{High Fever}] \approx 0.42$$

'Flu' is **Most Likely a Posteriori** (MAP) cause of high fever.

'Ebola' is **Maximum Likelihood Estimate** (MLE) of cause:
causes fever with largest probability.

Recall that


$$p_m = Pr[A_m], q_m = Pr[B|A_m], Pr[A_m|B] = \frac{p_m q_m}{p_1 q_1 + \dots + p_M q_M}.$$

Thus,


- ▶ MAP = value of m that maximizes $p_m q_m$.
- ▶ MLE = value of m that maximizes q_m .

Bayes' Rule Operations

Bayes' Rule Operations



Bayes' Rule Operations



Bayes' Rule: canonical example of how information changes our opinions.

Thomas Bayes

Thomas Bayes




Portrait used of Bayes in a 1936 book,^[1] but it is doubtful whether the portrait is actually of him.^[2]

No earlier portrait or claimed portrait survives.

Born	c. 1701 London, England
Died	7 April 1761 (aged 59) Tunbridge Wells, Kent , England
Residence	Tunbridge Wells, Kent, England
Nationality	English
Known for	Bayes' theorem

Thomas Bayes



A Bayesian picture of Thomas Bayes.

Testing for disease.

Random Experiment: Pick a random male.

Testing for disease.

Random Experiment: Pick a random male.

Outcomes: (*test*, *disease*)

Testing for disease.

Random Experiment: Pick a random male.

Outcomes: (*test*, *disease*)

A - prostate cancer.

B - positive PSA test.

Testing for disease.

Random Experiment: Pick a random male.

Outcomes: (*test, disease*)

A - prostate cancer.

B - positive PSA test.

- ▶ $Pr[A] = 0.0016$, (.16 % of the male population is affected.)
- ▶ $Pr[B|A] = 0.80$ (80% chance of positive test with disease.)
- ▶ $Pr[B|\bar{A}] = 0.10$ (10% chance of positive test without disease.)

Testing for disease.

Random Experiment: Pick a random male.

Outcomes: (*test, disease*)

A - prostate cancer.

B - positive PSA test.

- ▶ $Pr[A] = 0.0016$, (.16 % of the male population is affected.)
- ▶ $Pr[B|A] = 0.80$ (80% chance of positive test with disease.)
- ▶ $Pr[B|\bar{A}] = 0.10$ (10% chance of positive test without disease.)

From http://www.cpcn.org/01_psa_tests.htm and

<http://seer.cancer.gov/statfacts/html/prost.html> (10/12/2011.)

Testing for disease.

Random Experiment: Pick a random male.

Outcomes: (*test, disease*)

A - prostate cancer.

B - positive PSA test.

- ▶ $Pr[A] = 0.0016$, (.16 % of the male population is affected.)
- ▶ $Pr[B|A] = 0.80$ (80% chance of positive test with disease.)
- ▶ $Pr[B|\bar{A}] = 0.10$ (10% chance of positive test without disease.)

From http://www.cpcn.org/01_psa_tests.htm and
<http://seer.cancer.gov/statfacts/html/prost.html> (10/12/2011.)

Positive PSA test (B). Do I have disease?

Testing for disease.

Random Experiment: Pick a random male.

Outcomes: (*test, disease*)

A - prostate cancer.

B - positive PSA test.


- ▶ $Pr[A] = 0.0016$, (.16 % of the male population is affected.)
- ▶ $Pr[B|A] = 0.80$ (80% chance of positive test with disease.)
- ▶ $Pr[B|\bar{A}] = 0.10$ (10% chance of positive test without disease.)

From http://www.cpcn.org/01_psa_tests.htm and
<http://seer.cancer.gov/statfacts/html/prost.html> (10/12/2011.)


Positive PSA test (B). Do I have disease?

$$Pr[A|B]???$$

Bayes Rule.




Bayes Rule.



Using Bayes' rule, we find


Bayes Rule.



Using Bayes' rule, we find

$$P[A|B] = \frac{0.0016 \times 0.80}{0.0016 \times 0.80 + 0.9984 \times 0.10}$$


Bayes Rule.



Using Bayes' rule, we find

$$P[A|B] = \frac{0.0016 \times 0.80}{0.0016 \times 0.80 + 0.9984 \times 0.10} = .013.$$

Bayes Rule.




Using Bayes' rule, we find

$$P[A|B] = \frac{0.0016 \times 0.80}{0.0016 \times 0.80 + 0.9984 \times 0.10} = .013.$$

A 1.3% chance of prostate cancer with a positive PSA test.

Bayes Rule.




Using Bayes' rule, we find

$$P[A|B] = \frac{0.0016 \times 0.80}{0.0016 \times 0.80 + 0.9984 \times 0.10} = .013.$$

A 1.3% chance of prostate cancer with a positive PSA test.
Surgery anyone?

Bayes Rule.



Using Bayes' rule, we find


$$P[A|B] = \frac{0.0016 \times 0.80}{0.0016 \times 0.80 + 0.9984 \times 0.10} = .013.$$

A 1.3% chance of prostate cancer with a positive PSA test.

Surgery anyone?

Impotence...

Bayes Rule.



Using Bayes' rule, we find

$$P[A|B] = \frac{0.0016 \times 0.80}{0.0016 \times 0.80 + 0.9984 \times 0.10} = .013.$$


A 1.3% chance of prostate cancer with a positive PSA test.

Surgery anyone?

Impotence...

Incontinence..

Bayes Rule.



Using Bayes' rule, we find

$$P[A|B] = \frac{0.0016 \times 0.80}{0.0016 \times 0.80 + 0.9984 \times 0.10} = .013.$$

A 1.3% chance of prostate cancer with a positive PSA test.

Surgery anyone?

Impotence...

Incontinence..

Death.

Quick Review

Events, Conditional Probability, Independence, Bayes' Rule

Quick Review

Events, Conditional Probability, Independence, Bayes' Rule

Key Ideas:

- ▶ Conditional Probability:

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$$

Quick Review

Events, Conditional Probability, Independence, Bayes' Rule

Key Ideas:

- ▶ Conditional Probability:

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$$

- ▶ Independence: $Pr[A \cap B] = Pr[A]Pr[B]$.

Quick Review

Events, Conditional Probability, Independence, Bayes' Rule

Key Ideas:

- ▶ Conditional Probability:

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$$

- ▶ Independence: $Pr[A \cap B] = Pr[A]Pr[B]$.

- ▶ Bayes' Rule:

$$Pr[A_n|B] = \frac{Pr[A_n]Pr[B|A_n]}{\sum_m Pr[A_m]Pr[B|A_m]}.$$

Quick Review

Events, Conditional Probability, Independence, Bayes' Rule

Key Ideas:

- ▶ Conditional Probability:

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$$

- ▶ Independence: $Pr[A \cap B] = Pr[A]Pr[B]$.

- ▶ Bayes' Rule:

$$Pr[A_n|B] = \frac{Pr[A_n]Pr[B|A_n]}{\sum_m Pr[A_m]Pr[B|A_m]}.$$

$Pr[A_n|B]$ = posterior probability; $Pr[A_n]$ = prior probability .

Quick Review

Events, Conditional Probability, Independence, Bayes' Rule

Key Ideas:

- ▶ Conditional Probability:

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$$

- ▶ Independence: $Pr[A \cap B] = Pr[A]Pr[B]$.

- ▶ Bayes' Rule:

$$Pr[A_n|B] = \frac{Pr[A_n]Pr[B|A_n]}{\sum_m Pr[A_m]Pr[B|A_m]}.$$

$Pr[A_n|B]$ = posterior probability; $Pr[A_n]$ = prior probability .

- ▶ All these are possible:

$$Pr[A|B] < Pr[A]; Pr[A|B] > Pr[A]; Pr[A|B] = Pr[A].$$

Independence

Recall :

A and B are independent

Independence

Recall :

A and B are independent

$$\Leftrightarrow \Pr[A \cap B] = \Pr[A] \Pr[B]$$

Independence

Recall :

A and B are independent

$$\Leftrightarrow \Pr[A \cap B] = \Pr[A] \Pr[B]$$

$$\Leftrightarrow \Pr[A|B] = \Pr[A].$$

Independence


Recall :

A and B are independent

$$\Leftrightarrow \Pr[A \cap B] = \Pr[A] \Pr[B]$$

$$\Leftrightarrow \Pr[A|B] = \Pr[A].$$

Consider the example below:



Independence


Recall :

A and B are independent

$$\Leftrightarrow Pr[A \cap B] = Pr[A]Pr[B]$$

$$\Leftrightarrow Pr[A|B] = Pr[A].$$

Consider the example below:



Which are independent? (A) (A_2, B) (B) (A_2, \bar{B}) (C) (A_1, B) .

Independence


Recall :

A and B are independent

$$\Leftrightarrow \Pr[A \cap B] = \Pr[A]\Pr[B]$$

$$\Leftrightarrow \Pr[A|B] = \Pr[A].$$

Consider the example below:



Which are independent? (A) (A_2, B) (B) (A_2, \bar{B}) (C) (A_1, B) .
 (A_2, B) are independent:

Independence


Recall :

A and B are independent

$$\Leftrightarrow Pr[A \cap B] = Pr[A]Pr[B]$$

$$\Leftrightarrow Pr[A|B] = Pr[A].$$

Consider the example below:



Which are independent? (A) (A_2, B) (B) (A_2, \bar{B}) (C) (A_1, B) .
 (A_2, B) are independent: $Pr[A_2|B] = 0.5 = Pr[A_2]$.

Independence


Recall :

A and B are independent

$$\Leftrightarrow Pr[A \cap B] = Pr[A]Pr[B]$$

$$\Leftrightarrow Pr[A|B] = Pr[A].$$

Consider the example below:



Which are independent? (A) (A_2, B) (B) (A_2, \bar{B}) (C) (A_1, B) .
 (A_2, B) are independent: $Pr[A_2|B] = 0.5 = Pr[A_2]$.
 (A_2, \bar{B}) are independent:

Independence


Recall :

A and B are independent

$$\Leftrightarrow Pr[A \cap B] = Pr[A]Pr[B]$$

$$\Leftrightarrow Pr[A|B] = Pr[A].$$

Consider the example below:



Which are independent? (A) (A_2, B) (B) (A_2, \bar{B}) (C) (A_1, B) .

(A_2, B) are independent: $Pr[A_2|B] = 0.5 = Pr[A_2]$.

(A_2, \bar{B}) are independent: $Pr[A_2|\bar{B}] = 0.5 = Pr[A_2]$.

Independence


Recall :

A and B are independent

$$\Leftrightarrow Pr[A \cap B] = Pr[A]Pr[B]$$

$$\Leftrightarrow Pr[A|B] = Pr[A].$$

Consider the example below:



Which are independent? (A) (A_2, B) (B) (A_2, \bar{B}) (C) (A_1, B) .

(A_2, B) are independent: $Pr[A_2|B] = 0.5 = Pr[A_2]$.

(A_2, \bar{B}) are independent: $Pr[A_2|\bar{B}] = 0.5 = Pr[A_2]$.

(A_1, B) are not independent:

Independence


Recall :

A and B are independent

$$\Leftrightarrow Pr[A \cap B] = Pr[A]Pr[B]$$

$$\Leftrightarrow Pr[A|B] = Pr[A].$$

Consider the example below:



Which are independent? (A) (A_2, B) (B) (A_2, \bar{B}) (C) (A_1, B) .

(A_2, B) are independent: $Pr[A_2|B] = 0.5 = Pr[A_2]$.

(A_2, \bar{B}) are independent: $Pr[A_2|\bar{B}] = 0.5 = Pr[A_2]$.

(A_1, B) are not independent: $Pr[A_1|B] = \frac{0.1}{0.5} = 0.2 \neq Pr[A_1] = 0.25$.

Pairwise Independence


Flip two fair coins. Let

- ▶ $A = \text{'first coin is H'} = \{HT, HH\};$
- ▶ $B = \text{'second coin is H'} = \{TH, HH\};$
- ▶ $C = \text{'the two coins are different'} = \{TH, HT\}.$

Pairwise Independence

Flip two fair coins. Let


- ▶ $A = \text{'first coin is H'} = \{HT, HH\}$;
- ▶ $B = \text{'second coin is H'} = \{TH, HH\}$;
- ▶ $C = \text{'the two coins are different'} = \{TH, HT\}$.



Pairwise Independence

Flip two fair coins. Let

- ▶ $A = \text{'first coin is H'} = \{HT, HH\}$;
- ▶ $B = \text{'second coin is H'} = \{TH, HH\}$;
- ▶ $C = \text{'the two coins are different'} = \{TH, HT\}$.




A, C are independent;

Pairwise Independence

Flip two fair coins. Let

- ▶ $A = \text{'first coin is H'} = \{HT, HH\}$;
- ▶ $B = \text{'second coin is H'} = \{TH, HH\}$;
- ▶ $C = \text{'the two coins are different'} = \{TH, HT\}$.




A, C are independent; B, C are independent;

Pairwise Independence

Flip two fair coins. Let

- ▶ $A = \text{'first coin is H'} = \{HT, HH\}$;
- ▶ $B = \text{'second coin is H'} = \{TH, HH\}$;
- ▶ $C = \text{'the two coins are different'} = \{TH, HT\}$.




A, C are independent; B, C are independent;

$A \cap B, C$ are **not** independent.

Pairwise Independence

Flip two fair coins. Let

- ▶ $A = \text{'first coin is H'} = \{HT, HH\}$;
- ▶ $B = \text{'second coin is H'} = \{TH, HH\}$;
- ▶ $C = \text{'the two coins are different'} = \{TH, HT\}$.




A, C are independent; B, C are independent;

$A \cap B, C$ are **not** independent. ($Pr[A \cap B \cap C] = 0 \neq Pr[A \cap B]Pr[C]$.)

Pairwise Independence

Flip two fair coins. Let

- ▶ $A = \text{'first coin is H'} = \{HT, HH\}$;
- ▶ $B = \text{'second coin is H'} = \{TH, HH\}$;
- ▶ $C = \text{'the two coins are different'} = \{TH, HT\}$.



A, C are independent; B, C are independent;

$A \cap B, C$ are **not** independent. ($Pr[A \cap B \cap C] = 0 \neq Pr[A \cap B]Pr[C]$.)

False: If A did not say anything about C and B did not say anything about C , then $A \cap B$ would not say anything about C .

Example 2

Flip a fair coin 5 times.

Example 2

Flip a fair coin 5 times. Let A_n = 'coin n is H', for $n = 1, \dots, 5$.

Example 2

Flip a fair coin 5 times. Let A_n = 'coin n is H', for $n = 1, \dots, 5$.

Then,

A_m, A_n are independent for all $m \neq n$.

Example 2

Flip a fair coin 5 times. Let A_n = 'coin n is H', for $n = 1, \dots, 5$.

Then,

A_m, A_n are independent for all $m \neq n$.

Also,

A_1 and $A_3 \cap A_5$ are independent.

Example 2

Flip a fair coin 5 times. Let A_n = 'coin n is H', for $n = 1, \dots, 5$.

Then,

A_m, A_n are independent for all $m \neq n$.

Also,

A_1 and $A_3 \cap A_5$ are independent.

Indeed,

$$Pr[A_1 \cap (A_3 \cap A_5)] = \frac{1}{8} = Pr[A_1]Pr[A_3 \cap A_5].$$

Example 2

Flip a fair coin 5 times. Let A_n = 'coin n is H', for $n = 1, \dots, 5$.

Then,

A_m, A_n are independent for all $m \neq n$.

Also,

A_1 and $A_3 \cap A_5$ are independent.

Indeed,

$$Pr[A_1 \cap (A_3 \cap A_5)] = \frac{1}{8} = Pr[A_1]Pr[A_3 \cap A_5].$$

Similarly,

$A_1 \cap A_2$ and $A_3 \cap A_4 \cap A_5$ are independent.

Example 2

Flip a fair coin 5 times. Let A_n = 'coin n is H', for $n = 1, \dots, 5$.

Then,

A_m, A_n are independent for all $m \neq n$.

Also,

A_1 and $A_3 \cap A_5$ are independent.

Indeed,

$$Pr[A_1 \cap (A_3 \cap A_5)] = \frac{1}{8} = Pr[A_1]Pr[A_3 \cap A_5].$$

Similarly,

$A_1 \cap A_2$ and $A_3 \cap A_4 \cap A_5$ are independent.

This leads to a definition

Mutual Independence

Definition Mutual Independence

Mutual Independence

Definition Mutual Independence

(a) The events A_1, \dots, A_5 are mutually independent if

Mutual Independence

Definition Mutual Independence

(a) The events A_1, \dots, A_5 are **mutually independent** if

$$Pr[\cap_{k \in K} A_k] = \prod_{k \in K} Pr[A_k], \text{ for all } K \subseteq \{1, \dots, 5\}.$$

Mutual Independence

Definition Mutual Independence

(a) The events A_1, \dots, A_5 are **mutually independent** if

$$Pr[\cap_{k \in K} A_k] = \prod_{k \in K} Pr[A_k], \text{ for all } K \subseteq \{1, \dots, 5\}.$$

(b) More generally, the events $\{A_j, j \in J\}$ are **mutually independent** if

Mutual Independence

Definition Mutual Independence

(a) The events A_1, \dots, A_5 are **mutually independent** if

$$Pr[\cap_{k \in K} A_k] = \prod_{k \in K} Pr[A_k], \text{ for all } K \subseteq \{1, \dots, 5\}.$$

(b) More generally, the events $\{A_j, j \in J\}$ are **mutually independent** if

$$Pr[\cap_{k \in K} A_k] = \prod_{k \in K} Pr[A_k], \text{ for all finite } K \subseteq J.$$

Mutual Independence

Definition Mutual Independence

(a) The events A_1, \dots, A_5 are **mutually independent** if

$$Pr[\cap_{k \in K} A_k] = \prod_{k \in K} Pr[A_k], \text{ for all } K \subseteq \{1, \dots, 5\}.$$

(b) More generally, the events $\{A_j, j \in J\}$ are **mutually independent** if

$$Pr[\cap_{k \in K} A_k] = \prod_{k \in K} Pr[A_k], \text{ for all finite } K \subseteq J.$$

Example: Flip a fair coin forever. Let $A_n =$ 'coin n is H.' Then the events A_n are mutually independent.

Mutual Independence

Theorem

Mutual Independence

Theorem

(a) If the events $\{A_j, j \in J\}$ are mutually independent and if K_1 and K_2 are disjoint finite subsets of J , then

Mutual Independence

Theorem

(a) If the events $\{A_j, j \in J\}$ are mutually independent and if K_1 and K_2 are disjoint finite subsets of J , then

$\cap_{k \in K_1} A_k$ and $\cap_{k \in K_2} A_k$ are independent.

Mutual Independence

Theorem

(a) If the events $\{A_j, j \in J\}$ are mutually independent and if K_1 and K_2 are disjoint finite subsets of J , then

$\cap_{k \in K_1} A_k$ and $\cap_{k \in K_2} A_k$ are independent.

(b) More generally, if the K_n are pairwise disjoint finite subsets of J , then the events

$\cap_{k \in K_n} A_k$ are mutually independent.

Mutual Independence

Theorem

(a) If the events $\{A_j, j \in J\}$ are mutually independent and if K_1 and K_2 are disjoint finite subsets of J , then

$\cap_{k \in K_1} A_k$ and $\cap_{k \in K_2} A_k$ are independent.

(b) More generally, if the K_n are pairwise disjoint finite subsets of J , then the events

$\cap_{k \in K_n} A_k$ are mutually independent.

(c) Also, the same is true if we replace some of the A_k by \bar{A}_k .

Mutual Independence

Theorem

(a) If the events $\{A_j, j \in J\}$ are mutually independent and if K_1 and K_2 are disjoint finite subsets of J , then

$\cap_{k \in K_1} A_k$ and $\cap_{k \in K_2} A_k$ are independent.

(b) More generally, if the K_n are pairwise disjoint finite subsets of J , then the events

$\cap_{k \in K_n} A_k$ are mutually independent.

(c) Also, the same is true if we replace some of the A_k by \bar{A}_k .

Conditional Probability: Review

Conditional Probability: Review

Recall:

$$\blacktriangleright \Pr[A|B] = \frac{\Pr[A \cap B]}{\Pr[B]}.$$

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A],$

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,
i.e., if $Pr[A \cap B] < Pr[A]Pr[B].$

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,
i.e., if $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶ A and B are *independent* if $Pr[A|B] = Pr[A],$

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,
i.e., if $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶ A and B are *independent* if $Pr[A|B] = Pr[A]$,
i.e., if $Pr[A \cap B] = Pr[A]Pr[B].$

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,
i.e., if $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶ A and B are *independent* if $Pr[A|B] = Pr[A]$,
i.e., if $Pr[A \cap B] = Pr[A]Pr[B].$
- ▶ Note: $B \subset A \Rightarrow A$ and B are

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,
i.e., if $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶ A and B are *independent* if $Pr[A|B] = Pr[A]$,
i.e., if $Pr[A \cap B] = Pr[A]Pr[B].$
- ▶ Note: $B \subset A \Rightarrow A$ and B are positively correlated.

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,
i.e., if $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶ A and B are *independent* if $Pr[A|B] = Pr[A]$,
i.e., if $Pr[A \cap B] = Pr[A]Pr[B].$
- ▶ Note: $B \subset A \Rightarrow A$ and B are positively correlated.
($Pr[A|B] = 1 > Pr[A]$)

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,
i.e., if $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶ A and B are *independent* if $Pr[A|B] = Pr[A]$,
i.e., if $Pr[A \cap B] = Pr[A]Pr[B].$
- ▶ Note: $B \subset A \Rightarrow A$ and B are positively correlated.
($Pr[A|B] = 1 > Pr[A]$)
- ▶ Note: $A \cap B = \emptyset \Rightarrow A$ and B are

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,
i.e., if $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶ A and B are *independent* if $Pr[A|B] = Pr[A]$,
i.e., if $Pr[A \cap B] = Pr[A]Pr[B].$
- ▶ Note: $B \subset A \Rightarrow A$ and B are positively correlated.
($Pr[A|B] = 1 > Pr[A]$)
- ▶ Note: $A \cap B = \emptyset \Rightarrow A$ and B are negatively correlated.

Conditional Probability: Review

Recall:

- ▶ $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence, $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶ A and B are *positively correlated* if $Pr[A|B] > Pr[A]$,
i.e., if $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶ A and B are *negatively correlated* if $Pr[A|B] < Pr[A]$,
i.e., if $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶ A and B are *independent* if $Pr[A|B] = Pr[A]$,
i.e., if $Pr[A \cap B] = Pr[A]Pr[B].$
- ▶ Note: $B \subset A \Rightarrow A$ and B are positively correlated.
($Pr[A|B] = 1 > Pr[A]$)
- ▶ Note: $A \cap B = \emptyset \Rightarrow A$ and B are negatively correlated.
($Pr[A|B] = 0 < Pr[A]$)

Quick Review.

Bayes' Rule, Mutual Independence, Collisions and Collecting

Quick Review.

Bayes' Rule, Mutual Independence, Collisions and Collecting

Main results:

- **Bayes' Rule:** $Pr[A_m|B] = p_m q_m / (p_1 q_1 + \cdots + p_M q_M)$.

Quick Review.

Bayes' Rule, Mutual Independence, Collisions and Collecting

Main results:

- ▶ **Bayes' Rule:** $Pr[A_m|B] = p_m q_m / (p_1 q_1 + \cdots + p_M q_M)$.
- ▶ **Product Rule:**
 $Pr[A_1 \cap \cdots \cap A_n] = Pr[A_1] Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \cdots \cap A_{n-1}]$.

Quick Review.

Bayes' Rule, Mutual Independence, Collisions and Collecting

Main results:

- ▶ **Bayes' Rule:** $Pr[A_m|B] = p_m q_m / (p_1 q_1 + \cdots + p_M q_M)$.
- ▶ **Product Rule:**
 $Pr[A_1 \cap \cdots \cap A_n] = Pr[A_1] Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \cdots \cap A_{n-1}]$.

Quick Review.

Bayes' Rule, Mutual Independence, Collisions and Collecting

Main results:

- ▶ **Bayes' Rule:** $Pr[A_m|B] = p_m q_m / (p_1 q_1 + \cdots + p_M q_M)$.
- ▶ **Product Rule:**
 $Pr[A_1 \cap \cdots \cap A_n] = Pr[A_1] Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \cdots \cap A_{n-1}]$.