

Coupon Collecting: Fun with harmonic numbers!

Coupon Collecting: Fun with harmonic numbers!
Memoryless Property.

Coupon Collecting: Fun with harmonic numbers!

Memoryless Property.

Law of the unconscious statistician. (Hmmm.)

Coupon Collecting: Fun with harmonic numbers!

Memoryless Property.

Law of the unconscious statistician. (Hmmm.)

Variance/ Covariance.

Time to collect coupons

X -time to get n coupons.

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon.

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$.

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

$Pr[\text{"get second coupon"} | \text{"got milk"}]$

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

$Pr[\text{"get second coupon"} | \text{"got milk first coupon"}] = \frac{n-1}{n}$

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

$Pr[\text{"get second coupon"} | \text{"got milk first coupon"}] = \frac{n-1}{n}$

$E[X_2]$?

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

$Pr[\text{"get second coupon"} | \text{"got milk first coupon"}] = \frac{n-1}{n}$

$E[X_2]$? Geometric

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

$Pr[\text{"get second coupon"} | \text{"got milk first coupon"}] = \frac{n-1}{n}$

$E[X_2]$? Geometric !

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

$Pr[\text{"get second coupon"} | \text{"got milk first coupon"}] = \frac{n-1}{n}$

$E[X_2]$? **Geometric ! !**

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

$Pr[\text{"get second coupon"} | \text{"got milk first coupon"}] = \frac{n-1}{n}$

$E[X_2]$? **Geometric ! ! !**

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

$Pr[\text{"get second coupon"} | \text{"got milk first coupon"}] = \frac{n-1}{n}$

$E[X_2]$? Geometric !!! $\implies E[X_2] = \frac{1}{p} =$

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

$Pr[\text{"get second coupon"} | \text{"got milk first coupon"}] = \frac{n-1}{n}$

$E[X_2]$? Geometric !!! $\implies E[X_2] = \frac{1}{p} = \frac{1}{\frac{n-1}{n}}$

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

$Pr[\text{"get second coupon"} | \text{"got milk first coupon"}] = \frac{n-1}{n}$

$E[X_2]$? Geometric !!! $\implies E[X_2] = \frac{1}{p} = \frac{1}{\frac{n-1}{n}} = \frac{n}{n-1}$.

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

$$Pr[\text{"get second coupon"} | \text{"got ~~red~~ first coupon"}] = \frac{n-1}{n}$$

$$E[X_2]? \text{ Geometric !!! } \implies E[X_2] = \frac{1}{p} = \frac{1}{\frac{n-1}{n}} = \frac{n}{n-1}.$$

$$Pr[\text{"getting } i\text{th coupon"} | \text{"got } i-1 \text{st coupons"}] = \frac{n-(i-1)}{n} = \frac{n-i+1}{n}$$

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

$$Pr[\text{"get second coupon"} | \text{"got milk first coupon"}] = \frac{n-1}{n}$$

$$E[X_2]? \text{ Geometric !!! } \implies E[X_2] = \frac{1}{p} = \frac{1}{\frac{n-1}{n}} = \frac{n}{n-1}.$$

$$Pr[\text{"getting } i\text{th coupon"} | \text{"got } i-1 \text{st coupons"}] = \frac{n-(i-1)}{n} = \frac{n-i+1}{n}$$

$$E[X_i]$$

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

$$\Pr[\text{"get second coupon"} | \text{"got ~~milk~~ first coupon"}] = \frac{n-1}{n}$$

$$E[X_2]? \text{ Geometric !!! } \implies E[X_2] = \frac{1}{p} = \frac{1}{\frac{n-1}{n}} = \frac{n}{n-1}.$$

$$\Pr[\text{"getting } i\text{th coupon"} | \text{"got } i-1 \text{st coupons"}] = \frac{n-(i-1)}{n} = \frac{n-i+1}{n}$$

$$E[X_i] = \frac{1}{p}$$

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

$$Pr[\text{"get second coupon"} | \text{"got ~~milk~~ first coupon"}] = \frac{n-1}{n}$$

$$E[X_2]? \text{ Geometric !!! } \implies E[X_2] = \frac{1}{p} = \frac{1}{\frac{n-1}{n}} = \frac{n}{n-1}.$$

$$Pr[\text{"getting } i\text{th coupon"} | \text{"got } i-1 \text{st coupons"}] = \frac{n-(i-1)}{n} = \frac{n-i+1}{n}$$

$$E[X_i] = \frac{1}{p} = \frac{n}{n-i+1},$$

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

$$\Pr[\text{"get second coupon"} | \text{"got milk first coupon"}] = \frac{n-1}{n}$$

$$E[X_2]? \text{ Geometric !!! } \implies E[X_2] = \frac{1}{p} = \frac{1}{\frac{n-1}{n}} = \frac{n}{n-1}.$$

$$\Pr[\text{"getting } i\text{th coupon"} | \text{"got } i-1 \text{st coupons"}] = \frac{n-(i-1)}{n} = \frac{n-i+1}{n}$$

$$E[X_i] = \frac{1}{p} = \frac{n}{n-i+1}, i = 1, 2, \dots, n.$$

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

$$Pr[\text{"get second coupon"} | \text{"got milk first coupon"}] = \frac{n-1}{n}$$

$$E[X_2]? \text{ Geometric !!! } \implies E[X_2] = \frac{1}{p} = \frac{1}{\frac{n-1}{n}} = \frac{n}{n-1}.$$

$$Pr[\text{"getting } i\text{th coupon"} | \text{"got } i-1 \text{st coupons"}] = \frac{n-(i-1)}{n} = \frac{n-i+1}{n}$$

$$E[X_i] = \frac{1}{p} = \frac{n}{n-i+1}, i = 1, 2, \dots, n.$$

$$E[X] = E[X_1] + \dots + E[X_n] =$$

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

$$\Pr[\text{"get second coupon"} | \text{"got milk first coupon"}] = \frac{n-1}{n}$$

$$E[X_2]? \text{ Geometric !!! } \implies E[X_2] = \frac{1}{p} = \frac{1}{\frac{n-1}{n}} = \frac{n}{n-1}.$$

$$\Pr[\text{"getting } i\text{th coupon"} | \text{"got } i-1 \text{st coupons"}] = \frac{n-(i-1)}{n} = \frac{n-i+1}{n}$$

$$E[X_i] = \frac{1}{p} = \frac{n}{n-i+1}, i = 1, 2, \dots, n.$$

$$E[X] = E[X_1] + \dots + E[X_n] = \frac{n}{n} + \frac{n}{n-1} + \frac{n}{n-2} + \dots + \frac{n}{1}$$

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

$$\Pr[\text{"get second coupon"} | \text{"got milk first coupon"}] = \frac{n-1}{n}$$

$$E[X_2]? \text{ Geometric !!! } \implies E[X_2] = \frac{1}{p} = \frac{1}{\frac{n-1}{n}} = \frac{n}{n-1}.$$

$$\Pr[\text{"getting } i\text{th coupon"} | \text{"got } i-1 \text{st coupons"}] = \frac{n-(i-1)}{n} = \frac{n-i+1}{n}$$

$$E[X_i] = \frac{1}{p} = \frac{n}{n-i+1}, i = 1, 2, \dots, n.$$

$$\begin{aligned} E[X] &= E[X_1] + \dots + E[X_n] = \frac{n}{n} + \frac{n}{n-1} + \frac{n}{n-2} + \dots + \frac{n}{1} \\ &= n\left(1 + \frac{1}{2} + \dots + \frac{1}{n}\right) =: nH(n) \end{aligned}$$

Time to collect coupons

X -time to get n coupons.

X_1 - time to get first coupon. Note: $X_1 = 1$. $E(X_1) = 1$.

X_2 - time to get second coupon after getting first.

$$\Pr[\text{"get second coupon"} | \text{"got milk first coupon"}] = \frac{n-1}{n}$$

$$E[X_2]? \text{ Geometric !!! } \implies E[X_2] = \frac{1}{p} = \frac{1}{\frac{n-1}{n}} = \frac{n}{n-1}.$$

$$\Pr[\text{"getting } i\text{th coupon"} | \text{"got } i-1 \text{st coupons"}] = \frac{n-(i-1)}{n} = \frac{n-i+1}{n}$$

$$E[X_i] = \frac{1}{p} = \frac{n}{n-i+1}, i = 1, 2, \dots, n.$$

$$\begin{aligned} E[X] &= E[X_1] + \dots + E[X_n] = \frac{n}{n} + \frac{n}{n-1} + \frac{n}{n-2} + \dots + \frac{n}{1} \\ &= n\left(1 + \frac{1}{2} + \dots + \frac{1}{n}\right) =: nH(n) \approx n(\ln n + \gamma) \end{aligned}$$


Review: Harmonic sum

$$H(n) = 1 + \frac{1}{2} + \cdots + \frac{1}{n} \approx \int_1^n \frac{1}{x} dx = \ln(n).$$

.


Review: Harmonic sum

$$H(n) = 1 + \frac{1}{2} + \cdots + \frac{1}{n} \approx \int_1^n \frac{1}{x} dx = \ln(n).$$



Review: Harmonic sum

$$H(n) = 1 + \frac{1}{2} + \cdots + \frac{1}{n} \approx \int_1^n \frac{1}{x} dx = \ln(n).$$



A good approximation is


$$H(n) \approx \ln(n) + \gamma \text{ where } \gamma \approx 0.58 \text{ (Euler-Mascheroni constant).}$$

Harmonic sum: Paradox

Consider this stack of cards (no glue!):


Harmonic sum: Paradox

Consider this stack of cards (no glue!):



Harmonic sum: Paradox


Consider this stack of cards (no glue!):



If each card has length 2, the stack can extend $H(n)$ to the right of the table.

Harmonic sum: Paradox

Consider this stack of cards (no glue!):



If each card has length 2, the stack can extend $H(n)$ to the right of the table. As n increases, you can go as far as you want!

Paradox

par·a·dox

/ˈperəˌdäks/


noun

a statement or proposition that, despite sound (or apparently sound) reasoning from acceptable premises, leads to a conclusion that seems senseless, logically unacceptable, or self-contradictory.


"a potentially serious conflict between quantum mechanics and the general theory of relativity known as the information paradox"

- a seemingly absurd or self-contradictory statement or proposition that when investigated or explained may prove to be well founded or true.
"in a paradox, he has discovered that stepping back from his job has increased the rewards he gleans from it"
synonyms: [contradiction](#), contradiction in terms, [self-contradiction](#), [inconsistency](#), [incongruity](#); [More](#)
- a situation, person, or thing that combines contradictory features or qualities.
"the mingling of deciduous trees with elements of desert flora forms a fascinating ecological paradox"

Stacking




Stacking



The cards have width 2.

Stacking



The cards have width 2. Induction shows that the center of gravity after n cards is $H(n)$ away from the right-most edge.

[Video.](#)

Calculating $E[g(X)]$: LOTUS

Calculating $E[g(X)]$: LOTUS

Let $Y = g(X)$.

Calculating $E[g(X)]$: LOTUS

Let $Y = g(X)$. Assume that we know the distribution of X .

Calculating $E[g(X)]$: LOTUS

Let $Y = g(X)$. Assume that we know the distribution of X .

We want to calculate $E[Y]$.

Calculating $E[g(X)]$: LOTUS

Let $Y = g(X)$. Assume that we know the distribution of X .

We want to calculate $E[Y]$.

Method 1:

Calculating $E[g(X)]$: LOTUS

Let $Y = g(X)$. Assume that we know the distribution of X .

We want to calculate $E[Y]$.

Method 1: We calculate the distribution of Y :

Calculating $E[g(X)]$: LOTUS

Let $Y = g(X)$. Assume that we know the distribution of X .

We want to calculate $E[Y]$.

Method 1: We calculate the distribution of Y :

$$Pr[Y = y] = Pr[X \in g^{-1}(y)] \text{ where } g^{-1}(x) = \{x \in \mathfrak{X} : g(x) = y\}.$$

Calculating $E[g(X)]$: LOTUS

Let $Y = g(X)$. Assume that we know the distribution of X .

We want to calculate $E[Y]$.

Method 1: We calculate the distribution of Y :

$$Pr[Y = y] = Pr[X \in g^{-1}(y)] \text{ where } g^{-1}(x) = \{x \in \mathfrak{X} : g(x) = y\}.$$

This is typically rather tedious!

Calculating $E[g(X)]$: LOTUS

Let $Y = g(X)$. Assume that we know the distribution of X .

We want to calculate $E[Y]$.

Method 1: We calculate the distribution of Y :

$$Pr[Y = y] = Pr[X \in g^{-1}(y)] \text{ where } g^{-1}(x) = \{x \in \mathfrak{X} : g(x) = y\}.$$

This is typically rather tedious!

Method 2: We use the following result.

Calculating $E[g(X)]$: LOTUS

Let $Y = g(X)$. Assume that we know the distribution of X .

We want to calculate $E[Y]$.

Method 1: We calculate the distribution of Y :

$$Pr[Y = y] = Pr[X \in g^{-1}(y)] \text{ where } g^{-1}(x) = \{x \in \mathfrak{X} : g(x) = y\}.$$

This is typically rather tedious!

Method 2: We use the following result.

Called “Law of the unconscious statistician.”

Theorem:

$$E[g(X)] = \sum_x g(x) Pr[X = x].$$

Calculating $E[g(X)]$: LOTUS

Let $Y = g(X)$. Assume that we know the distribution of X .

We want to calculate $E[Y]$.

Method 1: We calculate the distribution of Y :

$$Pr[Y = y] = Pr[X \in g^{-1}(y)] \text{ where } g^{-1}(x) = \{x \in \mathfrak{X} : g(x) = y\}.$$

This is typically rather tedious!

Method 2: We use the following result.

Called “Law of the unconscious statistician.”

Theorem:

$$E[g(X)] = \sum_x g(x) Pr[X = x].$$

Proof:

Calculating $E[g(X)]$: LOTUS

Let $Y = g(X)$. Assume that we know the distribution of X .

We want to calculate $E[Y]$.

Method 1: We calculate the distribution of Y :

$$Pr[Y = y] = Pr[X \in g^{-1}(y)] \text{ where } g^{-1}(x) = \{x \in \mathfrak{X} : g(x) = y\}.$$

This is typically rather tedious!

Method 2: We use the following result.

Called “Law of the unconscious statistician.”

Theorem:

$$E[g(X)] = \sum_x g(x) Pr[X = x].$$

Proof:

$$E[g(X)] = \sum_{\omega} g(X(\omega)) Pr[\omega]$$

Calculating $E[g(X)]$: LOTUS

Let $Y = g(X)$. Assume that we know the distribution of X .

We want to calculate $E[Y]$.

Method 1: We calculate the distribution of Y :

$$Pr[Y = y] = Pr[X \in g^{-1}(y)] \text{ where } g^{-1}(x) = \{x \in \mathfrak{X} : g(x) = y\}.$$

This is typically rather tedious!

Method 2: We use the following result.

Called “Law of the unconscious statistician.”

Theorem:

$$E[g(X)] = \sum_x g(x) Pr[X = x].$$

Proof:

$$E[g(X)] = \sum_{\omega} g(X(\omega)) Pr[\omega] = \sum_x \sum_{\omega \in X^{-1}(x)} g(X(\omega)) Pr[\omega]$$

Calculating $E[g(X)]$: LOTUS

Let $Y = g(X)$. Assume that we know the distribution of X .

We want to calculate $E[Y]$.

Method 1: We calculate the distribution of Y :

$$Pr[Y = y] = Pr[X \in g^{-1}(y)] \text{ where } g^{-1}(x) = \{x \in \mathfrak{X} : g(x) = y\}.$$

This is typically rather tedious!

Method 2: We use the following result.

Called “Law of the unconscious statistician.”

Theorem:

$$E[g(X)] = \sum_x g(x) Pr[X = x].$$

Proof:

$$\begin{aligned} E[g(X)] &= \sum_{\omega} g(X(\omega)) Pr[\omega] = \sum_x \sum_{\omega \in X^{-1}(x)} g(X(\omega)) Pr[\omega] \\ &= \sum_x \sum_{\omega \in X^{-1}(x)} g(x) Pr[\omega] \end{aligned}$$

Calculating $E[g(X)]$: LOTUS

Let $Y = g(X)$. Assume that we know the distribution of X .

We want to calculate $E[Y]$.

Method 1: We calculate the distribution of Y :

$$Pr[Y = y] = Pr[X \in g^{-1}(y)] \text{ where } g^{-1}(x) = \{x \in \mathfrak{X} : g(x) = y\}.$$

This is typically rather tedious!

Method 2: We use the following result.

Called “Law of the unconscious statistician.”

Theorem:

$$E[g(X)] = \sum_x g(x) Pr[X = x].$$

Proof:

$$\begin{aligned} E[g(X)] &= \sum_{\omega} g(X(\omega)) Pr[\omega] = \sum_x \sum_{\omega \in X^{-1}(x)} g(X(\omega)) Pr[\omega] \\ &= \sum_x \sum_{\omega \in X^{-1}(x)} g(x) Pr[\omega] = \sum_x g(x) \sum_{\omega \in X^{-1}(x)} Pr[\omega] \end{aligned}$$

Calculating $E[g(X)]$: LOTUS

Let $Y = g(X)$. Assume that we know the distribution of X .

We want to calculate $E[Y]$.

Method 1: We calculate the distribution of Y :

$$Pr[Y = y] = Pr[X \in g^{-1}(y)] \text{ where } g^{-1}(x) = \{x \in \mathfrak{X} : g(x) = y\}.$$

This is typically rather tedious!

Method 2: We use the following result.

Called “Law of the unconscious statistician.”

Theorem:

$$E[g(X)] = \sum_x g(x) Pr[X = x].$$

Proof:

$$\begin{aligned} E[g(X)] &= \sum_{\omega} g(X(\omega)) Pr[\omega] = \sum_x \sum_{\omega \in X^{-1}(x)} g(X(\omega)) Pr[\omega] \\ &= \sum_x \sum_{\omega \in X^{-1}(x)} g(x) Pr[\omega] = \sum_x g(x) \sum_{\omega \in X^{-1}(x)} Pr[\omega] \\ &= \sum_x g(x) Pr[X = x]. \end{aligned}$$



Poll.

Which is LOTUS?

Poll.

Which is LOTUS?

(A) $E[X] = \sum_{x \in \text{Range}(X)} g(x) Pr[g(X) = g(x)]$

(B) $E[X] = \sum_{x \in \text{Range}(X)} g(x) Pr[X = x]$

(C) $E[X] = \sum_{x \in \text{Range}(g)} x Pr[g(X) = x]$

Geometric Distribution.

Experiment: flip a coin with heads prob. p . until Heads.

Random Variable X : number of flips.

Geometric Distribution.

Experiment: flip a coin with heads prob. p . until Heads.

Random Variable X : number of flips.

And distribution is:

Geometric Distribution.

Experiment: flip a coin with heads prob. p . until Heads.

Random Variable X : number of flips.

And distribution is:

$$(A) X \sim G(p) : Pr[X = i] = (1 - p)^{i-1} p.$$

$$(B) X \sim B(p, n) : Pr[X = i] = \binom{n}{i} p^i (1 - p)^{n-i}.$$

Geometric Distribution.

Experiment: flip a coin with heads prob. p . until Heads.

Random Variable X : number of flips.

And distribution is:

$$(A) X \sim G(p) : Pr[X = i] = (1 - p)^{i-1} p.$$

$$(B) X \sim B(p, n) : Pr[X = i] = \binom{n}{i} p^i (1 - p)^{n-i}.$$


$$(A) \text{ Distribution of } X \sim G(p): Pr[X = i] = (1 - p)^{i-1} p.$$

Geometric Distribution: Memoryless - Interpretation

$$Pr[X > n + m | X > n] = Pr[X > m], m, n \geq 0.$$


Geometric Distribution: Memoryless - Interpretation

$$Pr[X > n + m | X > n] = Pr[X > m], m, n \geq 0.$$



Geometric Distribution: Memoryless - Interpretation


$$Pr[X > n + m | X > n] = Pr[X > m], m, n \geq 0.$$



$$Pr[X > n + m | X > n] = Pr[A|B] = Pr[A'] = Pr[X > m].$$

Geometric Distribution: Memoryless - Interpretation

$$Pr[X > n + m | X > n] = Pr[X > m], m, n \geq 0.$$




$$Pr[X > n + m | X > n] = Pr[A|B] = Pr[A'] = Pr[X > m].$$

A' : is m coin tosses before heads.

$A|B$: m 'more' coin tosses before heads.

Geometric Distribution: Memoryless - Interpretation

$$Pr[X > n + m | X > n] = Pr[X > m], m, n \geq 0.$$



$$Pr[X > n + m | X > n] = Pr[A|B] = Pr[A'] = Pr[X > m].$$

A' : is m coin tosses before heads.

$A|B$: m 'more' coin tosses before heads.

The coin is memoryless, therefore, so is X .

Independent coin: $Pr[H | \text{any previous set of coin tosses}] = p$

Geometric Distribution: Memoryless by derivation.

Let X be $G(p)$. Then, for $n \geq 0$,

Geometric Distribution: Memoryless by derivation.

Let X be $G(p)$. Then, for $n \geq 0$,

$$Pr[X > n] = Pr[\text{first } n \text{ flips are } T] =$$

Geometric Distribution: Memoryless by derivation.

Let X be $G(p)$. Then, for $n \geq 0$,

$$Pr[X > n] = Pr[\text{first } n \text{ flips are } T] = (1 - p)^n.$$

Geometric Distribution: Memoryless by derivation.

Let X be $G(p)$. Then, for $n \geq 0$,

$$Pr[X > n] = Pr[\text{first } n \text{ flips are } T] = (1 - p)^n.$$

Theorem

$$Pr[X > n + m | X > n] = Pr[X > m], m, n \geq 0.$$

Geometric Distribution: Memoryless by derivation.

Let X be $G(p)$. Then, for $n \geq 0$,

$$Pr[X > n] = Pr[\text{first } n \text{ flips are } T] = (1 - p)^n.$$

Theorem

$$Pr[X > n + m | X > n] = Pr[X > m], m, n \geq 0.$$

Proof:

$$Pr[X > n + m | X > n] =$$

Geometric Distribution: Memoryless by derivation.

Let X be $G(p)$. Then, for $n \geq 0$,

$$Pr[X > n] = Pr[\text{first } n \text{ flips are } T] = (1 - p)^n.$$

Theorem

$$Pr[X > n + m | X > n] = Pr[X > m], m, n \geq 0.$$

Proof:

$$Pr[X > n + m | X > n] = \frac{Pr[X > n + m \text{ and } X > n]}{Pr[X > n]}$$

Geometric Distribution: Memoryless by derivation.

Let X be $G(p)$. Then, for $n \geq 0$,

$$Pr[X > n] = Pr[\text{first } n \text{ flips are } T] = (1 - p)^n.$$

Theorem

$$Pr[X > n + m | X > n] = Pr[X > m], m, n \geq 0.$$

Proof:

$$\begin{aligned} Pr[X > n + m | X > n] &= \frac{Pr[X > n + m \text{ and } X > n]}{Pr[X > n]} \\ &= \frac{Pr[X > n + m]}{Pr[X > n]} \end{aligned}$$

Geometric Distribution: Memoryless by derivation.

Let X be $G(p)$. Then, for $n \geq 0$,

$$\Pr[X > n] = \Pr[\text{first } n \text{ flips are } T] = (1 - p)^n.$$

Theorem

$$\Pr[X > n + m | X > n] = \Pr[X > m], m, n \geq 0.$$

Proof:

$$\begin{aligned}\Pr[X > n + m | X > n] &= \frac{\Pr[X > n + m \text{ and } X > n]}{\Pr[X > n]} \\ &= \frac{\Pr[X > n + m]}{\Pr[X > n]} \\ &= \frac{(1 - p)^{n+m}}{(1 - p)^n} =\end{aligned}$$

Geometric Distribution: Memoryless by derivation.

Let X be $G(p)$. Then, for $n \geq 0$,

$$Pr[X > n] = Pr[\text{first } n \text{ flips are } T] = (1 - p)^n.$$

Theorem

$$Pr[X > n + m | X > n] = Pr[X > m], m, n \geq 0.$$

Proof:

$$\begin{aligned} Pr[X > n + m | X > n] &= \frac{Pr[X > n + m \text{ and } X > n]}{Pr[X > n]} \\ &= \frac{Pr[X > n + m]}{Pr[X > n]} \\ &= \frac{(1 - p)^{n+m}}{(1 - p)^n} = (1 - p)^m \end{aligned}$$

Geometric Distribution: Memoryless by derivation.

Let X be $G(p)$. Then, for $n \geq 0$,

$$Pr[X > n] = Pr[\text{first } n \text{ flips are } T] = (1 - p)^n.$$

Theorem


$$Pr[X > n + m | X > n] = Pr[X > m], m, n \geq 0.$$

Proof:


$$\begin{aligned} Pr[X > n + m | X > n] &= \frac{Pr[X > n + m \text{ and } X > n]}{Pr[X > n]} \\ &= \frac{Pr[X > n + m]}{Pr[X > n]} \\ &= \frac{(1 - p)^{n+m}}{(1 - p)^n} = (1 - p)^m \\ &= Pr[X > m]. \end{aligned}$$

Variance

Variance




Variance



The variance measures the deviation from the mean value.


Variance



The variance measures the deviation from the mean value.

Definition: The **variance** of X is

Variance




The variance measures the deviation from the mean value.

Definition: The **variance** of X is

$$\sigma^2(X) := \text{var}[X] = E[(X - E[X])^2].$$

Variance




The variance measures the deviation from the mean value.

Definition: The **variance** of X is

$$\sigma^2(X) := \text{var}[X] = E[(X - E[X])^2].$$

$\sigma(X)$ is called the **standard deviation** of X .

Variance



The variance measures the deviation from the mean value.

Definition: The **variance** of X is

$$\sigma^2(X) := \text{var}[X] = E[(X - E[X])^2].$$

$\sigma(X)$ is called the **standard deviation** of X .

Variance and Standard Deviation

Fact:

$$\text{var}[X] = E[X^2] - E[X]^2.$$

Variance and Standard Deviation

Fact:

$$\text{var}[X] = E[X^2] - E[X]^2.$$

Indeed:

$$\text{var}(X) = E[(X - E[X])^2]$$

Variance and Standard Deviation

Fact:

$$\text{var}[X] = E[X^2] - E[X]^2.$$

Indeed:

$$\begin{aligned}\text{var}(X) &= E[(X - E[X])^2] \\ &= E[X^2 - 2XE[X] + E[X]^2]\end{aligned}$$

Variance and Standard Deviation

Fact:

$$\text{var}[X] = E[X^2] - E[X]^2.$$

Indeed:

$$\begin{aligned}\text{var}(X) &= E[(X - E[X])^2] \\ &= E[X^2 - 2XE[X] + E[X]^2] \\ &= E[X^2] - 2E[X]E[X] + E[X]^2,\end{aligned}$$

Variance and Standard Deviation

Fact:

$$\text{var}[X] = E[X^2] - E[X]^2.$$

Indeed:

$$\begin{aligned}\text{var}(X) &= E[(X - E[X])^2] \\ &= E[X^2 - 2XE[X] + E[X]^2] \\ &= E[X^2] - 2E[X]E[X] + E[X]^2, \text{ by linearity}\end{aligned}$$

Variance and Standard Deviation

Fact:

$$\text{var}[X] = E[X^2] - E[X]^2.$$

Indeed:


$$\begin{aligned}\text{var}(X) &= E[(X - E[X])^2] \\ &= E[X^2 - 2XE[X] + E[X]^2] \\ &= E[X^2] - 2E[X]E[X] + E[X]^2, \text{ by linearity} \\ &= E[X^2] - E[X]^2.\end{aligned}$$

A simple example

This example illustrates the term ‘standard deviation.’


A simple example

This example illustrates the term 'standard deviation.'



A simple example

This example illustrates the term ‘standard deviation.’




Consider the random variable X such that

$$X = \begin{cases} \mu - \sigma, & \text{w.p. } 1/2 \\ \mu + \sigma, & \text{w.p. } 1/2. \end{cases}$$

A simple example

This example illustrates the term ‘standard deviation.’




Consider the random variable X such that

$$X = \begin{cases} \mu - \sigma, & \text{w.p. } 1/2 \\ \mu + \sigma, & \text{w.p. } 1/2. \end{cases}$$

Then, $E[X] = \mu$ and $(X - E[X])^2 = \sigma^2$.

A simple example

This example illustrates the term ‘standard deviation.’



Consider the random variable X such that

$$X = \begin{cases} \mu - \sigma, & \text{w.p. } 1/2 \\ \mu + \sigma, & \text{w.p. } 1/2. \end{cases}$$

Then, $E[X] = \mu$ and $(X - E[X])^2 = \sigma^2$. Hence,

$$\text{var}(X) = \sigma^2 \text{ and } \sigma(X) = \sigma.$$

Example

Consider X with

$$X = \begin{cases} -1, & \text{w. p. } 0.99 \\ 99, & \text{w. p. } 0.01. \end{cases}$$

Example

Consider X with

$$X = \begin{cases} -1, & \text{w. p. } 0.99 \\ 99, & \text{w. p. } 0.01. \end{cases}$$

Then

$$E[X] = -1 \times 0.99 + 99 \times 0.01 = 0.$$

Example

Consider X with

$$X = \begin{cases} -1, & \text{w. p. } 0.99 \\ 99, & \text{w. p. } 0.01. \end{cases}$$

Then

$$\begin{aligned} E[X] &= -1 \times 0.99 + 99 \times 0.01 = 0. \\ E[X^2] &= 1 \times 0.99 + (99)^2 \times 0.01 \approx 100. \end{aligned}$$

Example

Consider X with

$$X = \begin{cases} -1, & \text{w. p. } 0.99 \\ 99, & \text{w. p. } 0.01. \end{cases}$$

Then

$$E[X] = -1 \times 0.99 + 99 \times 0.01 = 0.$$

$$E[X^2] = 1 \times 0.99 + (99)^2 \times 0.01 \approx 100.$$

$$\text{Var}(X) \approx 100 \implies \sigma(X) \approx 10.$$

Example

Consider X with

$$X = \begin{cases} -1, & \text{w. p. } 0.99 \\ 99, & \text{w. p. } 0.01. \end{cases}$$

Then

$$E[X] = -1 \times 0.99 + 99 \times 0.01 = 0.$$

$$E[X^2] = 1 \times 0.99 + (99)^2 \times 0.01 \approx 100.$$

$$\text{Var}(X) \approx 100 \implies \sigma(X) \approx 10.$$

Also,

$$E(|X|) = 1 \times 0.99 + 99 \times 0.01 = 1.98.$$

Example

Consider X with

$$X = \begin{cases} -1, & \text{w. p. } 0.99 \\ 99, & \text{w. p. } 0.01. \end{cases}$$

Then

$$E[X] = -1 \times 0.99 + 99 \times 0.01 = 0.$$

$$E[X^2] = 1 \times 0.99 + (99)^2 \times 0.01 \approx 100.$$

$$\text{Var}(X) \approx 100 \implies \sigma(X) \approx 10.$$

Also,

$$E(|X|) = 1 \times 0.99 + 99 \times 0.01 = 1.98.$$

Thus, $\sigma(X) = \sqrt{E[(X - E(X))^2]} \neq E[|X - E[X]|]!$

Example

Consider X with

$$X = \begin{cases} -1, & \text{w. p. } 0.99 \\ 99, & \text{w. p. } 0.01. \end{cases}$$

Then

$$E[X] = -1 \times 0.99 + 99 \times 0.01 = 0.$$

$$E[X^2] = 1 \times 0.99 + (99)^2 \times 0.01 \approx 100.$$

$$\text{Var}(X) \approx 100 \implies \sigma(X) \approx 10.$$

Also,

$$E(|X|) = 1 \times 0.99 + 99 \times 0.01 = 1.98.$$

Thus, $\sigma(X) = \sqrt{E[(X - E(X))^2]} \neq E[|X - E[X]|]!$

Exercise: How big can you make $\frac{\sigma(X)}{E[|X - E[X]|]}$?

Uniform

Assume that $\Pr[X = i] = \frac{1}{n}$ for $i \in \{1, \dots, n\}$. Then

Uniform

Assume that $Pr[X = i] = \frac{1}{n}$ for $i \in \{1, \dots, n\}$. Then

$$E[X] = \sum_{i=1}^n i \times Pr[X = i] = \frac{1}{n} \sum_{i=1}^n i$$

Uniform

Assume that $Pr[X = i] = \frac{1}{n}$ for $i \in \{1, \dots, n\}$. Then

$$\begin{aligned} E[X] &= \sum_{i=1}^n i \times Pr[X = i] = \frac{1}{n} \sum_{i=1}^n i \\ &= \frac{1}{n} \frac{n(n+1)}{2} = \frac{n+1}{2}. \end{aligned}$$

Uniform

Assume that $Pr[X = i] = \frac{1}{n}$ for $i \in \{1, \dots, n\}$. Then

$$\begin{aligned} E[X] &= \sum_{i=1}^n i \times Pr[X = i] = \frac{1}{n} \sum_{i=1}^n i \\ &= \frac{1}{n} \frac{n(n+1)}{2} = \frac{n+1}{2}. \end{aligned}$$

Also,

$$E[X^2] = \sum_{i=1}^n i^2 Pr[X = i] = \frac{1}{n} \sum_{i=1}^n i^2$$

Uniform

Assume that $Pr[X = i] = \frac{1}{n}$ for $i \in \{1, \dots, n\}$. Then

$$\begin{aligned} E[X] &= \sum_{i=1}^n i \times Pr[X = i] = \frac{1}{n} \sum_{i=1}^n i \\ &= \frac{1}{n} \frac{n(n+1)}{2} = \frac{n+1}{2}. \end{aligned}$$

Also,

$$\begin{aligned} E[X^2] &= \sum_{i=1}^n i^2 Pr[X = i] = \frac{1}{n} \sum_{i=1}^n i^2 \\ &= \frac{1}{n} \frac{(n)(n+1)(n+2)}{6} = \frac{1+3n+2n^2}{6}, \end{aligned}$$

Uniform

Assume that $Pr[X = i] = \frac{1}{n}$ for $i \in \{1, \dots, n\}$. Then

$$\begin{aligned} E[X] &= \sum_{i=1}^n i \times Pr[X = i] = \frac{1}{n} \sum_{i=1}^n i \\ &= \frac{1}{n} \frac{n(n+1)}{2} = \frac{n+1}{2}. \end{aligned}$$

Also,

$$\begin{aligned} E[X^2] &= \sum_{i=1}^n i^2 Pr[X = i] = \frac{1}{n} \sum_{i=1}^n i^2 \\ &= \frac{1}{n} \frac{(n)(n+1)(n+2)}{6} = \frac{1+3n+2n^2}{6}, \text{ as you can verify.} \end{aligned}$$

Uniform

Assume that $Pr[X = i] = \frac{1}{n}$ for $i \in \{1, \dots, n\}$. Then

$$\begin{aligned} E[X] &= \sum_{i=1}^n i \times Pr[X = i] = \frac{1}{n} \sum_{i=1}^n i \\ &= \frac{1}{n} \frac{n(n+1)}{2} = \frac{n+1}{2}. \end{aligned}$$

Also,

$$\begin{aligned} E[X^2] &= \sum_{i=1}^n i^2 Pr[X = i] = \frac{1}{n} \sum_{i=1}^n i^2 \\ &= \frac{1}{n} \frac{(n)(n+1)(n+2)}{6} = \frac{1+3n+2n^2}{6}, \text{ as you can verify.} \end{aligned}$$

This gives

$$var(X) = \frac{1+3n+2n^2}{6} - \frac{(n+1)^2}{4} = \frac{n^2-1}{12}.$$

(Sort of $\int_0^{1/2} x^2 dx = \frac{x^3}{3}$.)

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $Pr[X = n] = (1 - p)^{n-1}p$ for $n \geq 1$.

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $Pr[X = n] = (1 - p)^{n-1}p$ for $n \geq 1$. Recall $E[X] = 1/p$.

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $\Pr[X = n] = (1 - p)^{n-1}p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$E[X^2] = p + 4p(1 - p) + 9p(1 - p)^2 + \dots$$

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $Pr[X = n] = (1 - p)^{n-1}p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$\begin{aligned} E[X^2] &= p + 4p(1 - p) + 9p(1 - p)^2 + \dots \\ -(1 - p)E[X^2] &= -[p(1 - p) + 4p(1 - p)^2 + \dots] \end{aligned}$$

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $Pr[X = n] = (1 - p)^{n-1}p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$\begin{aligned}E[X^2] &= p + 4p(1 - p) + 9p(1 - p)^2 + \dots \\-(1 - p)E[X^2] &= -[p(1 - p) + 4p(1 - p)^2 + \dots] \\pE[X^2] &= p + 3p(1 - p) + 5p(1 - p)^2 + \dots\end{aligned}$$

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $\Pr[X = n] = (1 - p)^{n-1}p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$\begin{aligned}E[X^2] &= p + 4p(1 - p) + 9p(1 - p)^2 + \dots \\-(1 - p)E[X^2] &= -[p(1 - p) + 4p(1 - p)^2 + \dots] \\pE[X^2] &= p + 3p(1 - p) + 5p(1 - p)^2 + \dots \\&= 2(p + 2p(1 - p) + 3p(1 - p)^2 + \dots) \quad E[X]! \\&\quad - (p + p(1 - p) + p(1 - p)^2 + \dots) \quad \text{Distribution.}\end{aligned}$$

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $\Pr[X = n] = (1 - p)^{n-1}p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$\begin{aligned}E[X^2] &= p + 4p(1 - p) + 9p(1 - p)^2 + \dots \\-(1 - p)E[X^2] &= -[p(1 - p) + 4p(1 - p)^2 + \dots] \\pE[X^2] &= p + 3p(1 - p) + 5p(1 - p)^2 + \dots \\&= 2(p + 2p(1 - p) + 3p(1 - p)^2 + \dots) \quad E[X]! \\&\quad - (p + p(1 - p) + p(1 - p)^2 + \dots) \quad \text{Distribution.}\end{aligned}$$

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $\Pr[X = n] = (1 - p)^{n-1}p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$\begin{aligned}E[X^2] &= p + 4p(1 - p) + 9p(1 - p)^2 + \dots \\-(1 - p)E[X^2] &= -[p(1 - p) + 4p(1 - p)^2 + \dots] \\pE[X^2] &= p + 3p(1 - p) + 5p(1 - p)^2 + \dots \\&= 2(p + 2p(1 - p) + 3p(1 - p)^2 + \dots) \quad E[X]! \\&\quad -(p + p(1 - p) + p(1 - p)^2 + \dots) \quad \text{Distribution.}\end{aligned}$$

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $\Pr[X = n] = (1 - p)^{n-1}p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$\begin{aligned}E[X^2] &= p + 4p(1 - p) + 9p(1 - p)^2 + \dots \\-(1 - p)E[X^2] &= -[p(1 - p) + 4p(1 - p)^2 + \dots] \\pE[X^2] &= p + 3p(1 - p) + 5p(1 - p)^2 + \dots \\&= 2(p + 2p(1 - p) + 3p(1 - p)^2 + \dots) \quad E[X]! \\&\quad -(p + p(1 - p) + p(1 - p)^2 + \dots) \quad \text{Distribution.} \\pE[X^2] &= 2E[X] - 1\end{aligned}$$

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $\Pr[X = n] = (1 - p)^{n-1}p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$\begin{aligned}E[X^2] &= p + 4p(1 - p) + 9p(1 - p)^2 + \dots \\-(1 - p)E[X^2] &= -[p(1 - p) + 4p(1 - p)^2 + \dots] \\pE[X^2] &= p + 3p(1 - p) + 5p(1 - p)^2 + \dots \\&= 2(p + 2p(1 - p) + 3p(1 - p)^2 + \dots) \quad E[X]! \\&\quad -(p + p(1 - p) + p(1 - p)^2 + \dots) \quad \text{Distribution.} \\pE[X^2] &= 2E[X] - 1 \\&= 2\left(\frac{1}{p}\right) - 1\end{aligned}$$

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $\Pr[X = n] = (1 - p)^{n-1}p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$\begin{aligned}E[X^2] &= p + 4p(1 - p) + 9p(1 - p)^2 + \dots \\-(1 - p)E[X^2] &= -[p(1 - p) + 4p(1 - p)^2 + \dots] \\pE[X^2] &= p + 3p(1 - p) + 5p(1 - p)^2 + \dots \\&= 2(p + 2p(1 - p) + 3p(1 - p)^2 + \dots) \quad E[X]! \\&\quad -(p + p(1 - p) + p(1 - p)^2 + \dots) \quad \text{Distribution.} \\pE[X^2] &= 2E[X] - 1 \\&= 2\left(\frac{1}{p}\right) - 1 = \frac{2 - p}{p}\end{aligned}$$

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $\Pr[X = n] = (1 - p)^{n-1}p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$E[X^2] = p + 4p(1 - p) + 9p(1 - p)^2 + \dots$$

$$-(1 - p)E[X^2] = -[p(1 - p) + 4p(1 - p)^2 + \dots]$$

$$pE[X^2] = p + 3p(1 - p) + 5p(1 - p)^2 + \dots$$

$$= 2(p + 2p(1 - p) + 3p(1 - p)^2 + \dots) \quad E[X]!$$

$$-(p + p(1 - p) + p(1 - p)^2 + \dots) \quad \text{Distribution.}$$

$$pE[X^2] = 2E[X] - 1$$

$$= 2\left(\frac{1}{p}\right) - 1 = \frac{2 - p}{p}$$

$$\implies E[X^2] = (2 - p)/p^2$$

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $\Pr[X = n] = (1 - p)^{n-1}p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$\begin{aligned}E[X^2] &= p + 4p(1-p) + 9p(1-p)^2 + \dots \\-(1-p)E[X^2] &= -[p(1-p) + 4p(1-p)^2 + \dots] \\pE[X^2] &= p + 3p(1-p) + 5p(1-p)^2 + \dots \\&= 2(p + 2p(1-p) + 3p(1-p)^2 + \dots) \quad E[X]! \\&\quad -(p + p(1-p) + p(1-p)^2 + \dots) \quad \text{Distribution.} \\pE[X^2] &= 2E[X] - 1 \\&= 2\left(\frac{1}{p}\right) - 1 = \frac{2-p}{p}\end{aligned}$$

$$\begin{aligned}\implies E[X^2] &= (2-p)/p^2 \text{ and} \\ \text{var}[X] &= E[X^2] - E[X]^2\end{aligned}$$

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $\Pr[X = n] = (1 - p)^{n-1} p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$\begin{aligned} E[X^2] &= p + 4p(1-p) + 9p(1-p)^2 + \dots \\ -(1-p)E[X^2] &= -[p(1-p) + 4p(1-p)^2 + \dots] \\ pE[X^2] &= p + 3p(1-p) + 5p(1-p)^2 + \dots \\ &= 2(p + 2p(1-p) + 3p(1-p)^2 + \dots) \quad E[X]! \\ &\quad -(p + p(1-p) + p(1-p)^2 + \dots) \quad \text{Distribution.} \\ pE[X^2] &= 2E[X] - 1 \\ &= 2\left(\frac{1}{p}\right) - 1 = \frac{2-p}{p} \end{aligned}$$

$$\begin{aligned} \implies E[X^2] &= (2-p)/p^2 \text{ and} \\ \text{var}[X] &= E[X^2] - E[X]^2 = \frac{2-p}{p^2} - \frac{1}{p^2} \end{aligned}$$

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $\Pr[X = n] = (1 - p)^{n-1} p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$\begin{aligned}E[X^2] &= p + 4p(1-p) + 9p(1-p)^2 + \dots \\-(1-p)E[X^2] &= -[p(1-p) + 4p(1-p)^2 + \dots] \\pE[X^2] &= p + 3p(1-p) + 5p(1-p)^2 + \dots \\&= 2(p + 2p(1-p) + 3p(1-p)^2 + \dots) \quad E[X]! \\&\quad -(p + p(1-p) + p(1-p)^2 + \dots) \quad \text{Distribution.} \\pE[X^2] &= 2E[X] - 1 \\&= 2\left(\frac{1}{p}\right) - 1 = \frac{2-p}{p}\end{aligned}$$

$$\begin{aligned}\implies E[X^2] &= (2-p)/p^2 \text{ and} \\var[X] &= E[X^2] - E[X]^2 = \frac{2-p}{p^2} - \frac{1}{p^2} = \frac{1-p}{p^2}.\end{aligned}$$

$$\sigma(X) = \frac{\sqrt{1-p}}{p}$$

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $\Pr[X = n] = (1 - p)^{n-1} p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$\begin{aligned}E[X^2] &= p + 4p(1-p) + 9p(1-p)^2 + \dots \\-(1-p)E[X^2] &= -[p(1-p) + 4p(1-p)^2 + \dots] \\pE[X^2] &= p + 3p(1-p) + 5p(1-p)^2 + \dots \\&= 2(p + 2p(1-p) + 3p(1-p)^2 + \dots) \quad E[X]! \\&\quad -(p + p(1-p) + p(1-p)^2 + \dots) \quad \text{Distribution.} \\pE[X^2] &= 2E[X] - 1 \\&= 2\left(\frac{1}{p}\right) - 1 = \frac{2-p}{p}\end{aligned}$$

$$\begin{aligned}\implies E[X^2] &= (2-p)/p^2 \text{ and} \\var[X] &= E[X^2] - E[X]^2 = \frac{2-p}{p^2} - \frac{1}{p^2} = \frac{1-p}{p^2}. \\ \sigma(X) &= \frac{\sqrt{1-p}}{p} \approx E[X] \text{ when } p \text{ is small(ish).}\end{aligned}$$

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $\Pr[X = n] = (1 - p)^{n-1} p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$\begin{aligned}E[X^2] &= p + 4p(1-p) + 9p(1-p)^2 + \dots \\-(1-p)E[X^2] &= -[p(1-p) + 4p(1-p)^2 + \dots] \\pE[X^2] &= p + 3p(1-p) + 5p(1-p)^2 + \dots \\&= 2(p + 2p(1-p) + 3p(1-p)^2 + \dots) \quad E[X]! \\&\quad -(p + p(1-p) + p(1-p)^2 + \dots) \quad \text{Distribution.} \\pE[X^2] &= 2E[X] - 1 \\&= 2\left(\frac{1}{p}\right) - 1 = \frac{2-p}{p}\end{aligned}$$

$$\begin{aligned}\implies E[X^2] &= (2-p)/p^2 \text{ and} \\var[X] &= E[X^2] - E[X]^2 = \frac{2-p}{p^2} - \frac{1}{p^2} = \frac{1-p}{p^2}. \\ \sigma(X) &= \frac{\sqrt{1-p}}{p} \approx E[X] \text{ when } p \text{ is small(ish).}\end{aligned}$$

Fixed points.

Number of fixed points in a random permutation of n items.

Fixed points.

Number of fixed points in a random permutation of n items.

“Number of student that get homework back.”

Fixed points.

Number of fixed points in a random permutation of n items.

“Number of student that get homework back.”

$$X = X_1 + X_2 \cdots + X_n$$

Fixed points.

Number of fixed points in a random permutation of n items.

“Number of student that get homework back.”

$$X = X_1 + X_2 \cdots + X_n$$

where X_i is indicator variable for i th student getting hw back.

Fixed points.

Number of fixed points in a random permutation of n items.

“Number of student that get homework back.”

$$X = X_1 + X_2 + \cdots + X_n$$

where X_i is indicator variable for i th student getting hw back.

$$\begin{aligned} E(X^2) &= \sum_i E(X_i^2) + \sum_{i \neq j} E(X_i X_j). \\ &= \quad \quad \quad + \end{aligned}$$

Fixed points.

Number of fixed points in a random permutation of n items.

“Number of student that get homework back.”

$$X = X_1 + X_2 + \cdots + X_n$$

where X_i is indicator variable for i th student getting hw back.

$$\begin{aligned} E(X^2) &= \sum_i E(X_i^2) + \sum_{i \neq j} E(X_i X_j). \\ &= \quad + \end{aligned}$$

$$E(X_i^2) = 1 \times \Pr[X_i = 1] + 0 \times \Pr[X_i = 0]$$

Fixed points.

Number of fixed points in a random permutation of n items.

“Number of student that get homework back.”

$$X = X_1 + X_2 + \cdots + X_n$$

where X_i is indicator variable for i th student getting hw back.

$$\begin{aligned} E(X^2) &= \sum_i E(X_i^2) + \sum_{i \neq j} E(X_i X_j). \\ &= \quad + \end{aligned}$$

$$\begin{aligned} E(X_i^2) &= 1 \times \Pr[X_i = 1] + 0 \times \Pr[X_i = 0] \\ &= \frac{1}{n} \end{aligned}$$

Fixed points.

Number of fixed points in a random permutation of n items.

“Number of student that get homework back.”

$$X = X_1 + X_2 \cdots + X_n$$

where X_i is indicator variable for i th student getting hw back.

$$\begin{aligned} E(X^2) &= \sum_i E(X_i^2) + \sum_{i \neq j} E(X_i X_j). \\ &= \quad \quad \quad + \end{aligned}$$

$$\begin{aligned} E(X_i^2) &= 1 \times \Pr[X_i = 1] + 0 \times \Pr[X_i = 0] \\ &= \frac{1}{n} \end{aligned}$$

Fixed points.

Number of fixed points in a random permutation of n items.

“Number of student that get homework back.”

$$X = X_1 + X_2 + \cdots + X_n$$

where X_i is indicator variable for i th student getting hw back.

$$\begin{aligned} E(X^2) &= \sum_i E(X_i^2) + \sum_{i \neq j} E(X_i X_j). \\ &= n \times \frac{1}{n} + \end{aligned}$$

$$\begin{aligned} E(X_i^2) &= 1 \times \Pr[X_i = 1] + 0 \times \Pr[X_i = 0] \\ &= \frac{1}{n} \end{aligned}$$

Fixed points.

Number of fixed points in a random permutation of n items.

“Number of student that get homework back.”

$$X = X_1 + X_2 + \cdots + X_n$$

where X_i is indicator variable for i th student getting hw back.

$$\begin{aligned} E(X^2) &= \sum_i E(X_i^2) + \sum_{i \neq j} E(X_i X_j). \\ &= n \times \frac{1}{n} + \end{aligned}$$

$$E(X_i^2) = 1 \times \Pr[X_i = 1] + 0 \times \Pr[X_i = 0]$$

$$= \frac{1}{n}$$

$$E(X_i X_j) = 1 \times \Pr[X_i = 1 \cap X_j = 1] + 0 \times \Pr[\text{“anything else”}]$$

Fixed points.

Number of fixed points in a random permutation of n items.

“Number of student that get homework back.”

$$X = X_1 + X_2 \cdots + X_n$$

where X_i is indicator variable for i th student getting hw back.

$$\begin{aligned} E(X^2) &= \sum_i E(X_i^2) + \sum_{i \neq j} E(X_i X_j). \\ &= n \times \frac{1}{n} + \end{aligned}$$

$$E(X_i^2) = 1 \times \Pr[X_i = 1] + 0 \times \Pr[X_i = 0]$$

$$= \frac{1}{n}$$

$$E(X_i X_j) = 1 \times \Pr[X_i = 1 \cap X_j = 1] + 0 \times \Pr[\text{“anything else”}]$$

$$= 1 \times \frac{(n-2)!}{n!}$$

Fixed points.

Number of fixed points in a random permutation of n items.

“Number of student that get homework back.”

$$X = X_1 + X_2 + \cdots + X_n$$

where X_i is indicator variable for i th student getting hw back.

$$\begin{aligned} E(X^2) &= \sum_i E(X_i^2) + \sum_{i \neq j} E(X_i X_j). \\ &= n \times \frac{1}{n} + \end{aligned}$$

$$E(X_i^2) = 1 \times \Pr[X_i = 1] + 0 \times \Pr[X_i = 0]$$

$$= \frac{1}{n}$$

$$E(X_i X_j) = 1 \times \Pr[X_i = 1 \cap X_j = 1] + 0 \times \Pr[\text{“anything else”}]$$

$$= 1 \times \frac{(n-2)!}{n!} = \frac{1}{n(n-1)}$$

Fixed points.

Number of fixed points in a random permutation of n items.

“Number of student that get homework back.”

$$X = X_1 + X_2 \cdots + X_n$$

where X_i is indicator variable for i th student getting hw back.

$$\begin{aligned} E(X^2) &= \sum_i E(X_i^2) + \sum_{i \neq j} E(X_i X_j). \\ &= n \times \frac{1}{n} + (n)(n-1) \times \frac{1}{n(n-1)} \end{aligned}$$

$$E(X_i^2) = 1 \times \Pr[X_i = 1] + 0 \times \Pr[X_i = 0]$$

$$\begin{aligned} E(X_i X_j) &= \frac{1}{n} \times \Pr[X_i = 1 \cap X_j = 1] + 0 \times \Pr[\text{“anything else”}] \\ &= 1 \times \frac{(n-2)!}{n!} = \frac{1}{n(n-1)} \end{aligned}$$

Fixed points.

Number of fixed points in a random permutation of n items.

“Number of student that get homework back.”

$$X = X_1 + X_2 \cdots + X_n$$

where X_i is indicator variable for i th student getting hw back.

$$\begin{aligned} E(X^2) &= \sum_i E(X_i^2) + \sum_{i \neq j} E(X_i X_j). \\ &= n \times \frac{1}{n} + (n)(n-1) \times \frac{1}{n(n-1)} \\ &= 1 + 1 = 2. \end{aligned}$$

$$E(X_i^2) = 1 \times \Pr[X_i = 1] + 0 \times \Pr[X_i = 0]$$

$$\begin{aligned} E(X_i X_j) &= \frac{1}{n} \times \Pr[X_i = 1 \cap X_j = 1] + 0 \times \Pr[\text{“anything else”}] \\ &= 1 \times \frac{(n-2)!}{n!} = \frac{1}{n(n-1)} \end{aligned}$$

Fixed points.

Number of fixed points in a random permutation of n items.

“Number of student that get homework back.”

$$X = X_1 + X_2 \cdots + X_n$$

where X_i is indicator variable for i th student getting hw back.

$$\begin{aligned} E(X^2) &= \sum_i E(X_i^2) + \sum_{i \neq j} E(X_i X_j). \\ &= n \times \frac{1}{n} + (n)(n-1) \times \frac{1}{n(n-1)} \\ &= 1 + 1 = 2. \end{aligned}$$

$$E(X_i^2) = 1 \times \Pr[X_i = 1] + 0 \times \Pr[X_i = 0]$$

$$\begin{aligned} E(X_i X_j) &= \frac{1}{n} \times \Pr[X_i = 1 \cap X_j = 1] + 0 \times \Pr[\text{“anything else”}] \\ &= 1 \times \frac{(n-2)!}{n!} = \frac{1}{n(n-1)} \end{aligned}$$

$$\text{Var}(X) = E(X^2) - (E(X))^2 = 2 - 1 = 1.$$

Poll: fixed points.

What's true?

Poll: fixed points.

What's true?

(A) X_i and X_j are independent.

(B) $E[X_i X_j] = Pr[X_i X_j = 1]$

(C) $Pr[X_i X_j] = \frac{(n-2)!}{n!}$

(D) $X_i^2 = X_i$.

Variance: binomial.

$$E[X^2] = \sum_{i=0}^n i^2 \binom{n}{i} p^i (1-p)^{n-i}.$$

Variance: binomial.

$$\begin{aligned} E[X^2] &= \sum_{i=0}^n i^2 \binom{n}{i} p^i (1-p)^{n-i}. \\ &= \end{aligned}$$

Variance: binomial.

$$\begin{aligned} E[X^2] &= \sum_{i=0}^n i^2 \binom{n}{i} p^i (1-p)^{n-i}. \\ &= \text{Really???!!##...} \end{aligned}$$

Too hard!

Variance: binomial.

$$\begin{aligned} E[X^2] &= \sum_{i=0}^n i^2 \binom{n}{i} p^i (1-p)^{n-i}. \\ &= \text{Really???!!##...} \end{aligned}$$

Too hard!

Ok..

Variance: binomial.

$$\begin{aligned} E[X^2] &= \sum_{i=0}^n i^2 \binom{n}{i} p^i (1-p)^{n-i}. \\ &= \text{Really???!!##...} \end{aligned}$$

Too hard!

Ok.. fine.

Variance: binomial.

$$\begin{aligned} E[X^2] &= \sum_{i=0}^n i^2 \binom{n}{i} p^i (1-p)^{n-i}. \\ &= \text{Really???!!##...} \end{aligned}$$

Too hard!

Ok.. fine.

Let's do something else.

Variance: binomial.

$$\begin{aligned} E[X^2] &= \sum_{i=0}^n i^2 \binom{n}{i} p^i (1-p)^{n-i}. \\ &= \text{Really???!!##...} \end{aligned}$$

Too hard!

Ok.. fine.

Let's do something else.

Maybe not much easier...

Variance: binomial.

$$\begin{aligned} E[X^2] &= \sum_{i=0}^n i^2 \binom{n}{i} p^i (1-p)^{n-i}. \\ &= \text{Really???!!##...} \end{aligned}$$

Too hard!

Ok.. fine.

Let's do something else.

Maybe not much easier...but there is a payoff.

Properties of variance.

1. $\text{Var}(cX) = c^2 \text{Var}(X)$, where c is a constant.

Properties of variance.

1. $\text{Var}(cX) = c^2 \text{Var}(X)$, where c is a constant.
Scales by c^2 .

Properties of variance.

1. $\text{Var}(cX) = c^2 \text{Var}(X)$, where c is a constant.
Scales by c^2 .
2. $\text{Var}(X + c) = \text{Var}(X)$, where c is a constant.

Properties of variance.

1. $\text{Var}(cX) = c^2 \text{Var}(X)$, where c is a constant.
Scales by c^2 .
2. $\text{Var}(X + c) = \text{Var}(X)$, where c is a constant.
Shifts center.

Properties of variance.

1. $Var(cX) = c^2 Var(X)$, where c is a constant.
Scales by c^2 .
2. $Var(X + c) = Var(X)$, where c is a constant.
Shifts center.

Proof:

$$Var(cX) = E((cX)^2) - (E(cX))^2$$

Properties of variance.

1. $Var(cX) = c^2 Var(X)$, where c is a constant.
Scales by c^2 .
2. $Var(X + c) = Var(X)$, where c is a constant.
Shifts center.

Proof:

$$\begin{aligned} Var(cX) &= E((cX)^2) - (E(cX))^2 \\ &= c^2 E(X^2) - c^2 (E(X))^2 \end{aligned}$$

Properties of variance.

1. $Var(cX) = c^2 Var(X)$, where c is a constant.
Scales by c^2 .
2. $Var(X + c) = Var(X)$, where c is a constant.
Shifts center.

Proof:

$$\begin{aligned} Var(cX) &= E((cX)^2) - (E(cX))^2 \\ &= c^2 E(X^2) - c^2 (E(X))^2 = c^2 (E(X^2) - E(X)^2) \end{aligned}$$

Properties of variance.

1. $Var(cX) = c^2 Var(X)$, where c is a constant.
Scales by c^2 .
2. $Var(X + c) = Var(X)$, where c is a constant.
Shifts center.

Proof:

$$\begin{aligned} Var(cX) &= E((cX)^2) - (E(cX))^2 \\ &= c^2 E(X^2) - c^2 (E(X))^2 = c^2 (E(X^2) - E(X)^2) \\ &= c^2 Var(X) \end{aligned}$$

Properties of variance.

1. $Var(cX) = c^2 Var(X)$, where c is a constant.
Scales by c^2 .
2. $Var(X + c) = Var(X)$, where c is a constant.
Shifts center.

Proof:

$$\begin{aligned} Var(cX) &= E((cX)^2) - (E(cX))^2 \\ &= c^2 E(X^2) - c^2 (E(X))^2 = c^2 (E(X^2) - E(X)^2) \\ &= c^2 Var(X) \\ Var(X + c) &= E((X + c - E(X + c))^2) \end{aligned}$$

Properties of variance.

1. $Var(cX) = c^2 Var(X)$, where c is a constant.
Scales by c^2 .
2. $Var(X + c) = Var(X)$, where c is a constant.
Shifts center.

Proof:

$$\begin{aligned} Var(cX) &= E((cX)^2) - (E(cX))^2 \\ &= c^2 E(X^2) - c^2 (E(X))^2 = c^2 (E(X^2) - E(X)^2) \\ &= c^2 Var(X) \\ Var(X + c) &= E((X + c - E(X + c))^2) \\ &= E((X + c - E(X) - c)^2) \end{aligned}$$

Properties of variance.

1. $Var(cX) = c^2 Var(X)$, where c is a constant.
Scales by c^2 .
2. $Var(X + c) = Var(X)$, where c is a constant.
Shifts center.

Proof:

$$\begin{aligned} Var(cX) &= E((cX)^2) - (E(cX))^2 \\ &= c^2 E(X^2) - c^2 (E(X))^2 = c^2 (E(X^2) - E(X)^2) \\ &= c^2 Var(X) \end{aligned}$$

$$\begin{aligned} Var(X + c) &= E((X + c - E(X + c))^2) \\ &= E((X + c - E(X) - c)^2) \\ &= E((X - E(X))^2) \end{aligned}$$

Properties of variance.

1. $Var(cX) = c^2 Var(X)$, where c is a constant.
Scales by c^2 .
2. $Var(X + c) = Var(X)$, where c is a constant.
Shifts center.

Proof:

$$\begin{aligned} Var(cX) &= E((cX)^2) - (E(cX))^2 \\ &= c^2 E(X^2) - c^2 (E(X))^2 = c^2 (E(X^2) - E(X)^2) \\ &= c^2 Var(X) \end{aligned}$$

$$\begin{aligned} Var(X + c) &= E((X + c - E(X + c))^2) \\ &= E((X + c - E(X) - c)^2) \\ &= E((X - E(X))^2) = Var(X) \end{aligned}$$

Properties of variance.

1. $Var(cX) = c^2 Var(X)$, where c is a constant.
Scales by c^2 .
2. $Var(X + c) = Var(X)$, where c is a constant.
Shifts center.

Proof:

$$\begin{aligned} Var(cX) &= E((cX)^2) - (E(cX))^2 \\ &= c^2 E(X^2) - c^2 (E(X))^2 = c^2 (E(X^2) - E(X)^2) \\ &= c^2 Var(X) \\ Var(X + c) &= E((X + c - E(X + c))^2) \\ &= E((X + c - E(X) - c)^2) \\ &= E((X - E(X))^2) = Var(X) \end{aligned}$$



Independent random variables.

Independent: $P[X = a, Y = b] = Pr[X = a]Pr[Y = b]$

Independent random variables.

Independent: $P[X = a, Y = b] = Pr[X = a]Pr[Y = b]$

Fact: $E[XY] = E[X]E[Y]$ for independent random variables.

Independent random variables.

Independent: $P[X = a, Y = b] = Pr[X = a]Pr[Y = b]$

Fact: $E[XY] = E[X]E[Y]$ for independent random variables.

$$\begin{aligned} E[XY] &= \sum_a \sum_b a \times b \times Pr[X = a, Y = b] \\ &= \sum_a \sum_b a \times b \times Pr[X = a]Pr[Y = b] \\ &= \left(\sum_a a Pr[X = a]\right) \left(\sum_b b Pr[Y = b]\right) \\ &= E[X]E[Y] \end{aligned}$$

Variance of sum of two independent random variables

Variance of sum of two independent random variables

Theorem:

If X and Y are independent, then

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y).$$

Variance of sum of two independent random variables

Theorem:

If X and Y are independent, then

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y).$$

Proof:

Since shifting the random variables does not change their variance, let us subtract their means.

Variance of sum of two independent random variables

Theorem:

If X and Y are independent, then

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y).$$

Proof:

Since shifting the random variables does not change their variance, let us subtract their means.

That is, we assume that $E(X) = 0$ and $E(Y) = 0$.

Variance of sum of two independent random variables

Theorem:

If X and Y are independent, then

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y).$$

Proof:

Since shifting the random variables does not change their variance, let us subtract their means.

That is, we assume that $E(X) = 0$ and $E(Y) = 0$.

Then, by independence,

$$E(XY) = E(X)E(Y) = 0.$$

Variance of sum of two independent random variables

Theorem:

If X and Y are independent, then

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y).$$

Proof:

Since shifting the random variables does not change their variance, let us subtract their means.

That is, we assume that $E(X) = 0$ and $E(Y) = 0$.

Then, by independence,

$$E(XY) = E(X)E(Y) = 0.$$

Hence,

$$\text{var}(X + Y) = E((X + Y)^2)$$

Variance of sum of two independent random variables

Theorem:

If X and Y are independent, then

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y).$$

Proof:

Since shifting the random variables does not change their variance, let us subtract their means.

That is, we assume that $E(X) = 0$ and $E(Y) = 0$.

Then, by independence,

$$E(XY) = E(X)E(Y) = 0.$$

Hence,

$$\text{var}(X + Y) = E((X + Y)^2) = E(X^2 + 2XY + Y^2)$$

Variance of sum of two independent random variables

Theorem:

If X and Y are independent, then

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y).$$

Proof:

Since shifting the random variables does not change their variance, let us subtract their means.

That is, we assume that $E(X) = 0$ and $E(Y) = 0$.

Then, by independence,

$$E(XY) = E(X)E(Y) = 0.$$

Hence,

$$\begin{aligned}\text{var}(X + Y) &= E((X + Y)^2) = E(X^2 + 2XY + Y^2) \\ &= E(X^2) + 2E(XY) + E(Y^2)\end{aligned}$$

Variance of sum of two independent random variables

Theorem:

If X and Y are independent, then

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y).$$

Proof:

Since shifting the random variables does not change their variance, let us subtract their means.

That is, we assume that $E(X) = 0$ and $E(Y) = 0$.

Then, by independence,

$$E(XY) = E(X)E(Y) = 0.$$

Hence,

$$\begin{aligned}\text{var}(X + Y) &= E((X + Y)^2) = E(X^2 + 2XY + Y^2) \\ &= E(X^2) + 2E(XY) + E(Y^2) = E(X^2) + E(Y^2)\end{aligned}$$

Variance of sum of two independent random variables

Theorem:

If X and Y are independent, then

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y).$$

Proof:

Since shifting the random variables does not change their variance, let us subtract their means.

That is, we assume that $E(X) = 0$ and $E(Y) = 0$.

Then, by independence,

$$E(XY) = E(X)E(Y) = 0.$$

Hence,

$$\begin{aligned}\text{var}(X + Y) &= E((X + Y)^2) = E(X^2 + 2XY + Y^2) \\ &= E(X^2) + 2E(XY) + E(Y^2) = E(X^2) + E(Y^2) \\ &= \text{var}(X) + \text{var}(Y).\end{aligned}$$

Variance of sum of independent random variables

Variance of sum of independent random variables

Theorem:

If X, Y, Z, \dots are pairwise independent, then

$$\text{var}(X + Y + Z + \dots) = \text{var}(X) + \text{var}(Y) + \text{var}(Z) + \dots .$$

Variance of sum of independent random variables

Theorem:

If X, Y, Z, \dots are pairwise independent, then

$$\text{var}(X + Y + Z + \dots) = \text{var}(X) + \text{var}(Y) + \text{var}(Z) + \dots .$$

Proof:

Since shifting the random variables does not change their variance, let us subtract their means.

Variance of sum of independent random variables

Theorem:

If X, Y, Z, \dots are pairwise independent, then

$$\text{var}(X + Y + Z + \dots) = \text{var}(X) + \text{var}(Y) + \text{var}(Z) + \dots.$$

Proof:

Since shifting the random variables does not change their variance, let us subtract their means.

That is, we assume that $E[X] = E[Y] = \dots = 0$.

Variance of sum of independent random variables

Theorem:

If X, Y, Z, \dots are pairwise independent, then

$$\text{var}(X + Y + Z + \dots) = \text{var}(X) + \text{var}(Y) + \text{var}(Z) + \dots.$$

Proof:

Since shifting the random variables does not change their variance, let us subtract their means.

That is, we assume that $E[X] = E[Y] = \dots = 0$.

Then, by independence,

$$E[XY] = E[X]E[Y] = 0.$$

Variance of sum of independent random variables

Theorem:

If X, Y, Z, \dots are pairwise independent, then

$$\text{var}(X + Y + Z + \dots) = \text{var}(X) + \text{var}(Y) + \text{var}(Z) + \dots .$$

Proof:

Since shifting the random variables does not change their variance, let us subtract their means.

That is, we assume that $E[X] = E[Y] = \dots = 0$.

Then, by independence,

$$E[XY] = E[X]E[Y] = 0. \text{ Also, } E[XZ] = E[YZ] = \dots = 0.$$

Variance of sum of independent random variables

Theorem:

If X, Y, Z, \dots are pairwise independent, then

$$\text{var}(X + Y + Z + \dots) = \text{var}(X) + \text{var}(Y) + \text{var}(Z) + \dots.$$

Proof:

Since shifting the random variables does not change their variance, let us subtract their means.

That is, we assume that $E[X] = E[Y] = \dots = 0$.

Then, by independence,

$$E[XY] = E[X]E[Y] = 0. \text{ Also, } E[XZ] = E[YZ] = \dots = 0.$$

Hence,

$$\text{var}(X + Y + Z + \dots) = E((X + Y + Z + \dots)^2)$$

Variance of sum of independent random variables

Theorem:

If X, Y, Z, \dots are pairwise independent, then

$$\text{var}(X + Y + Z + \dots) = \text{var}(X) + \text{var}(Y) + \text{var}(Z) + \dots.$$

Proof:

Since shifting the random variables does not change their variance, let us subtract their means.

That is, we assume that $E[X] = E[Y] = \dots = 0$.

Then, by independence,

$$E[XY] = E[X]E[Y] = 0. \text{ Also, } E[XZ] = E[YZ] = \dots = 0.$$

Hence,

$$\begin{aligned}\text{var}(X + Y + Z + \dots) &= E((X + Y + Z + \dots)^2) \\ &= E(X^2 + Y^2 + Z^2 + \dots + 2XY + 2XZ + 2YZ + \dots)\end{aligned}$$

Variance of sum of independent random variables

Theorem:

If X, Y, Z, \dots are pairwise independent, then

$$\text{var}(X + Y + Z + \dots) = \text{var}(X) + \text{var}(Y) + \text{var}(Z) + \dots.$$

Proof:

Since shifting the random variables does not change their variance, let us subtract their means.

That is, we assume that $E[X] = E[Y] = \dots = 0$.

Then, by independence,

$$E[XY] = E[X]E[Y] = 0. \text{ Also, } E[XZ] = E[YZ] = \dots = 0.$$

Hence,

$$\begin{aligned}\text{var}(X + Y + Z + \dots) &= E((X + Y + Z + \dots)^2) \\ &= E(X^2 + Y^2 + Z^2 + \dots + 2XY + 2XZ + 2YZ + \dots) \\ &= E(X^2) + E(Y^2) + E(Z^2) + \dots + 0 + \dots + 0\end{aligned}$$

Variance of sum of independent random variables

Theorem:

If X, Y, Z, \dots are pairwise independent, then

$$\text{var}(X + Y + Z + \dots) = \text{var}(X) + \text{var}(Y) + \text{var}(Z) + \dots.$$

Proof:

Since shifting the random variables does not change their variance, let us subtract their means.

That is, we assume that $E[X] = E[Y] = \dots = 0$.

Then, by independence,

$$E[XY] = E[X]E[Y] = 0. \text{ Also, } E[XZ] = E[YZ] = \dots = 0.$$

Hence,

$$\begin{aligned} \text{var}(X + Y + Z + \dots) &= E((X + Y + Z + \dots)^2) \\ &= E(X^2 + Y^2 + Z^2 + \dots + 2XY + 2XZ + 2YZ + \dots) \\ &= E(X^2) + E(Y^2) + E(Z^2) + \dots + 0 + \dots + 0 \\ &= \text{var}(X) + \text{var}(Y) + \text{var}(Z) + \dots. \end{aligned}$$



Variance of Binomial Distribution.

Flip coin with heads probability p .

Variance of Binomial Distribution.

Flip coin with heads probability p .

X - how many heads?

Variance of Binomial Distribution.

Flip coin with heads probability p .

X - how many heads?

$$X_i = \begin{cases} 1 & \text{if } i\text{th flip is heads} \\ 0 & \text{otherwise} \end{cases}$$

Variance of Binomial Distribution.

Flip coin with heads probability p .

X - how many heads?

$$X_i = \begin{cases} 1 & \text{if } i\text{th flip is heads} \\ 0 & \text{otherwise} \end{cases}$$

$$E(X_i^2)$$

Variance of Binomial Distribution.

Flip coin with heads probability p .

X - how many heads?

$$X_i = \begin{cases} 1 & \text{if } i\text{th flip is heads} \\ 0 & \text{otherwise} \end{cases}$$

$$E(X_i^2) = 1^2 \times p + 0^2 \times (1 - p)$$

Variance of Binomial Distribution.

Flip coin with heads probability p .

X - how many heads?

$$X_i = \begin{cases} 1 & \text{if } i\text{th flip is heads} \\ 0 & \text{otherwise} \end{cases}$$

$$E(X_i^2) = 1^2 \times p + 0^2 \times (1 - p) = p.$$

Variance of Binomial Distribution.

Flip coin with heads probability p .

X - how many heads?

$$X_i = \begin{cases} 1 & \text{if } i\text{th flip is heads} \\ 0 & \text{otherwise} \end{cases}$$

$$E(X_i^2) = 1^2 \times p + 0^2 \times (1 - p) = p.$$

$$\text{Var}(X_i) = p - (E(X))^2$$

Variance of Binomial Distribution.

Flip coin with heads probability p .

X - how many heads?

$$X_i = \begin{cases} 1 & \text{if } i\text{th flip is heads} \\ 0 & \text{otherwise} \end{cases}$$

$$E(X_i^2) = 1^2 \times p + 0^2 \times (1 - p) = p.$$

$$\text{Var}(X_i) = p - (E(X))^2 = p - p^2$$

Variance of Binomial Distribution.

Flip coin with heads probability p .

X - how many heads?

$$X_i = \begin{cases} 1 & \text{if } i\text{th flip is heads} \\ 0 & \text{otherwise} \end{cases}$$

$$E(X_i^2) = 1^2 \times p + 0^2 \times (1 - p) = p.$$

$$\text{Var}(X_i) = p - (E(X_i))^2 = p - p^2 = p(1 - p).$$

Variance of Binomial Distribution.

Flip coin with heads probability p .

X - how many heads?

$$X_i = \begin{cases} 1 & \text{if } i\text{th flip is heads} \\ 0 & \text{otherwise} \end{cases}$$

$$E(X_i^2) = 1^2 \times p + 0^2 \times (1 - p) = p.$$

$$\text{Var}(X_i) = p - (E(X_i))^2 = p - p^2 = p(1 - p).$$

$$p = 0$$

Variance of Binomial Distribution.

Flip coin with heads probability p .

X - how many heads?

$$X_i = \begin{cases} 1 & \text{if } i\text{th flip is heads} \\ 0 & \text{otherwise} \end{cases}$$

$$E(X_i^2) = 1^2 \times p + 0^2 \times (1 - p) = p.$$

$$\text{Var}(X_i) = p - (E(X))^2 = p - p^2 = p(1 - p).$$

$$p = 0 \implies \text{Var}(X_i) = 0$$

Variance of Binomial Distribution.

Flip coin with heads probability p .

X - how many heads?

$$X_i = \begin{cases} 1 & \text{if } i\text{th flip is heads} \\ 0 & \text{otherwise} \end{cases}$$

$$E(X_i^2) = 1^2 \times p + 0^2 \times (1 - p) = p.$$

$$\text{Var}(X_i) = p - (E(X))^2 = p - p^2 = p(1 - p).$$

$$p = 0 \implies \text{Var}(X_i) = 0$$

$$p = 1$$

Variance of Binomial Distribution.

Flip coin with heads probability p .

X - how many heads?

$$X_i = \begin{cases} 1 & \text{if } i\text{th flip is heads} \\ 0 & \text{otherwise} \end{cases}$$

$$E(X_i^2) = 1^2 \times p + 0^2 \times (1 - p) = p.$$

$$\text{Var}(X_i) = p - (E(X))^2 = p - p^2 = p(1 - p).$$

$$p = 0 \implies \text{Var}(X_i) = 0$$

$$p = 1 \implies \text{Var}(X_i) = 0$$

Variance of Binomial Distribution.

Flip coin with heads probability p .

X - how many heads?

$$X_i = \begin{cases} 1 & \text{if } i\text{th flip is heads} \\ 0 & \text{otherwise} \end{cases}$$

$$E(X_i^2) = 1^2 \times p + 0^2 \times (1 - p) = p.$$

$$\text{Var}(X_i) = p - (E(X))^2 = p - p^2 = p(1 - p).$$

$$p = 0 \implies \text{Var}(X_i) = 0$$

$$p = 1 \implies \text{Var}(X_i) = 0$$

Variance of Binomial Distribution.

Flip coin with heads probability p .

X - how many heads?

$$X_i = \begin{cases} 1 & \text{if } i\text{th flip is heads} \\ 0 & \text{otherwise} \end{cases}$$

$$E(X_i^2) = 1^2 \times p + 0^2 \times (1 - p) = p.$$

$$\text{Var}(X_i) = p - (E(X))^2 = p - p^2 = p(1 - p).$$

$$p = 0 \implies \text{Var}(X_i) = 0$$

$$p = 1 \implies \text{Var}(X_i) = 0$$

$$X = X_1 + X_2 + \dots X_n.$$

Variance of Binomial Distribution.

Flip coin with heads probability p .

X - how many heads?

$$X_i = \begin{cases} 1 & \text{if } i\text{th flip is heads} \\ 0 & \text{otherwise} \end{cases}$$

$$E(X_i^2) = 1^2 \times p + 0^2 \times (1 - p) = p.$$

$$\text{Var}(X_i) = p - (E(X))^2 = p - p^2 = p(1 - p).$$

$$p = 0 \implies \text{Var}(X_i) = 0$$

$$p = 1 \implies \text{Var}(X_i) = 0$$

$$X = X_1 + X_2 + \dots X_n.$$

X_i and X_j are independent:

Variance of Binomial Distribution.

Flip coin with heads probability p .

X - how many heads?

$$X_i = \begin{cases} 1 & \text{if } i\text{th flip is heads} \\ 0 & \text{otherwise} \end{cases}$$

$$E(X_i^2) = 1^2 \times p + 0^2 \times (1 - p) = p.$$

$$\text{Var}(X_i) = p - (E(X))^2 = p - p^2 = p(1 - p).$$

$$p = 0 \implies \text{Var}(X_i) = 0$$

$$p = 1 \implies \text{Var}(X_i) = 0$$

$$X = X_1 + X_2 + \dots X_n.$$

X_i and X_j are independent: $\text{Pr}[X_i = 1 | X_j = 1] = \text{Pr}[X_i = 1]$.

Variance of Binomial Distribution.

Flip coin with heads probability p .

X - how many heads?

$$X_i = \begin{cases} 1 & \text{if } i\text{th flip is heads} \\ 0 & \text{otherwise} \end{cases}$$

$$E(X_i^2) = 1^2 \times p + 0^2 \times (1 - p) = p.$$

$$\text{Var}(X_i) = p - (E(X))^2 = p - p^2 = p(1 - p).$$

$$p = 0 \implies \text{Var}(X_i) = 0$$

$$p = 1 \implies \text{Var}(X_i) = 0$$

$$X = X_1 + X_2 + \dots X_n.$$

X_i and X_j are independent: $\text{Pr}[X_i = 1 | X_j = 1] = \text{Pr}[X_i = 1]$.

$$\text{Var}(X) = \text{Var}(X_1 + \dots X_n)$$

Variance of Binomial Distribution.

Flip coin with heads probability p .

X - how many heads?

$$X_i = \begin{cases} 1 & \text{if } i\text{th flip is heads} \\ 0 & \text{otherwise} \end{cases}$$

$$E(X_i^2) = 1^2 \times p + 0^2 \times (1 - p) = p.$$

$$\text{Var}(X_i) = p - (E(X))^2 = p - p^2 = p(1 - p).$$

$$p = 0 \implies \text{Var}(X_i) = 0$$

$$p = 1 \implies \text{Var}(X_i) = 0$$

$$X = X_1 + X_2 + \dots X_n.$$

X_i and X_j are independent: $\text{Pr}[X_i = 1 | X_j = 1] = \text{Pr}[X_i = 1]$.

$$\text{Var}(X) = \text{Var}(X_1 + \dots X_n) = np(1 - p).$$

Variance of Binomial Distribution.

Flip coin with heads probability p .

X - how many heads?

$$X_i = \begin{cases} 1 & \text{if } i\text{th flip is heads} \\ 0 & \text{otherwise} \end{cases}$$

$$E(X_i^2) = 1^2 \times p + 0^2 \times (1 - p) = p.$$

$$\text{Var}(X_i) = p - (E(X))^2 = p - p^2 = p(1 - p).$$

$$p = 0 \implies \text{Var}(X_i) = 0$$

$$p = 1 \implies \text{Var}(X_i) = 0$$

$$X = X_1 + X_2 + \dots X_n.$$

X_i and X_j are independent: $\text{Pr}[X_i = 1 | X_j = 1] = \text{Pr}[X_i = 1]$.

$$\text{Var}(X) = \text{Var}(X_1 + \dots X_n) = np(1 - p).$$

Poisson Distribution: Variance.

Definition Poisson Distribution with parameter $\lambda > 0$

$$X = P(\lambda) \Leftrightarrow \Pr[X = m] = \frac{\lambda^m}{m!} e^{-\lambda}, m \geq 0.$$

Poisson Distribution: Variance.

Definition Poisson Distribution with parameter $\lambda > 0$

$$X = P(\lambda) \Leftrightarrow \Pr[X = m] = \frac{\lambda^m}{m!} e^{-\lambda}, m \geq 0.$$

Mean, Variance?

Poisson Distribution: Variance.

Definition Poisson Distribution with parameter $\lambda > 0$

$$X = P(\lambda) \Leftrightarrow \Pr[X = m] = \frac{\lambda^m}{m!} e^{-\lambda}, m \geq 0.$$

Mean, Variance?

Ugh.

Poisson Distribution: Variance.

Definition Poisson Distribution with parameter $\lambda > 0$

$$X = P(\lambda) \Leftrightarrow \Pr[X = m] = \frac{\lambda^m}{m!} e^{-\lambda}, m \geq 0.$$

Mean, Variance?

Ugh.

Recall that Poisson is the limit of the Binomial with $p = \lambda/n$ as $n \rightarrow \infty$.

Poisson Distribution: Variance.

Definition Poisson Distribution with parameter $\lambda > 0$

$$X = P(\lambda) \Leftrightarrow \Pr[X = m] = \frac{\lambda^m}{m!} e^{-\lambda}, m \geq 0.$$

Mean, Variance?

Ugh.

Recall that Poisson is the limit of the Binomial with $p = \lambda/n$ as $n \rightarrow \infty$.

Mean: $pn = \lambda$

Poisson Distribution: Variance.

Definition Poisson Distribution with parameter $\lambda > 0$

$$X = P(\lambda) \Leftrightarrow \Pr[X = m] = \frac{\lambda^m}{m!} e^{-\lambda}, m \geq 0.$$

Mean, Variance?

Ugh.

Recall that Poisson is the limit of the Binomial with $p = \lambda/n$ as $n \rightarrow \infty$.

Mean: $pn = \lambda$

Variance: $p(1-p)n = \lambda - \lambda^2/n \rightarrow \lambda$

Poisson Distribution: Variance.

Definition Poisson Distribution with parameter $\lambda > 0$

$$X = P(\lambda) \Leftrightarrow \Pr[X = m] = \frac{\lambda^m}{m!} e^{-\lambda}, m \geq 0.$$

Mean, Variance?

Ugh.

Recall that Poisson is the limit of the Binomial with $p = \lambda/n$ as $n \rightarrow \infty$.

Mean: $pn = \lambda$

Variance: $p(1-p)n = \lambda - \lambda^2/n \rightarrow \lambda$.

Poisson Distribution: Variance.

Definition Poisson Distribution with parameter $\lambda > 0$

$$X = P(\lambda) \Leftrightarrow \Pr[X = m] = \frac{\lambda^m}{m!} e^{-\lambda}, m \geq 0.$$

Mean, Variance?

Ugh.

Recall that Poisson is the limit of the Binomial with $p = \lambda/n$ as $n \rightarrow \infty$.

Mean: $pn = \lambda$

Variance: $p(1-p)n = \lambda - \lambda^2/n \rightarrow \lambda$.

$E(X^2)$?

Poisson Distribution: Variance.

Definition Poisson Distribution with parameter $\lambda > 0$

$$X = P(\lambda) \Leftrightarrow \Pr[X = m] = \frac{\lambda^m}{m!} e^{-\lambda}, m \geq 0.$$

Mean, Variance?

Ugh.

Recall that Poisson is the limit of the Binomial with $p = \lambda/n$ as $n \rightarrow \infty$.

Mean: $pn = \lambda$

Variance: $p(1-p)n = \lambda - \lambda^2/n \rightarrow \lambda$.

$E(X^2)$? $\text{Var}(X) = E(X^2) - (E(X))^2$ or $E(X^2) = \text{Var}(X) + E(X)^2$.

Poisson Distribution: Variance.

Definition Poisson Distribution with parameter $\lambda > 0$

$$X = P(\lambda) \Leftrightarrow \Pr[X = m] = \frac{\lambda^m}{m!} e^{-\lambda}, m \geq 0.$$

Mean, Variance?

Ugh.

Recall that Poisson is the limit of the Binomial with $p = \lambda/n$ as $n \rightarrow \infty$.

Mean: $pn = \lambda$

Variance: $p(1-p)n = \lambda - \lambda^2/n \rightarrow \lambda$.

$E(X^2)$? $\text{Var}(X) = E(X^2) - (E(X))^2$ or $E(X^2) = \text{Var}(X) + E(X)^2$.

$E(X^2) = \lambda + \lambda^2$.

Covariance

Definition The covariance of X and Y is

$$\text{cov}(X, Y) := E[(X - E[X])(Y - E[Y])].$$

Covariance

Definition The covariance of X and Y is

$$\text{cov}(X, Y) := E[(X - E[X])(Y - E[Y])].$$

Fact

$$\text{cov}(X, Y) = E[XY] - E[X]E[Y].$$

Covariance

Definition The covariance of X and Y is

$$\text{cov}(X, Y) := E[(X - E[X])(Y - E[Y])].$$

Fact

$$\text{cov}(X, Y) = E[XY] - E[X]E[Y].$$

Proof:

Think about $E[X] = E[Y] = 0$. Just $E[XY]$.

Covariance

Definition The covariance of X and Y is

$$\text{cov}(X, Y) := E[(X - E[X])(Y - E[Y])].$$

Fact

$$\text{cov}(X, Y) = E[XY] - E[X]E[Y].$$

Proof:

Think about $E[X] = E[Y] = 0$. Just $E[XY]$.



Covariance

Definition The covariance of X and Y is

$$\text{cov}(X, Y) := E[(X - E[X])(Y - E[Y])].$$

Fact

$$\text{cov}(X, Y) = E[XY] - E[X]E[Y].$$

Proof:

Think about $E[X] = E[Y] = 0$. Just $E[XY]$.

□ish.

Covariance

Definition The covariance of X and Y is

$$\text{cov}(X, Y) := E[(X - E[X])(Y - E[Y])].$$

Fact

$$\text{cov}(X, Y) = E[XY] - E[X]E[Y].$$

Proof:

Think about $E[X] = E[Y] = 0$. Just $E[XY]$.

□ish.

For the sake of completeness.

Covariance

Definition The covariance of X and Y is

$$\text{cov}(X, Y) := E[(X - E[X])(Y - E[Y])].$$

Fact

$$\text{cov}(X, Y) = E[XY] - E[X]E[Y].$$

Proof:

Think about $E[X] = E[Y] = 0$. Just $E[XY]$.

□ish.

For the sake of completeness.

$$\begin{aligned} E[(X - E[X])(Y - E[Y])] &= E[XY - E[X]Y - XE[Y] + E[X]E[Y]] \\ &= E[XY] - E[X]E[Y] - E[X]E[Y] + E[X]E[Y] \\ &= E[XY] - E[X]E[Y]. \end{aligned}$$

Correlation

Definition The correlation of X, Y , $Cor(X, Y)$ is

$$corr(X, Y) : \frac{cov(X, Y)}{\sigma(X)\sigma(Y)}.$$

Theorem: $-1 \leq corr(X, Y) \leq 1$.

Proof: Idea: $(a - b)^2 > 0$

Correlation

Definition The correlation of X, Y , $Cor(X, Y)$ is

$$corr(X, Y) : \frac{cov(X, Y)}{\sigma(X)\sigma(Y)}.$$

Theorem: $-1 \leq corr(X, Y) \leq 1$.

Proof: Idea: $(a - b)^2 > 0 \rightarrow a^2 + b^2 \geq 2ab$.

Correlation

Definition The correlation of X, Y , $Cor(X, Y)$ is

$$corr(X, Y) : \frac{cov(X, Y)}{\sigma(X)\sigma(Y)}.$$

Theorem: $-1 \leq corr(X, Y) \leq 1$.

Proof: Idea: $(a - b)^2 > 0 \rightarrow a^2 + b^2 \geq 2ab$.

Simple case: $E[X] = E[Y] = 0$ and $E[X^2] = E[Y^2] = 1$.

Correlation

Definition The correlation of X, Y , $Cor(X, Y)$ is

$$corr(X, Y) : \frac{cov(X, Y)}{\sigma(X)\sigma(Y)}.$$

Theorem: $-1 \leq corr(X, Y) \leq 1$.

Proof: Idea: $(a - b)^2 > 0 \rightarrow a^2 + b^2 \geq 2ab$.

Simple case: $E[X] = E[Y] = 0$ and $E[X^2] = E[Y^2] = 1$.

$Cor(X, Y) = E[XY]$.

Correlation

Definition The correlation of X, Y , $Cor(X, Y)$ is

$$corr(X, Y) : \frac{cov(X, Y)}{\sigma(X)\sigma(Y)}.$$

Theorem: $-1 \leq corr(X, Y) \leq 1$.

Proof: Idea: $(a - b)^2 > 0 \rightarrow a^2 + b^2 \geq 2ab$.

Simple case: $E[X] = E[Y] = 0$ and $E[X^2] = E[Y^2] = 1$.

$Cor(X, Y) = E[XY]$.

$$E[(X - Y)^2] = E[X^2] + E[Y^2] - 2E[XY] = 2(1 - E[XY]) \geq 0$$

Correlation

Definition The correlation of X, Y , $Cor(X, Y)$ is

$$corr(X, Y) : \frac{cov(X, Y)}{\sigma(X)\sigma(Y)}.$$

Theorem: $-1 \leq corr(X, Y) \leq 1$.

Proof: Idea: $(a - b)^2 > 0 \rightarrow a^2 + b^2 \geq 2ab$.

Simple case: $E[X] = E[Y] = 0$ and $E[X^2] = E[Y^2] = 1$.

$Cor(X, Y) = E[XY]$.

$E[(X - Y)^2] = E[X^2] + E[Y^2] - 2E[XY] = 2(1 - E[XY]) \geq 0$
 $\rightarrow E[XY] \leq 1$.

Correlation

Definition The correlation of X, Y , $Cor(X, Y)$ is

$$corr(X, Y) : \frac{cov(X, Y)}{\sigma(X)\sigma(Y)}.$$

Theorem: $-1 \leq corr(X, Y) \leq 1$.

Proof: Idea: $(a - b)^2 > 0 \rightarrow a^2 + b^2 \geq 2ab$.

Simple case: $E[X] = E[Y] = 0$ and $E[X^2] = E[Y^2] = 1$.

$Cor(X, Y) = E[XY]$.

$E[(X - Y)^2] = E[X^2] + E[Y^2] - 2E[XY] = 2(1 - E[XY]) \geq 0$
 $\rightarrow E[XY] \leq 1$.

$E[(X + Y)^2] = E[X^2] + E[Y^2] + 2E[XY] = 2(1 + E[XY]) \geq 0$

Correlation

Definition The correlation of X, Y , $Cor(X, Y)$ is

$$corr(X, Y) : \frac{cov(X, Y)}{\sigma(X)\sigma(Y)}.$$

Theorem: $-1 \leq corr(X, Y) \leq 1$.

Proof: Idea: $(a - b)^2 > 0 \rightarrow a^2 + b^2 \geq 2ab$.

Simple case: $E[X] = E[Y] = 0$ and $E[X^2] = E[Y^2] = 1$.

$$Cor(X, Y) = E[XY].$$

$$\begin{aligned} E[(X - Y)^2] &= E[X^2] + E[Y^2] - 2E[XY] = 2(1 - E[XY]) \geq 0 \\ \rightarrow E[XY] &\leq 1. \end{aligned}$$


$$\begin{aligned} E[(X + Y)^2] &= E[X^2] + E[Y^2] + 2E[XY] = 2(1 + E[XY]) \geq 0 \\ \rightarrow E[XY] &\geq -1. \end{aligned}$$



Shifting and scaling doesn't change correlation.


Examples of Covariance

Four equally likely pairs of values




Examples of Covariance

Four equally likely pairs of values



Note that $E[X] = 0$ and $E[Y] = 0$ in these examples. Then $cov(X, Y) = E[XY]$.


Examples of Covariance



Note that $E[X] = 0$ and $E[Y] = 0$ in these examples. Then $cov(X, Y) = E[XY]$.

When $cov(X, Y) > 0$, the RVs X and Y tend to be large or small together.


Examples of Covariance



Note that $E[X] = 0$ and $E[Y] = 0$ in these examples. Then $cov(X, Y) = E[XY]$.

When $cov(X, Y) > 0$, the RVs X and Y tend to be large or small together. X and Y are said to be **positively correlated**.

Examples of Covariance




Note that $E[X] = 0$ and $E[Y] = 0$ in these examples. Then $cov(X, Y) = E[XY]$.

When $cov(X, Y) > 0$, the RVs X and Y tend to be large or small together. X and Y are said to be **positively correlated**.

When $cov(X, Y) < 0$, when X is larger, Y tends to be smaller.

Examples of Covariance




Note that $E[X] = 0$ and $E[Y] = 0$ in these examples. Then $cov(X, Y) = E[XY]$.

When $cov(X, Y) > 0$, the RVs X and Y tend to be large or small together. X and Y are said to be **positively correlated**.

When $cov(X, Y) < 0$, when X is larger, Y tends to be smaller. X and Y are said to be **negatively correlated**.

Examples of Covariance




Note that $E[X] = 0$ and $E[Y] = 0$ in these examples. Then $cov(X, Y) = E[XY]$.

When $cov(X, Y) > 0$, the RVs X and Y tend to be large or small together. X and Y are said to be **positively correlated**.


When $cov(X, Y) < 0$, when X is larger, Y tends to be smaller. X and Y are said to be **negatively correlated**.

When $cov(X, Y) = 0$, we say that X and Y are **uncorrelated**.

Examples of Covariance




Examples of Covariance



$$E[X] = 1 \times 0.15 + 2 \times 0.4 + 3 \times 0.45 = 2.3$$


Examples of Covariance



$$E[X] = 1 \times 0.15 + 2 \times 0.4 + 3 \times 0.45 = 2.3$$

$$E[X^2] = 1^2 \times 0.15 + 2^2 \times 0.4 + 3^2 \times 0.45 = 5.8$$

Examples of Covariance




$$E[X] = 1 \times 0.15 + 2 \times 0.4 + 3 \times 0.45 = 2.3$$

$$E[X^2] = 1^2 \times 0.15 + 2^2 \times 0.4 + 3^2 \times 0.45 = 5.8$$

$$E[Y] = 1 \times 0.2 + 2 \times 0.6 + 3 \times 0.2 = 2$$

Examples of Covariance




$$E[X] = 1 \times 0.15 + 2 \times 0.4 + 3 \times 0.45 = 2.3$$

$$E[X^2] = 1^2 \times 0.15 + 2^2 \times 0.4 + 3^2 \times 0.45 = 5.8$$

$$E[Y] = 1 \times 0.2 + 2 \times 0.6 + 3 \times 0.2 = 2$$

$$E[Y^2] = 1 \times 0.2 + 4 \times 0.6 + 9 \times 0.2 = 4.4$$

Examples of Covariance



$$E[X] = 1 \times 0.15 + 2 \times 0.4 + 3 \times 0.45 = 2.3$$


$$E[X^2] = 1^2 \times 0.15 + 2^2 \times 0.4 + 3^2 \times 0.45 = 5.8$$

$$E[Y] = 1 \times 0.2 + 2 \times 0.6 + 3 \times 0.2 = 2$$

$$E[Y^2] = 1 \times 0.2 + 4 \times 0.6 + 9 \times 0.2 = 4.4$$

$$E[XY] = 1 \times 0.05 + 1 \times 2 \times 0.1 + \cdots + 3 \times 3 \times 0.2 = 4.85$$

Examples of Covariance



$$E[X] = 1 \times 0.15 + 2 \times 0.4 + 3 \times 0.45 = 2.3$$

$$E[X^2] = 1^2 \times 0.15 + 2^2 \times 0.4 + 3^2 \times 0.45 = 5.8$$


$$E[Y] = 1 \times 0.2 + 2 \times 0.6 + 3 \times 0.2 = 2$$

$$E[Y^2] = 1 \times 0.2 + 4 \times 0.6 + 9 \times 0.2 = 4.4$$

$$E[XY] = 1 \times 0.05 + 1 \times 2 \times 0.1 + \cdots + 3 \times 3 \times 0.2 = 4.85$$

$$\text{cov}(X, Y) = E[XY] - E[X]E[Y] = .25$$

Examples of Covariance



$$E[X] = 1 \times 0.15 + 2 \times 0.4 + 3 \times 0.45 = 2.3$$

$$E[X^2] = 1^2 \times 0.15 + 2^2 \times 0.4 + 3^2 \times 0.45 = 5.8$$

$$E[Y] = 1 \times 0.2 + 2 \times 0.6 + 3 \times 0.2 = 2$$


$$E[Y^2] = 1 \times 0.2 + 4 \times 0.6 + 9 \times 0.2 = 4.4$$

$$E[XY] = 1 \times 0.05 + 1 \times 2 \times 0.1 + \dots + 3 \times 3 \times 0.2 = 4.85$$

$$\text{cov}(X, Y) = E[XY] - E[X]E[Y] = .25$$

$$\text{var}[X] = E[X^2] - E[X]^2 = .51$$

Examples of Covariance



$$E[X] = 1 \times 0.15 + 2 \times 0.4 + 3 \times 0.45 = 2.3$$

$$E[X^2] = 1^2 \times 0.15 + 2^2 \times 0.4 + 3^2 \times 0.45 = 5.8$$

$$E[Y] = 1 \times 0.2 + 2 \times 0.6 + 3 \times 0.2 = 2$$

$$E[Y^2] = 1 \times 0.2 + 4 \times 0.6 + 9 \times 0.2 = 4.4$$


$$E[XY] = 1 \times 0.05 + 1 \times 2 \times 0.1 + \cdots + 3 \times 3 \times 0.2 = 4.85$$

$$\text{cov}(X, Y) = E[XY] - E[X]E[Y] = .25$$

$$\text{var}[X] = E[X^2] - E[X]^2 = .51$$

$$\text{var}[Y] = E[Y^2] - E[Y]^2 = .4$$

Examples of Covariance



$$E[X] = 1 \times 0.15 + 2 \times 0.4 + 3 \times 0.45 = 2.3$$

$$E[X^2] = 1^2 \times 0.15 + 2^2 \times 0.4 + 3^2 \times 0.45 = 5.8$$

$$E[Y] = 1 \times 0.2 + 2 \times 0.6 + 3 \times 0.2 = 2$$

$$E[Y^2] = 1 \times 0.2 + 4 \times 0.6 + 9 \times 0.2 = 4.4$$

$$E[XY] = 1 \times 0.05 + 1 \times 2 \times 0.1 + \dots + 3 \times 3 \times 0.2 = 4.85$$


$$\text{cov}(X, Y) = E[XY] - E[X]E[Y] = .25$$

$$\text{var}[X] = E[X^2] - E[X]^2 = .51$$

$$\text{var}[Y] = E[Y^2] - E[Y]^2 = .4$$

$$\text{corr}(X, Y) \approx 0.55$$

Examples of Covariance



$$E[X] = 1 \times 0.15 + 2 \times 0.4 + 3 \times 0.45 = 2.3$$

$$E[X^2] = 1^2 \times 0.15 + 2^2 \times 0.4 + 3^2 \times 0.45 = 5.8$$

$$E[Y] = 1 \times 0.2 + 2 \times 0.6 + 3 \times 0.2 = 2$$

$$E[Y^2] = 1 \times 0.2 + 4 \times 0.6 + 9 \times 0.2 = 4.4$$

$$E[XY] = 1 \times 0.05 + 1 \times 2 \times 0.1 + \dots + 3 \times 3 \times 0.2 = 4.85$$

$$\text{cov}(X, Y) = E[XY] - E[X]E[Y] = .25$$

$$\text{var}[X] = E[X^2] - E[X]^2 = .51$$

$$\text{var}[Y] = E[Y^2] - E[Y]^2 = .4$$

$$\text{corr}(X, Y) \approx 0.55$$

Properties of Covariance

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])] = E[XY] - E[X]E[Y].$$

Properties of Covariance

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])] = E[XY] - E[X]E[Y].$$

Fact

(a) $\text{var}[X] = \text{cov}(X, X)$

Properties of Covariance

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])] = E[XY] - E[X]E[Y].$$

Fact

(a) $\text{var}[X] = \text{cov}(X, X)$

(b) X, Y independent $\Rightarrow \text{cov}(X, Y) =$

Properties of Covariance

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])] = E[XY] - E[X]E[Y].$$

Fact

(a) $\text{var}[X] = \text{cov}(X, X)$

(b) X, Y independent $\Rightarrow \text{cov}(X, Y) = 0$

Properties of Covariance

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])] = E[XY] - E[X]E[Y].$$

Fact

- (a) $\text{var}[X] = \text{cov}(X, X)$
- (b) X, Y independent $\Rightarrow \text{cov}(X, Y) = 0$
- (c) $\text{cov}(a + X, b + Y) = \text{cov}(X, Y)$

Properties of Covariance

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])] = E[XY] - E[X]E[Y].$$

Fact

- (a) $\text{var}[X] = \text{cov}(X, X)$
- (b) X, Y independent $\Rightarrow \text{cov}(X, Y) = 0$
- (c) $\text{cov}(a + X, b + Y) = \text{cov}(X, Y)$
- (d) $\text{cov}(aX + bY, cU + dV) = ac \cdot \text{cov}(X, U) + ad \cdot \text{cov}(X, V) + bc \cdot \text{cov}(Y, U) + bd \cdot \text{cov}(Y, V).$

Properties of Covariance

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])] = E[XY] - E[X]E[Y].$$

Fact

- (a) $\text{var}[X] = \text{cov}(X, X)$
- (b) X, Y independent $\Rightarrow \text{cov}(X, Y) = 0$
- (c) $\text{cov}(a + X, b + Y) = \text{cov}(X, Y)$
- (d) $\text{cov}(aX + bY, cU + dV) = ac \cdot \text{cov}(X, U) + ad \cdot \text{cov}(X, V) + bc \cdot \text{cov}(Y, U) + bd \cdot \text{cov}(Y, V).$

Proof:

(a)-(b)-(c) are obvious.

Properties of Covariance

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])] = E[XY] - E[X]E[Y].$$

Fact

- (a) $\text{var}[X] = \text{cov}(X, X)$
- (b) X, Y independent $\Rightarrow \text{cov}(X, Y) = 0$
- (c) $\text{cov}(a + X, b + Y) = \text{cov}(X, Y)$
- (d) $\text{cov}(aX + bY, cU + dV) = ac \cdot \text{cov}(X, U) + ad \cdot \text{cov}(X, V) + bc \cdot \text{cov}(Y, U) + bd \cdot \text{cov}(Y, V).$

Proof:

- (a)-(b)-(c) are obvious.
- (d) In view of (c), one can subtract the means and assume that the RVs are zero-mean.

Properties of Covariance

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])] = E[XY] - E[X]E[Y].$$

Fact

- (a) $\text{var}[X] = \text{cov}(X, X)$
- (b) X, Y independent $\Rightarrow \text{cov}(X, Y) = 0$
- (c) $\text{cov}(a + X, b + Y) = \text{cov}(X, Y)$
- (d) $\text{cov}(aX + bY, cU + dV) = ac \cdot \text{cov}(X, U) + ad \cdot \text{cov}(X, V) + bc \cdot \text{cov}(Y, U) + bd \cdot \text{cov}(Y, V).$

Proof:

- (a)-(b)-(c) are obvious.
- (d) In view of (c), one can subtract the means and assume that the RVs are zero-mean. Then,

$$\text{cov}(aX + bY, cU + dV) = E[(aX + bY)(cU + dV)]$$

Properties of Covariance

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])] = E[XY] - E[X]E[Y].$$

Fact

- (a) $\text{var}[X] = \text{cov}(X, X)$
- (b) X, Y independent $\Rightarrow \text{cov}(X, Y) = 0$
- (c) $\text{cov}(a + X, b + Y) = \text{cov}(X, Y)$
- (d) $\text{cov}(aX + bY, cU + dV) = ac \cdot \text{cov}(X, U) + ad \cdot \text{cov}(X, V) + bc \cdot \text{cov}(Y, U) + bd \cdot \text{cov}(Y, V).$

Proof:

- (a)-(b)-(c) are obvious.
- (d) In view of (c), one can subtract the means and assume that the RVs are zero-mean. Then,

$$\begin{aligned}\text{cov}(aX + bY, cU + dV) &= E[(aX + bY)(cU + dV)] \\ &= ac \cdot E[XU] + ad \cdot E[XV] + bc \cdot E[YU] + bd \cdot E[YV]\end{aligned}$$

Properties of Covariance

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])] = E[XY] - E[X]E[Y].$$

Fact

- (a) $\text{var}[X] = \text{cov}(X, X)$
- (b) X, Y independent $\Rightarrow \text{cov}(X, Y) = 0$
- (c) $\text{cov}(a + X, b + Y) = \text{cov}(X, Y)$
- (d) $\text{cov}(aX + bY, cU + dV) = ac \cdot \text{cov}(X, U) + ad \cdot \text{cov}(X, V) + bc \cdot \text{cov}(Y, U) + bd \cdot \text{cov}(Y, V).$

Proof:

- (a)-(b)-(c) are obvious.
- (d) In view of (c), one can subtract the means and assume that the RVs are zero-mean. Then,

$$\begin{aligned}\text{cov}(aX + bY, cU + dV) &= E[(aX + bY)(cU + dV)] \\ &= ac \cdot E[XU] + ad \cdot E[XV] + bc \cdot E[YU] + bd \cdot E[YV] \\ &= ac \cdot \text{cov}(X, U) + ad \cdot \text{cov}(X, V) + bc \cdot \text{cov}(Y, U) + bd \cdot \text{cov}(Y, V).\end{aligned}$$

Properties of Covariance

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])] = E[XY] - E[X]E[Y].$$

Fact

- (a) $\text{var}[X] = \text{cov}(X, X)$
- (b) X, Y independent $\Rightarrow \text{cov}(X, Y) = 0$
- (c) $\text{cov}(a + X, b + Y) = \text{cov}(X, Y)$
- (d) $\text{cov}(aX + bY, cU + dV) = ac \cdot \text{cov}(X, U) + ad \cdot \text{cov}(X, V) + bc \cdot \text{cov}(Y, U) + bd \cdot \text{cov}(Y, V).$

Proof:

- (a)-(b)-(c) are obvious.
- (d) In view of (c), one can subtract the means and assume that the RVs are zero-mean. Then,

$$\begin{aligned}\text{cov}(aX + bY, cU + dV) &= E[(aX + bY)(cU + dV)] \\ &= ac \cdot E[XU] + ad \cdot E[XV] + bc \cdot E[YU] + bd \cdot E[YV] \\ &= ac \cdot \text{cov}(X, U) + ad \cdot \text{cov}(X, V) + bc \cdot \text{cov}(Y, U) + bd \cdot \text{cov}(Y, V).\end{aligned}$$



Summary

Variance

Summary

Variance

- **Variance:** $\text{var}[X] := E[(X - E[X])^2] = E[X^2] - E[X]^2$

Summary

Variance

- ▶ **Variance:** $\text{var}[X] := E[(X - E[X])^2] = E[X^2] - E[X]^2$
- ▶ **Fact:** $\text{var}[aX + b] = a^2 \text{var}[X]$

Summary

Variance

- ▶ **Variance:** $\text{var}[X] := E[(X - E[X])^2] = E[X^2] - E[X]^2$
- ▶ **Fact:** $\text{var}[aX + b] = a^2 \text{var}[X]$
- ▶ **Sum:** X, Y, Z pairwise ind. $\Rightarrow \text{var}[X + Y + Z] = \dots$

Summary

Variance

- ▶ **Variance:** $\text{var}[X] := E[(X - E[X])^2] = E[X^2] - E[X]^2$
- ▶ **Fact:** $\text{var}[aX + b] = a^2 \text{var}[X]$
- ▶ **Sum:** X, Y, Z pairwise ind. $\Rightarrow \text{var}[X + Y + Z] = \dots$

Random Variables so far.

Probability Space: Ω , $Pr : \Omega \rightarrow [0, 1]$, $\sum_{\omega \in \Omega} Pr(\omega) = 1$.

Random Variables so far.

Probability Space: Ω , $Pr : \Omega \rightarrow [0, 1]$, $\sum_{\omega \in \Omega} Pr(\omega) = 1$.

Random Variables: $X : \Omega \rightarrow \mathcal{R}$.

Random Variables so far.

Probability Space: Ω , $Pr : \Omega \rightarrow [0, 1]$, $\sum_{\omega \in \Omega} Pr(\omega) = 1$.

Random Variables: $X : \Omega \rightarrow R$.

Associated event: $Pr[X = a] = \sum_{\omega: X(\omega)=a} Pr(\omega)$

Random Variables so far.

Probability Space: Ω , $Pr : \Omega \rightarrow [0, 1]$, $\sum_{\omega \in \Omega} Pr(\omega) = 1$.

Random Variables: $X : \Omega \rightarrow R$.

Associated event: $Pr[X = a] = \sum_{\omega: X(\omega)=a} Pr(\omega)$

X and Y independent \iff all associated events are independent.

Random Variables so far.

Probability Space: Ω , $Pr : \Omega \rightarrow [0, 1]$, $\sum_{\omega \in \Omega} Pr(\omega) = 1$.

Random Variables: $X : \Omega \rightarrow R$.

Associated event: $Pr[X = a] = \sum_{\omega: X(\omega)=a} Pr(\omega)$

X and Y independent \iff all associated events are independent.

Expectation: $E[X] = \sum_a a Pr[X = a] = \sum_{\omega \in \Omega} X(\omega) Pr(\omega)$.

Random Variables so far.

Probability Space: Ω , $Pr : \Omega \rightarrow [0, 1]$, $\sum_{\omega \in \Omega} Pr(\omega) = 1$.

Random Variables: $X : \Omega \rightarrow R$.

Associated event: $Pr[X = a] = \sum_{\omega: X(\omega)=a} Pr(\omega)$

X and Y independent \iff all associated events are independent.

Expectation: $E[X] = \sum_a a Pr[X = a] = \sum_{\omega \in \Omega} X(\omega) Pr(\omega)$.

Linearity: $E[X + Y] = E[X] + E[Y]$.

Random Variables so far.

Probability Space: Ω , $Pr : \Omega \rightarrow [0, 1]$, $\sum_{\omega \in \Omega} Pr(\omega) = 1$.

Random Variables: $X : \Omega \rightarrow R$.

Associated event: $Pr[X = a] = \sum_{\omega: X(\omega)=a} Pr(\omega)$

X and Y independent \iff all associated events are independent.

Expectation: $E[X] = \sum_a a Pr[X = a] = \sum_{\omega \in \Omega} X(\omega) Pr(\omega)$.

Linearity: $E[X + Y] = E[X] + E[Y]$.

Variance: $Var(X) = E[(X - E[X])^2] = E[X^2] - (E(X))^2$

Random Variables so far.

Probability Space: Ω , $Pr : \Omega \rightarrow [0, 1]$, $\sum_{\omega \in \Omega} Pr(\omega) = 1$.

Random Variables: $X : \Omega \rightarrow R$.

Associated event: $Pr[X = a] = \sum_{\omega: X(\omega)=a} Pr(\omega)$

X and Y independent \iff all associated events are independent.

Expectation: $E[X] = \sum_a a Pr[X = a] = \sum_{\omega \in \Omega} X(\omega) Pr(\omega)$.

Linearity: $E[X + Y] = E[X] + E[Y]$.

Variance: $Var(X) = E[(X - E[X])^2] = E[X^2] - (E(X))^2$

For independent X, Y , $Var(X + Y) = Var(X) + Var(Y)$.

Random Variables so far.

Probability Space: Ω , $Pr : \Omega \rightarrow [0, 1]$, $\sum_{\omega \in \Omega} Pr(\omega) = 1$.

Random Variables: $X : \Omega \rightarrow R$.

Associated event: $Pr[X = a] = \sum_{\omega: X(\omega)=a} Pr(\omega)$

X and Y independent \iff all associated events are independent.

Expectation: $E[X] = \sum_a a Pr[X = a] = \sum_{\omega \in \Omega} X(\omega) Pr(\omega)$.

Linearity: $E[X + Y] = E[X] + E[Y]$.

Variance: $Var(X) = E[(X - E[X])^2] = E[X^2] - (E(X))^2$

For independent X, Y , $Var(X + Y) = Var(X) + Var(Y)$.

Also: $Var(cX) = c^2 Var(X)$ and $Var(X + b) = Var(X)$.

Random Variables so far.

Probability Space: Ω , $Pr : \Omega \rightarrow [0, 1]$, $\sum_{\omega \in \Omega} Pr(\omega) = 1$.

Random Variables: $X : \Omega \rightarrow R$.

Associated event: $Pr[X = a] = \sum_{\omega: X(\omega)=a} Pr(\omega)$

X and Y independent \iff all associated events are independent.

Expectation: $E[X] = \sum_a a Pr[X = a] = \sum_{\omega \in \Omega} X(\omega) Pr(\omega)$.

Linearity: $E[X + Y] = E[X] + E[Y]$.

Variance: $Var(X) = E[(X - E[X])^2] = E[X^2] - (E(X))^2$

For independent X, Y , $Var(X + Y) = Var(X) + Var(Y)$.

Also: $Var(cX) = c^2 Var(X)$ and $Var(X + b) = Var(X)$.

Poisson: $X \sim P(\lambda)$

Random Variables so far.

Probability Space: Ω , $Pr : \Omega \rightarrow [0, 1]$, $\sum_{\omega \in \Omega} Pr(\omega) = 1$.

Random Variables: $X : \Omega \rightarrow R$.

Associated event: $Pr[X = a] = \sum_{\omega: X(\omega)=a} Pr(\omega)$

X and Y independent \iff all associated events are independent.

Expectation: $E[X] = \sum_a a Pr[X = a] = \sum_{\omega \in \Omega} X(\omega) Pr(\omega)$.

Linearity: $E[X + Y] = E[X] + E[Y]$.

Variance: $Var(X) = E[(X - E[X])^2] = E[X^2] - (E(X))^2$

For independent X, Y , $Var(X + Y) = Var(X) + Var(Y)$.

Also: $Var(cX) = c^2 Var(X)$ and $Var(X + b) = Var(X)$.

Poisson: $X \sim P(\lambda)$ $E(X) = \lambda$, $Var(X) = \lambda$.

Random Variables so far.

Probability Space: Ω , $Pr : \Omega \rightarrow [0, 1]$, $\sum_{\omega \in \Omega} Pr(\omega) = 1$.

Random Variables: $X : \Omega \rightarrow R$.

Associated event: $Pr[X = a] = \sum_{\omega: X(\omega)=a} Pr(\omega)$

X and Y independent \iff all associated events are independent.

Expectation: $E[X] = \sum_a a Pr[X = a] = \sum_{\omega \in \Omega} X(\omega) Pr(\omega)$.

Linearity: $E[X + Y] = E[X] + E[Y]$.

Variance: $Var(X) = E[(X - E[X])^2] = E[X^2] - (E(X))^2$

For independent X, Y , $Var(X + Y) = Var(X) + Var(Y)$.

Also: $Var(cX) = c^2 Var(X)$ and $Var(X + b) = Var(X)$.

Poisson: $X \sim P(\lambda)$ $E(X) = \lambda$, $Var(X) = \lambda$.

Binomial: $X \sim B(n, p)$

Random Variables so far.

Probability Space: Ω , $Pr : \Omega \rightarrow [0, 1]$, $\sum_{\omega \in \Omega} Pr(\omega) = 1$.

Random Variables: $X : \Omega \rightarrow R$.

Associated event: $Pr[X = a] = \sum_{\omega: X(\omega)=a} Pr(\omega)$

X and Y independent \iff all associated events are independent.

Expectation: $E[X] = \sum_a a Pr[X = a] = \sum_{\omega \in \Omega} X(\omega) Pr(\omega)$.

Linearity: $E[X + Y] = E[X] + E[Y]$.

Variance: $Var(X) = E[(X - E[X])^2] = E[X^2] - (E(X))^2$

For independent X, Y , $Var(X + Y) = Var(X) + Var(Y)$.

Also: $Var(cX) = c^2 Var(X)$ and $Var(X + b) = Var(X)$.

Poisson: $X \sim P(\lambda)$ $E(X) = \lambda$, $Var(X) = \lambda$.

Binomial: $X \sim B(n, p)$ $E(X) = np$, $Var(X) = np(1 - p)$

Random Variables so far.

Probability Space: Ω , $Pr : \Omega \rightarrow [0, 1]$, $\sum_{\omega \in \Omega} Pr(\omega) = 1$.

Random Variables: $X : \Omega \rightarrow R$.

Associated event: $Pr[X = a] = \sum_{\omega: X(\omega)=a} Pr(\omega)$

X and Y independent \iff all associated events are independent.

Expectation: $E[X] = \sum_a a Pr[X = a] = \sum_{\omega \in \Omega} X(\omega) Pr(\omega)$.

Linearity: $E[X + Y] = E[X] + E[Y]$.

Variance: $Var(X) = E[(X - E[X])^2] = E[X^2] - (E(X))^2$

For independent X, Y , $Var(X + Y) = Var(X) + Var(Y)$.

Also: $Var(cX) = c^2 Var(X)$ and $Var(X + b) = Var(X)$.

Poisson: $X \sim P(\lambda)$ $E(X) = \lambda$, $Var(X) = \lambda$.

Binomial: $X \sim B(n, p)$ $E(X) = np$, $Var(X) = np(1 - p)$

Uniform: $X \sim U\{1, \dots, n\}$

Random Variables so far.

Probability Space: Ω , $Pr : \Omega \rightarrow [0, 1]$, $\sum_{\omega \in \Omega} Pr(\omega) = 1$.

Random Variables: $X : \Omega \rightarrow R$.

Associated event: $Pr[X = a] = \sum_{\omega: X(\omega)=a} Pr(\omega)$

X and Y independent \iff all associated events are independent.

Expectation: $E[X] = \sum_a a Pr[X = a] = \sum_{\omega \in \Omega} X(\omega) Pr(\omega)$.

Linearity: $E[X + Y] = E[X] + E[Y]$.

Variance: $Var(X) = E[(X - E[X])^2] = E[X^2] - (E(X))^2$

For independent X, Y , $Var(X + Y) = Var(X) + Var(Y)$.

Also: $Var(cX) = c^2 Var(X)$ and $Var(X + b) = Var(X)$.

Poisson: $X \sim P(\lambda)$ $E(X) = \lambda$, $Var(X) = \lambda$.

Binomial: $X \sim B(n, p)$ $E(X) = np$, $Var(X) = np(1 - p)$

Uniform: $X \sim U\{1, \dots, n\}$ $E[X] = \frac{n+1}{2}$, $Var(X) = \frac{n^2-1}{12}$.

Random Variables so far.

Probability Space: Ω , $Pr : \Omega \rightarrow [0, 1]$, $\sum_{\omega \in \Omega} Pr(\omega) = 1$.

Random Variables: $X : \Omega \rightarrow R$.

Associated event: $Pr[X = a] = \sum_{\omega: X(\omega)=a} Pr(\omega)$

X and Y independent \iff all associated events are independent.

Expectation: $E[X] = \sum_a a Pr[X = a] = \sum_{\omega \in \Omega} X(\omega) Pr(\omega)$.

Linearity: $E[X + Y] = E[X] + E[Y]$.

Variance: $Var(X) = E[(X - E[X])^2] = E[X^2] - (E(X))^2$

For independent X, Y , $Var(X + Y) = Var(X) + Var(Y)$.

Also: $Var(cX) = c^2 Var(X)$ and $Var(X + b) = Var(X)$.

Poisson: $X \sim P(\lambda)$ $E(X) = \lambda$, $Var(X) = \lambda$.

Binomial: $X \sim B(n, p)$ $E(X) = np$, $Var(X) = np(1 - p)$

Uniform: $X \sim U\{1, \dots, n\}$ $E[X] = \frac{n+1}{2}$, $Var(X) = \frac{n^2-1}{12}$.

Geometric: $X \sim G(p)$

Random Variables so far.

Probability Space: Ω , $Pr : \Omega \rightarrow [0, 1]$, $\sum_{\omega \in \Omega} Pr(\omega) = 1$.

Random Variables: $X : \Omega \rightarrow R$.

Associated event: $Pr[X = a] = \sum_{\omega: X(\omega)=a} Pr(\omega)$

X and Y independent \iff all associated events are independent.

Expectation: $E[X] = \sum_a a Pr[X = a] = \sum_{\omega \in \Omega} X(\omega) Pr(\omega)$.

Linearity: $E[X + Y] = E[X] + E[Y]$.

Variance: $Var(X) = E[(X - E[X])^2] = E[X^2] - (E(X))^2$

For independent X, Y , $Var(X + Y) = Var(X) + Var(Y)$.

Also: $Var(cX) = c^2 Var(X)$ and $Var(X + b) = Var(X)$.

Poisson: $X \sim P(\lambda)$ $E(X) = \lambda$, $Var(X) = \lambda$.

Binomial: $X \sim B(n, p)$ $E(X) = np$, $Var(X) = np(1 - p)$

Uniform: $X \sim U\{1, \dots, n\}$ $E[X] = \frac{n+1}{2}$, $Var(X) = \frac{n^2-1}{12}$.

Geometric: $X \sim G(p)$ $E(X) = \frac{1}{p}$, $Var(X) = \frac{1-p}{p^2}$