

Modern Physics

A Collection of Notes and Problems

Tarang Srivastava

Dr. Mesut Çakir
Modern Physics
South Brunswick High School
20 April 2019

Contents

1	Lies From Classical Physics	2
1.1	Mechanics	2
1.2	Electricity and Magnetism	2
1.3	Kinetic Theory of Matter	3
1.4	Failures of Classical Concepts of Space and Time	3
1.4.1	Failure in Concept of Time	3
1.4.2	Failure in Concept of Space	3
1.4.3	Failure in Concept of Velocity	3

Chapter 1

Lies From Classical Physics

1.1 Mechanics

kinetic energy is defined by

$$K = \frac{1}{2}mv^2$$

and *linear momentum* \vec{p} defined by

$$\vec{p} = m\vec{v}$$

In terms of the linear momentum, the kinetic energy can be written

$$K = \frac{p^2}{2m}$$

Corresponding to external force there is often a potential energy U such that (for one dimensional motion)

$$F = -\frac{dU}{dx}$$

The total energy

$$E = K + U$$

When a particle moving with linear momentum \vec{p} is at a displacement \vec{r} from the origin O , its *angular momentum* \vec{L} about the point O is defined by

$$\vec{L} = \vec{r} \times \vec{p}$$

1.2 Electricity and Magnetism

The electrostatic force exerted by a charged particle q_1 on another charge q_2 has magnitude

$$F = \frac{1}{4\pi\epsilon} \frac{|q_1||q_2|}{r^2}$$

Using the previous relationship between potential energy and external force

$$U = \frac{1}{4\pi\epsilon} \frac{q_1q_2}{r}$$

Electrostatic potential difference ΔV is given by

$$\Delta U = q\Delta V$$

For reference

$$1eV = 1.602 \times 10^{-19} J$$

A magnetic field \vec{B} can be produced by current i . The magnetic field through a loop of radius r .

$$B = \frac{\mu_0 i}{2r}$$

magnetic moment $\vec{\mu}$ of a current loop:

$$|\vec{\mu}| = iA$$

where A is the geometrical area enclosed by the loop.

When a current loop is placed in a uniform external magnetic field \vec{B}_{ext} there is a torque $\vec{\tau}$

$$\vec{\tau} = \vec{\mu} \times \vec{B}_{ext}$$

potential energy is given by

$$U = -\vec{\mu} \cdot \vec{B}_{ext}$$

Electromagnetic waves travel in free space with speed c (speed of light), which is related to the electromagnetic constants ϵ_0 and μ_0

$$c = (\epsilon_0 \mu_0)^{-1/2}$$

EM waves with frequency f and wavelength λ are related

$$c = \lambda f$$

1.3 Kinetic Theory of Matter

ideal gas equation

$$PV = NrT$$

The average kinetic energy of a molecule depends on its temperature

$$K_{av} = \frac{3}{2}kT$$

1.4 Failures of Classical Concepts of Space and Time

1.4.1 Failure in Concept of Time

Newton's laws are based on the assumption that time is the same, irrespective of the observer. This is shown to be incorrect.

1.4.2 Failure in Concept of Space

Galileo and Newton's theories are based on the assumption that space is the same, but this is not the case.

1.4.3 Failure in Concept of Velocity

If space and time are not the same no matter what the same can be said for velocity, since velocity is dependent on space and time. There is also a speed limit on the universe we call the *speed of light* c